

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12662 A2

(51) International Patent Classification⁷: C07K 14/00

(21) International Application Number: PCT/US00/22315

(22) International Filing Date: 14 August 2000 (14.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/149,641 17 August 1999 (17.08.1999) US
60/164,203 9 November 1999 (09.11.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US 60/149,641 (CIP)
Filed on 17 August 1999 (17.08.1999)
US 60/164,203 (CIP)
Filed on 9 November 1999 (09.11.1999)

(71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). BURFORD, Neil [GB/US]; 1308 4th Avenue, San

Francisco, CA 94122 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). PATTERSON, Chandra [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MEMBRANE ASSOCIATED PROTEINS

(57) Abstract: The invention provides human membrane associated proteins (MEMAP) and polynucleotides which identify and encode MEMAP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of MEMAP.



WO 01/12662 A2

MEMBRANE ASSOCIATED PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of membrane associated
5 proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cell
proliferative, autoimmune/inflammatory, neurological and gastrointestinal disorders.

BACKGROUND OF THE INVENTION

Eukaryotic cells are surrounded by plasma membranes which enclose the cell and maintain
10 an environment inside the cell that is distinct from its surroundings. In addition, eukaryotic
organisms are distinct from prokaryotes in possessing many intracellular organelle and vesicle
structures. Many of the metabolic reactions which distinguish eukaryotic biochemistry from
prokaryotic biochemistry take place within these structures. The plasma membrane and the
membranes surrounding organelles and vesicles are composed of phosphoglycerides, fatty acids,
15 cholesterol, phospholipids, glycolipids, proteoglycans, and proteins. These components confer
identity and functionality to the membranes with which they associate.

Integral Membrane Proteins

The majority of known integral membrane proteins are transmembrane proteins (TM) which
20 are characterized by an extracellular, a transmembrane, and an intracellular domain. TM domains are
typically comprised of 15 to 25 hydrophobic amino acids which are predicted to adopt an α -helical
conformation. TM proteins are classified as bitopic (Types I and II) and polytopic (Types III and IV)
(Singer, S.J. (1990) Annu. Rev. Cell Biol. 6:247-96). Bitopic proteins span the membrane once while
polytopic proteins contain multiple membrane-spanning segments. TM proteins that act as cell-
25 surface receptor proteins involved in signal transduction include growth and differentiation factor
receptors, and receptor-interacting proteins such as *Drosophila* pecanex and frizzled proteins, LIV-1
protein, NF2 protein, and GNS1/SUR4 eukaryotic integral membrane proteins. TM proteins also act
as transporters of ions or metabolites, such as gap junction channels (connexins) and ion channels,
and as cell anchoring proteins, such as lectins, integrins, and fibronectins. TM proteins act as vesicle
30 organelle-forming molecules, such as calveolins, or as cell recognition molecules, such as cluster of
differentiation (CD) antigens, glycoproteins, and mucins.

Many membrane proteins (MPs) contain amino acid sequence motifs that target these
proteins to specific subcellular sites. Examples of these motifs include PDZ domains, KDEL, RGD,
NGR, and GSL sequence motifs, von Willebrand factor A (vWFA) domains, and EGF-like domains.
35 RGD, NGR, and GSL motif-containing peptides have been used as drug delivery agents in cancer

treatments which target tumor vasculature (Arap, W. et al. (1998) Science, 279:377-380). Furthermore, MPs may also contain amino acid sequence motifs, such as the carbohydrate recognition domain (CRD), also known as the C-type lectin domain, that mediate interactions with extracellular or intracellular molecules.

5 Membrane proteins may also interact with and regulate the properties of the membrane lipids. Phospholipid scramblase, a type II plasma membrane protein, mediates calcium dependent movement of phospholipids (PL) between membrane leaflets. Calcium induced remodeling of plasma membrane PL plays a key role in expression of platelet anticoagulant activity and in clearance of injured or apoptotic cells (Zhou Q. et al. (1997) J. Biol. Chem. 272:18240-18244). Scott syndrome, a
10 bleeding disorder, is caused by an inherited deficiency in plasma membrane PL scramblase function (Online Mendelian Inheritance in Man (OMIM) *262890 Platelet Receptor for Factor X, Deficiency of).

Chemical modification of amino acid residue side chains alters the manner in which MPs interact with other molecules, such as phospholipid membranes. Examples of such chemical
15 modifications to amino acid residue side chains are covalent bond formation with glycosaminoglycans, oligosaccharides, phospholipids, acetyl and palmitoyl moieties, ADP-ribose, phosphate, and sulphate groups.

One function of TM proteins is to facilitate cell-cell communication. The slit proteins are extracellular matrix proteins expressed by cells at the ventral midline of the nervous system. Slit
20 proteins are ligands for the repulsive guidance receptor Robo and thus play a role in repulsive axon guidance (Brose, K. et al. (1999) Cell 96:795-806).

In some cases TM proteins serve as transporters or channels in the cell membrane. For example, the mouse transporter protein (MTP) has four transmembrane domains and resides in an intracellular membrane compartment. MTP can mediate transport of nucleosides in vitro. The role
25 of MTP in the cell may therefore be to transfer nucleosides between the cytosol and the lumen of intracellular organelles (Hogue, D. L. (1996) J. Biol. Chem. 271:9801-9808). The human stomatin-like protein (hSLP-1), expressed primarily in the brain, contains an N-terminal domain similar to the erythrocyte internal membrane protein stomatin, as well as a non-specific lipid transfer protein domain at the C-terminus. hSLP-1 is the human homologue of the C. elegans behavioral gene unc-
30 24, which is believed to be involved in lipid transfer between closely apposed membranes (Seidel, G. and Prohaska, R (1998) Gene 225:23-29).

The transmembrane 4 superfamily (TM4SF) or tetraspan family is a multigene family encoding type III integral membrane proteins (Wright, M.D. and Tomlinson, M.G. (1994) Immunol. Today 15:588-594). TM4SF is comprised of membrane proteins which traverse the cell membrane
35 four times. Members of the TM4SF include platelet and endothelial cell membrane proteins,

melanoma-associated antigens, leukocyte surface glycoproteins, colonal carcinoma antigens, tumor-associated antigens, and surface proteins of the schistosome parasites (Jankowski, S.A. (1994) *Oncogene* 9:1205-1211). Members of the TM4SF share about 25-30% amino acid sequence identity with one another.

5 A number of TM4SF members have been implicated in signal transduction, control of cell adhesion, regulation of cell growth and proliferation, including development and oncogenesis, and cell motility, including tumor cell metastasis. Expression of TM4SF proteins is associated with a variety of tumors and the level of expression may be altered when cells are growing or activated.

10 Tumor antigens are cell surface molecules that are differentially expressed in tumor cells relative to normal cells. Tumor antigens distinguish tumor cells immunologically from normal cells and provide diagnostic and therapeutic targets for human cancers (Takagi, S. et al. (1995) *Int. J. Cancer* 61: 706-715; Liu, E. et al. (1992) *Oncogene* 7: 1027-1032). For example, the biliary glycoprotein-encoding gene is a member of the human carcinoembryonic antigen family, which are important tumor markers for colorectal carcinomas (Hammarstrom, S. (1999) *Semin. Cancer Bio.* 9:67-81). Another example is the neuron and testis specific protein Ma1, a marker for paraneoplastic neuronal disorders (Dalmau, J. et al. (1999) *Brain* 122:27-39).

Other types of cell surface antigens include those identified on leukocytic cells of the immune system. These antigens have been identified using systematic, monoclonal antibody (mAb)-based "shot gun" techniques. These techniques have resulted in the production of hundreds of mAbs directed against unknown cell surface leukocytic antigens. These antigens have been grouped into "clusters of differentiation" based on common immunocytochemical localization patterns in various differentiated and undifferentiated leukocytic cell types. Antigens in a given cluster are presumed to identify a single cell surface protein and are assigned a "cluster of differentiation" or "CD" designation. Some of the genes encoding proteins identified by CD antigens have been cloned and verified by standard molecular biology techniques. CD antigens have been characterized as both transmembrane proteins and cell surface proteins anchored to the plasma membrane via covalent attachment to fatty acid-containing glycolipids such as glycosylphosphatidylinositol (GPI). (Reviewed in Barclay, A. N. et al. (1995) The Leucocyte Antigen Facts Book, Academic Press, San Diego, CA, pp. 17-20.)

30 The TM cell surface glycoprotein CD69 is an early activation antigen of T lymphocytes. CD69 is homologous to members of a supergene family of type II integral membrane proteins having C-type lectin domains. Although the precise functions of the CD-69 antigen is not known, evidence suggests that these proteins transmit mitogenic signals across the plasma membrane and are up-regulated in response to lymphocyte activation (Hamann, J. et. al. (1993) *J. Immunol.* 150:4920-4927).

Macrophages are involved in functions including clearance of senescent or apoptotic cells, cytokine production, hemopoiesis, bone resorption, antigen transport, and neuroendocrine regulation. These diverse roles are influenced by specialized macrophage plasma membrane proteins. The murine macrophage restricted C-type lectin is a type II integral membrane protein expressed exclusively in macrophages. The strong expression of this protein in bone marrow suggests a hemopoietic function, while the lectin domain suggests it may be involved in cell-cell recognition (Balch, S. G. et al. (1998) J. Biol. Chem. 273:18656-18664).

The surface of red blood cells is populated with characteristic glycoproteins, such as the major sialoglycoproteins glycophorin A and B. Red blood cells lacking either glycophorin A or B are resistant to infection with the malaria parasite *Plasmodium falciparum* (OMIM Entry 111300 Blood Group-MN Locus). White blood cells also possess characteristic surface glycoproteins, such as the plasma cell glycoprotein-1 (PC-1). PC-1 is expressed on the surface of plasma cells, which are terminally differentiated, antibody-secreting B-lymphocytes. The extracellular domain of PC-1 has nucleotide phosphodiesterase (pyrophosphatase) activity (Funakoshi, I. et al. (1992) Arch. Biochem. Biophys. 295:180-187). Phosphodiesterase activity is associated with the hydrolytic removal of nucleotide subunits from oligonucleotides. Although the precise physiological role of PC-1 is not clear, increased PC-1 phosphodiesterase activity has been correlated with insulin resistance in patients with noninsulin-dependent diabetes mellitus, with abnormalities of bone mineralization and calcification, and with defects in renal tubule function. In addition, it appears that hPC-1 and mPC-1 are members of a multigene family of transmembrane phosphodiesterases with extracellular active sites. These enzymes may play a role in regulating the concentration of pharmacologically active extracellular compounds such as adenosine or other nucleotide derivatives in a variety of tissues and cell types. (Reviewed in Goding, J. W. et al. (1998) Immunol. Rev. 161:11-26.)

25 Peripheral and Anchored Membrane Proteins

Some membrane proteins are not membrane-spanning but are attached to the plasma membrane via membrane anchors or interactions with integral membrane proteins. Membrane anchors are covalently joined to a protein post-translationally and include such moieties as prenyl, myristyl, and glycosylphosphatidyl inositol (GPI) groups. Membrane localization of peripheral and anchored proteins is important for their function in processes such as receptor-mediated signal transduction. For example, prenylation of Ras is required for its localization to the plasma membrane and for its normal and oncogenic functions in signal transduction.

The pancortins are a group of four glycoproteins which are predominantly expressed in the cerebral cortex of adult rodents. Immunological localization indicates that the pancortins are endoplasmic reticulum anchored proteins. The pancortins share a common sequence in the middle of

their structure, but have alternative sequences at both ends due to differential promoter usage and alternative splicing. Each pancortin appears to be differentially expressed and may perform different functions in the brain (Nagano, T. et al. (1998) Mol. Brain Res. 53:13-23).

The discovery of new membrane associated proteins and the polynucleotides encoding them
5 satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative, autoimmune/inflammatory, neurological and gastrointestinal disorders.

SUMMARY OF THE INVENTION

10 The invention features purified polypeptides, membrane associated proteins, referred to collectively as "MEMAP" and individually as "MEMAP-1," "MEMAP-2," "MEMAP-3," "MEMAP-4," "MEMAP-5," "MEMAP-6," "MEMAP-7," "MEMAP-8," "MEMAP-9," "MEMAP-10," "MEMAP-11," "MEMAP-12," "MEMAP-13," "MEMAP-14," "MEMAP-15," "MEMAP-16," "MEMAP-17," "MEMAP-18," "MEMAP-19," "MEMAP-20," "MEMAP-21," "MEMAP-22,"
15 "MEMAP-23," "MEMAP-24," "MEMAP-25," "MEMAP-26," "MEMAP-27," "MEMAP-28," "MEMAP-29," "MEMAP-30," "MEMAP-31," "MEMAP-32," "MEMAP-33," "MEMAP-34," "MEMAP-35," "MEMAP-36," and "MEMAP-37." In one aspect, the invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino
20 acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-37.

25 The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group
30 consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-37. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:38-74.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter
35 sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid

sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of
5 SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising an amino acid
10 sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of
15 SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a
20 polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino
25 acid sequence selected from the group consisting of SEQ ID NO:1-37.

The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID
30 NO:38-74, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide
35 sequence selected from the group consisting of a) a polynucleotide sequence selected from the group

consisting of SEQ ID NO:38-74, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional MEMAP, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence

selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional MEMAP, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional MEMAP, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, b) a naturally

occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-37. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:38-74, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, ii) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, ii) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38-74, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of the above polynucleotide sequence; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding MEMAP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of MEMAP.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding MEMAP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the

invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"MEMAP" refers to the amino acid sequences of substantially purified MEMAP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of MEMAP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of MEMAP either by directly interacting with MEMAP or by acting on components of the biological pathway in which MEMAP participates.

An "allelic variant" is an alternative form of the gene encoding MEMAP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding MEMAP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as MEMAP or a polypeptide with at least one functional characteristic of MEMAP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding MEMAP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding MEMAP. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent MEMAP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of MEMAP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring

protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of MEMAP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of MEMAP either by directly interacting with MEMAP or by acting on components of the biological pathway in which MEMAP participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind MEMAP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the

designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic MEMAP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding MEMAP or fragments of MEMAP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (PE Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

Original Residue	Conservative Substitution
Ala	Gly, Ser
Arg	His, Lys
Asn	Asp, Gln, His
Asp	Asn, Glu
Cys	Ala, Ser
Gln	Asn, Glu, His
Glu	Asp, Gln, His

	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
5	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
10	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of MEMAP or the polynucleotide encoding MEMAP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the

present embodiments.

A fragment of SEQ ID NO:38-74 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:38-74, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:38-74 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:38-74 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:38-74 and the region of SEQ ID NO:38-74 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-37 is encoded by a fragment of SEQ ID NO:38-74. A fragment of SEQ ID NO:1-37 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-37. For example, a fragment of SEQ ID NO:1-37 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-37. The precise length of a fragment of SEQ ID NO:1-37 and the region of SEQ ID NO:1-37 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full-length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full-length" polynucleotide sequence encodes a "full-length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms

is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence

5 analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST
10 programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

15 *Penalty for mismatch: -2*

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

20 *Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous
25 nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes
30 in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some
35 alignment methods take into account conservative amino acid substitutions. Such conservative

substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (Apr-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity.

Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide

sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of MEMAP which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of MEMAP which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of MEMAP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of MEMAP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an MEMAP may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of MEMAP.

"Probe" refers to nucleic acid sequences encoding MEMAP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

- 5 "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

10 Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

- 15 Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs
20 can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

- Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to
25 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from
30 megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection
35 programs may also be obtained from their respective sources and modified to meet the user's specific

needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both
5 unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

10 A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have
15 been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a
20 vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription,
25 translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

30 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic
35 acids encoding MEMAP, or fragments thereof, or MEMAP itself, may comprise a bodily fluid; an

extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with

a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants, and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook, J. et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human membrane associated proteins (MEMAP), the polynucleotides encoding MEMAP, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, autoimmune/inflammatory, neurological and

gastrointestinal disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding MEMAP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each MEMAP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each MEMAP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding MEMAP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:38-74 and to distinguish between SEQ ID NO:38-74 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express MEMAP as a fraction of total tissues expressing MEMAP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing MEMAP as a fraction of total tissues expressing MEMAP. Column 5 lists the vectors used to subclone each cDNA library. Of particular note is the expression of SEQ ID NO:41, SEQ ID NO:48, and SEQ ID NO:56 in nervous tissues, of SEQ ID NO:52, SEQ ID NO:65, and SEQ ID NO:74 in gastrointestinal tissues, and of SEQ ID NO:55 in hematopoietic/immune tissues.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding MEMAP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

SEQ ID NO:38 maps to chromosome 4 within the interval from 77.9 to 86.0 centiMorgans, to chromosome 6 within the interval from 132.7 to 144.4 centiMorgans, and to chromosome 14 within the interval from 89.4 to 103.7 centiMorgans. The interval on chromosome 4 from 77.9 to 86.0 centiMorgans also contains a gene associated with deoxycytidine kinase deficiency. The interval on chromosome 6 from 132.7 to 144.4 centiMorgans also contains genes associated with peroxisomal disorders and leukemia. The interval on chromosome 14 from 89.4 to 103.7 centiMorgans also contains genes associated with spinocerebellar ataxia and protease inhibitor deficiencies. SEQ ID NO:39 maps to chromosome 2 within the interval from 236.2 to 269.5 centiMorgans, and to the X chromosome within the interval from 94.4 to 97.4 centiMorgans. The interval on chromosome 2 from 236.2 to 269.5 centiMorgans also contains genes associated with Crigler-Najjar syndrome, Oguchi disease, and oxalosis I. The interval on the X chromosome from 94.4 to 97.4 centiMorgans also contains genes associated with Charcot-Marie tooth disease, X-linked severe combined immunodeficiency, alpha thalassemia/mental retardation syndrome, Menkes' syndrome, and choroideremia. SEQ ID NO:42 maps to chromosome 1 within the interval from 218.2 to 232.0 centiMorgans. This interval also contains genes associated with familial hypertrophic cardiomyopathy, malignant hyperthermia, and hypokalemic periodic paralysis. SEQ ID NO:44 maps to chromosome 7 within the interval from 136.4 to 145.8 centiMorgans, to chromosome 14 within the interval from 28.0 to 32.9 centiMorgans, and to chromosome 14 within the interval from 71.5 to 73.7 centiMorgans. The interval on chromosome 7 from 136.4 to 145.8 centiMorgans also contains genes associated with diphosphoglycerate mutase deficiency. SEQ ID NO:60 maps to chromosome 7 within the interval from 167.6 to 184.0 centiMorgans, and to chromosome 14 within the interval from 50.0 to 59.0 centiMorgans. SEQ ID NO:63 maps to chromosome 8 within the interval from 101.0 to 125.8 centiMorgans, and to chromosome 8 within the interval from 132.4 to 135.1 centiMorgans. SEQ ID NO:67 maps to chromosome 4 within the interval from 145.3 to 146.4 centiMorgans.

The invention also encompasses MEMAP variants. A preferred MEMAP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the MEMAP amino acid sequence, and which contains at least one functional or structural characteristic of MEMAP.

The invention also encompasses polynucleotides which encode MEMAP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:38-74, which encodes MEMAP. The polynucleotide sequences of SEQ ID NO:38-74, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding MEMAP.

In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding MEMAP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID
5 NO:38-74 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:38-74. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of MEMAP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the
10 genetic code, a multitude of polynucleotide sequences encoding MEMAP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the
15 polynucleotide sequence of naturally occurring MEMAP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode MEMAP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring MEMAP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding
20 MEMAP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding MEMAP and its derivatives without altering the encoded amino acid
25 sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode MEMAP and MEMAP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell
30 systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding MEMAP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:38-74 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and
35 S.L. Berger (1987) Methods Enzymol. 152:399-407; Kimmel, A.R. (1987) Methods Enzymol.

152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (PE Biosystems, Foster City CA), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (PE Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (PE Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding MEMAP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National

Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been
5 size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze
10 the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, PE Biosystems), and the entire
15 process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode MEMAP may be cloned in recombinant DNA molecules that direct expression of
20 MEMAP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express MEMAP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter MEMAP-encoding sequences for a variety of purposes including,
25 but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

30 The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of MEMAP, such as its biological or enzymatic activity or its
35 ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of

gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding MEMAP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser. 7*:215-223; Horn, T. et al. (1980) *Nucleic Acids Symp. Ser. 7*:225-232.) Alternatively, MEMAP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (PE Biosystems). Additionally, the amino acid sequence of MEMAP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active MEMAP, the nucleotide sequences encoding MEMAP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding MEMAP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding MEMAP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding MEMAP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no

additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic.

- 5 The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding MEMAP and appropriate transcriptional and translational
10 control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

- 15 A variety of expression vector/host systems may be utilized to contain and express sequences encoding MEMAP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower
20 mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-
25 1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses,
30 adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.)
- 35 The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding MEMAP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding MEMAP can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding MEMAP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of MEMAP are needed, e.g. for the production of antibodies, vectors which direct high level expression of MEMAP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of MEMAP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, *supra*; and Scorer, *supra*.)

Plant systems may also be used for expression of MEMAP. Transcription of sequences encoding MEMAP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, *supra*; Broglie, *supra*; and Winter, *supra*.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding MEMAP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses MEMAP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of

DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

5 For long term production of recombinant proteins in mammalian systems, stable expression of MEMAP in cell lines is preferred. For example, sequences encoding MEMAP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in
10 enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These
15 include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk*⁻ and *apr*⁻ cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat*
20 confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins
25 (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is
30 also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding MEMAP is inserted within a marker gene sequence, transformed cells containing sequences encoding MEMAP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding MEMAP under the control of a single promoter. Expression of the marker gene in response to induction or selection
35 usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding MEMAP and that express MEMAP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of MEMAP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on MEMAP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding MEMAP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding MEMAP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding MEMAP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode MEMAP may be designed to contain signal sequences which direct secretion of MEMAP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of

the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding MEMAP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric MEMAP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of MEMAP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the MEMAP encoding sequence and the heterologous protein sequence, so that MEMAP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled MEMAP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

MEMAP of the present invention or fragments thereof may be used to screen for compounds that specifically bind to MEMAP. At least one and up to a plurality of test compounds may be screened for specific binding to MEMAP. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of

MEMAP, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which MEMAP binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express MEMAP, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing MEMAP or cell membrane fractions which contain MEMAP are then contacted with a test compound and binding, stimulation, or inhibition of activity of either MEMAP or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with MEMAP, either in solution or affixed to a solid support, and detecting the binding of MEMAP to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

MEMAP of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of MEMAP. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for MEMAP activity, wherein MEMAP is combined with at least one test compound, and the activity of MEMAP in the presence of a test compound is compared with the activity of MEMAP in the absence of the test compound. A change in the activity of MEMAP in the presence of the test compound is indicative of a compound that modulates the activity of MEMAP. Alternatively, a test compound is combined with an in vitro or cell-free system comprising MEMAP under conditions suitable for MEMAP activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of MEMAP may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding MEMAP or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of

interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding MEMAP may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding MEMAP can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding MEMAP is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress MEMAP, e.g., by secreting MEMAP in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

25 THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of MEMAP and membrane associated proteins. In addition, the expression of MEMAP is closely associated with neurological and gastrointestinal tissues, cancer, cell proliferation, and inflammation/trauma. Therefore, MEMAP appears to play a role in cell proliferative, autoimmune/inflammatory, neurological and gastrointestinal disorders. In the treatment of disorders associated with increased MEMAP expression or activity, it is desirable to decrease the expression or activity of MEMAP. In the treatment of disorders associated with decreased MEMAP expression or activity, it is desirable to increase the expression or activity of MEMAP.

Therefore, in one embodiment, MEMAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or

activity of MEMAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including

mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas.

In another embodiment, a vector capable of expressing MEMAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of MEMAP including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified MEMAP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of MEMAP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of MEMAP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of MEMAP including, but not limited to, those listed above.

In a further embodiment, an antagonist of MEMAP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of MEMAP. Examples of such disorders include, but are not limited to, those cell proliferative, autoimmune/inflammatory, neurological and gastrointestinal disorders described above. In one aspect, an antibody which specifically binds MEMAP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express MEMAP.

In an additional embodiment, a vector expressing the complement of the polynucleotide

encoding MEMAP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of MEMAP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of MEMAP may be produced using methods which are generally known in the art. In particular, purified MEMAP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind MEMAP. Antibodies to MEMAP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with MEMAP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to MEMAP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of MEMAP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to MEMAP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce MEMAP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for MEMAP may also be generated. For example, such fragments include, but are not limited to, $F(ab')_2$ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between MEMAP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering MEMAP epitopes is generally used, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for MEMAP. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of MEMAP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple MEMAP epitopes, represents the average affinity, or avidity, of the antibodies for MEMAP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular MEMAP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in

which the MEMAP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of MEMAP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of MEMAP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al., supra.)

In another embodiment of the invention, the polynucleotides encoding MEMAP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding MEMAP. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding MEMAP. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding MEMAP may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency

(e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in MEMAP expression or regulation causes disease, the expression of MEMAP from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in MEMAP are treated by constructing mammalian expression vectors encoding MEMAP and introducing these vectors by mechanical means into MEMAP-deficient cells. Mechanical transfer technologies for use with cells *in vivo* or *ex vitro* include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J.-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of MEMAP include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). MEMAP may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the

FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, *supra*)), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding MEMAP from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to MEMAP expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding MEMAP under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et al. (1997) *Blood* 89:2259-2267; Bonyhadi, M.L. (1997) *J. Virol.* 71:4707-4716; Ranga, U. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:1201-1206; Su, L. (1997) *Blood* 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding MEMAP to cells which have one or more genetic abnormalities with respect to the expression of MEMAP. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have

proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) *Transplantation* 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) *Annu. Rev. Nutr.* 19:511-544; and Verma, I.M. and N. Somia (1997) *Nature* 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding MEMAP to target cells which have one or more genetic abnormalities with respect to the expression of MEMAP. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing MEMAP to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) *Exp. Eye Res.* 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) *J. Virol.* 73:519-532 and Xu, H. et al. (1994) *Dev. Biol.* 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding MEMAP to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) *Curr. Opin. Biotechnol.* 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full-length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for MEMAP into the alphavirus genome in place of the capsid-coding region results in the production of a large number of MEMAP-coding RNAs and the synthesis of high levels of MEMAP in vector transduced cells. While

alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of MEMAP into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding MEMAP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding MEMAP. Such DNA sequences may be incorporated into a wide variety of

vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding MEMAP. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased MEMAP expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding MEMAP may be therapeutically useful, and in the treatment of disorders associated with decreased MEMAP expression or activity, a compound which specifically promotes expression of the polynucleotide encoding MEMAP may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding MEMAP is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding MEMAP are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence

complementary to the sequence of the polynucleotide encoding MEMAP. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of MEMAP, antibodies to MEMAP, and mimetics, agonists, antagonists, or inhibitors of MEMAP.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the

case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising MEMAP or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, MEMAP or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example MEMAP or fragments thereof, antibodies of MEMAP, and agonists, antagonists or inhibitors of MEMAP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the

active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind MEMAP may be used for the diagnosis of disorders characterized by expression of MEMAP, or in assays to monitor patients being treated with MEMAP or agonists, antagonists, or inhibitors of MEMAP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for MEMAP include methods which utilize the antibody and a label to detect MEMAP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring MEMAP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of MEMAP expression. Normal or standard values for MEMAP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to MEMAP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of MEMAP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding MEMAP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of MEMAP may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess

expression of MEMAP, and to monitor regulation of MEMAP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding MEMAP or closely related molecules may be used to identify nucleic acid sequences which encode MEMAP. The specificity of the probe, whether
5 it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding MEMAP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50%
10 sequence identity to any of the MEMAP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:38-74 or from genomic sequences including promoters, enhancers, and introns of the MEMAP gene.

Means for producing specific hybridization probes for DNAs encoding MEMAP include the cloning of polynucleotide sequences encoding MEMAP or MEMAP derivatives into vectors for the
15 production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding MEMAP may be used for the diagnosis of disorders
20 associated with expression of MEMAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including
25 adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS),
30 Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis,
35 erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves'

disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a gastrointestinal disorder such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis,

peliosis hepatitis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas. The polynucleotide sequences encoding MEMAP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR
5 technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered MEMAP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding MEMAP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide
10 sequences encoding MEMAP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding MEMAP in the
15 sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of MEMAP, a normal or standard profile for expression is established. This may be accomplished by
20 combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding MEMAP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values
25 obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from
30 successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance
35 of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals

to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding MEMAP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding MEMAP, or a fragment of a polynucleotide complementary to the polynucleotide encoding MEMAP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding MEMAP may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding MEMAP are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of MEMAP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the

polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described in Seilhamer, J.J. et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, incorporated herein by reference. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, antibodies specific for MEMAP, or MEMAP or fragments thereof may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and

toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The

optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic
5 cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for MEMAP to quantify
10 the levels of MEMAP expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendozze, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a
15 thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson,
20 N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

25 In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic
30 response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are
35 incubated with antibodies specific to the polypeptides of the present invention. The amount of

protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

5 Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are
10 well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding MEMAP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may
15 be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes
20 (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism
25 (RFLP). (See, e.g., Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding MEMAP on a
30 physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse,
35 may reveal associated markers even if the exact chromosomal locus is not known. This information is

valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, MEMAP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between MEMAP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with MEMAP, or fragments thereof, and washed. Bound MEMAP is then detected by methods well known in the art. Purified MEMAP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding MEMAP specifically compete with a test compound for binding MEMAP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with MEMAP.

In additional embodiments, the nucleotide sequences which encode MEMAP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/149,641 and U.S. Ser. No. 60/164,203 are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Genomics, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1

ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) *Anal. Biochem.* 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incye cDNA recovered in plasmids as described in Example II were sequenced as follows.

Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (PE Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (PE Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned

sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programing, and dinucleotide nearest neighbor analysis. The sequences were then queried
 5 against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA.
 10 The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene
 15 families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:38-74. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above:

20 IV. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

25 Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$30 \quad \frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is
 35 calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the

product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding MEMAP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Chromosomal Mapping of MEMAP Encoding Polynucleotides

The cDNA sequences which were used to assemble SEQ ID NO:38-74 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:38-74 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

The genetic map locations of SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:44, SEQ ID NO:60, SEQ ID NO:63, and SEQ ID NO:67 are described in The Invention as ranges, or intervals, of human chromosomes. More than one map location is reported for SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:60, and SEQ ID NO:63, indicating that previously mapped sequences having similarity, but not complete identity, to SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:44, SEQ ID NO:60, and SEQ ID NO:63 were assembled into their respective

clusters. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.)

- 5 The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Diseases associated with the public and Incyte sequences located within the indicated intervals are also reported in the Invention where applicable. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

VI. Extension of MEMAP Encoding Polynucleotides

- The full length nucleic acid sequences of SEQ ID NO:38-74 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

- High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

- The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar,

Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems).

In like manner, the polynucleotide sequences of SEQ ID NO:38-74 are used to obtain 5' regulatory sequences using the procedure above, along with oligonucleotides designed for such extension, and an appropriate genomic library.

VII. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:38-74 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a

SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

5 The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and
10 compared.

VIII. Microarrays

 The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the
15 aforementioned technologies should be uniform and solid with a non-porous surface (Skena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to
20 those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Skena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

 Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be
25 selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser
30 desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

35 Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and

poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40 μ M dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37 °C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85 °C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μ l 5X SSC/0.2% SDS.

15 Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60 °C followed by washes in

0.2% SDS and distilled water as before.

Hybridization

Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital

(A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

IX. Complementary Polynucleotides

Sequences complementary to the MEMAP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring MEMAP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of MEMAP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the MEMAP-encoding transcript.

X. Expression of MEMAP

Expression and purification of MEMAP is achieved using bacterial or virus-based expression systems. For expression of MEMAP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express MEMAP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of MEMAP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding MEMAP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases.

Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

- In most expression systems, MEMAP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from *Schistosoma japonicum*, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from MEMAP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10 and 16). Purified MEMAP obtained by these methods can be used directly in the assays shown in Examples XI and XV.

XI. Demonstration of MEMAP Activity

- MEMAP activity is demonstrated using a generic immunoblotting strategy or through a MEMAP-specific activity assay as outlined below. As a general approach, cell lines or tissues transformed with a vector containing MEMAP coding sequences can be assayed for MEMAP activity by immunoblotting. Transformed cells are denatured in SDS in the presence of β -mercaptoethanol, nucleic acids are removed by ethanol precipitation, and proteins are purified by acetone precipitation. Pellets are resuspended in 20 mM tris buffer at pH 7.5 and incubated with Protein G-Sepharose pre-coated with an antibody specific for MEMAP. After washing, the Sepharose beads are boiled in electrophoresis sample buffer, and the eluted proteins subjected to SDS-PAGE. Proteins are transferred from the SDS-PAGE gel to a membrane for immunoblotting, and the MEMAP activity is assessed by visualizing and quantifying bands on the blot using antibody specific for MEMAP as the primary antibody and 125 I-labeled IgG specific for the primary antibody as the secondary antibody.

- A specific assay for MEMAP activity measures the expression of MEMAP on the cell surface. cDNA encoding MEMAP is transfected into a mammalian (non-human) cell line. Cell surface proteins are labeled with biotin as described in de la Fuente, M.A. et al. ((1997) Blood 90:2398-2405). Immunoprecipitations are performed using MEMAP-specific antibodies, and immunoprecipitated samples are analyzed using SDS-PAGE and immunoblotting techniques. The ratio of labeled immunoprecipitant to unlabeled immunoprecipitant is proportional to the amount of MEMAP expressed on the cell surface.

- In an alternative specific assay, MEMAP transport activity is assayed by measuring uptake of

labeled substrates into Xenopus laevis oocytes. Oocytes at stages V and VI are injected with MEMAP mRNA (10 ng per oocyte) and incubated for 3 days at 18°C in OR2 medium (82.5mM NaCl, 2.5 mM KCl, 1mM CaCl₂, 1mM MgCl₂, 1mM Na₂HPO₄, 5 mM Hepes, 3.8 mM NaOH , 50µg/ml gentamycin, pH 7.8) to allow expression of MEMAP protein. Oocytes are then transferred to standard uptake medium (100mM NaCl, 2 mM KCl, 1mM CaCl₂, 1mM MgCl₂, 10 mM Hepes/Tris pH 7.5). Uptake of various substrates (e.g., amino acids, sugars, drugs, and neurotransmitters) is initiated by adding a ³H substrate to the oocytes. After incubating for 30 minutes, uptake is terminated by washing the oocytes three times in Na⁺-free medium, measuring the incorporated ³H, and comparing with controls. MEMAP activity is proportional to the level of internalized ³H substrate.

XII. Functional Assays

MEMAP function is assessed by expressing the sequences encoding MEMAP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10 µg of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 µg of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of MEMAP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding MEMAP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions

of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding MEMAP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIII. Production of MEMAP Specific Antibodies

MEMAP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the MEMAP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (PE Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-MEMAP activity by, for example, binding the peptide or MEMAP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIV. Purification of Naturally Occurring MEMAP Using Specific Antibodies

Naturally occurring or recombinant MEMAP is substantially purified by immunoaffinity chromatography using antibodies specific for MEMAP. An immunoaffinity column is constructed by covalently coupling anti-MEMAP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing MEMAP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of MEMAP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/MEMAP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and MEMAP is collected.

XV. Identification of Molecules Which Interact with MEMAP

MEMAP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate

molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled MEMAP, washed, and any wells with labeled MEMAP complex are assayed. Data obtained using different concentrations of MEMAP are used to calculate values for the number, affinity, and association of MEMAP with the candidate molecules.

5 Alternatively, molecules interacting with MEMAP are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, Nature 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

 MEMAP may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions
10 between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

 Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the
15 invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	38	112301	PITUNOT01	003382R1 (HMC1NOT01), 094523R1 (PITUNOT01), 112301H1 (PITUNOT01), 301778X11 (TESTNOT04), 320551X13 (EOSIHET02), 1368852R1 (SCORN02), 1800784H1 (COLNNOT27), 2117174X17C1 (BRSTTUT02), 2526345F6 (BRAITUT21), 4333609H1 (KIDCTMT01)
2	39	997947	KIDNTUT01	997947H1 (KIDNTUT01), 997947T6 (KIDNTUT01), 1417936X306D1 (KIDNNOT09), 1672062X307V1 (BLADNOT05), 3738956T6 (MENTNOT01), SCCA01437V1, SCCA05013V1, SCCA01691V1, SCCA02873V1
3	40	1521513	BLADTUT04	1222062H1 (NEUTGMT01), 1521513H1 (BLADTUT04), 1521513T1 (BLADTUT04), 3558522F6 (LUNGNOT31), 3558522T6 (LUNGNOT31)
4	41	1863994	PROSNOT19	265171R6 (HNT2AGT01), 1863994H1 (PROSNOT19), 3750444F6 (UTRSNOT18), 4177677F6 (BRAINOT22), 4697638F6 (BRALNOT01), 4774040F6 (BRAQNOT01), SCEA02960V1
5	42	2071941	ISLTNOT01	286350R1 (EOSIHET02), 491305R1 (HNT2AGT01), 724168R1 (SYNOOAT01), 1466668F1 (PANCUTUT02), 2071941H1 (ISLTNOT01), 2071941X11C1 (ISLTNOT01), 3579445H1 (293TF3T01)
6	43	2172512	ENDCNOT03	2172512H1 (ENDCNOT03), 2544419F6 (UTRSNOT11), 2798626H1 (NPOLNOT01), 3203359H1 (PENCNOT02), 91241299
7	44	2483172	SMCANOT01	217987F1 (STOMNOT01), 1289703F6 (BRAINOT11), 1289703T6 (BRAINOT11), 2211377F6 (SINPFET03), 2483172H1 (SMCANOT01), 2493236H1 (ADRETUT05), 3274006F6 (PROSBPT06)

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
8	45	2656128	THYMN0T04	2654722T6 (THYMN0T04), 2656128H1 (THYMN0T04), 2837168F6 (THYMN0T03)
9	46	5855841	FIBAUNT02	894553T1 (BRSTN0T05), 1296289F1 (PGANN0T03), 1466541T1 (PANCUT02), 2046927F6 (THP1T7T01), 2058873R6 (OVARN0T03), 3800875F6 (SPLN0T12), 5855841H1 (FIBAUNT02)
10	47	603462	BRSTTUT01	603462H1 (BRSTTUT01), 1487733H1 (UCMCL5T01), 1750451F6 (STOMTUT02), 5182853T6 (LUNGTM0T03)
11	48	747681	BRAITUT01	747681H1 (BRAITUT01), 752009R1 (BRAITUT01), 1267874F1 (BRAIN0T09), 1833308R6 (BRAIN0N01), 2673538X19F1 (KIDN0T19), SB0A07003F3, SCDA07521V1, SCDA04982V1, SCDA07275V1
12	49	919469	RATRN0T02	153337R6 (THP1PLB02), 1525415F6 (UCMCL5T01), 1527804F1 (UCMCL5T01), 1985565R6 (LUNGAST01), 2397811T6 (THP1A2T01), SAR01416F1, SARA03198F1
13	50	977658	BRSTN0T02	977658H1 (BRSTN0T02), 1873689F6 (LEUKN0T02), 2155095F6 (BRAIN0T09), 2186432F6 (PROSN0T26), 2204117F6 (SPLNFET02), 2206291F6 (SPLNFET02), 3255048R6 (OVARTUN01), 3501520H1 (ADREN0T11), 3743427H1 (THYMN0T08)
14	51	1004703	BRSTN0T03	742178H1 (PANCN0T04), 1444583F6 (THYRN0T03), 2068902X15C1 (ISLTN0T01), 2616367F6 (GBLAN0T01), SBVA02190V1
15	52	1334051	COLNN0T13	3222815T6 (COLNN0N03), SXBC00794V1, SXBC00639V1
16	53	1336728	COLNN0T13	630458R6 (KIDN0T05), 1336728H1 (COLNN0T13), SXBC00758V1, SXBC01825V1, SXBC01531V1, SXBC01624V1, SXBC00128V1

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
17	54	1452856	PENITUT01	873008R1 (LUNGAST01), 1452856H1 (PENITUT01), 2433573H1 (BRAVUNT02), 2444932F6 (THP1NOT03), 2858295F6 (SININOT03)
18	55	1562471	SPLNNOT04	286237F1 (EOSIHET02), 1562471H1 (SPLNNOT04), 1880730F6 (LEUKNOT03), 3420608F6 (UCMCNOT04), SBWA00968V1, SXBC01387V1, SBWA02301V1
19	56	1618158	BRAITUT12	967563R1 (BRSTNOT05), 1618158H1 (BRAITUT12), 1785271F6 (BRAINOT10), 2074680F6 (ISLTNOT01), 2822196H1 (ADRETUT06)
20	57	1656935	URETTUT01	1656935F6 (URETTUT01), 1656935H1 (URETTUT01), 2827605F6 (TLYMNOT03), 5272146H1 (OVARINOT02), g1482116
21	58	1859305	PROSNOT18	079372F1 (SYNORAB01), 639845R1 (BRSTNOT03), 1859305H1 (PROSNOT18), 3328091F6 (HEACNOT04), 3354812F6 (PROSNOT28), 5510642H1 (BRADDIR01)
22	59	1949083	PITUNOT01	1287161H1 (BRAINOT11), 1949083H1 (PITUNOT01), 1949083R6 (PITUNOT01), 1949083T6 (PITUNOT01), 3814131F6 (TONSNOT03)
23	60	1996357	BRSTTUT03	260527R6 (HNT2RAT01), 260527T6 (HNT2RAT01), 1313441F1 (BLADTUT02), 1442781R1 (THYRNOT03), 1996357H1 (BRSTTUT03), 1996357T6 (BRSTTUT03), 4262451H1 (BSCNDIT02), SAZA00147F1
24	61	2061330	OVARNOT03	2061330H1 (OVARNOT03), 2724233T6 (LUNGUTUT10), 5104031T6 (PROSTUS20)
25	62	2346947	TESTTUT02	2346947F6 (TESTTUT02), 2346947H1 (TESTTUT02), 4051345F6 (SINTNOT18)

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
26	63	2795577	NPOLNOT01	867213R6 (BRAITUT03), 2381770H1 (ISLTNOT01), 2795577CT1 (NPOLNOT01), 2795577H1 (NPOLNOT01)
27	64	3255825	OVARTUN01	3255825CT1 (OVARTUN01), 3255825H1 (OVARTUN01)
28	65	3393430	LUNGNOT28	2187169H1 (PROSNOT26), 3393256H1 (LUNGNOT28), 3393430H1 (LUNGNOT28), 3395774H1 (LUNGNOT28), 4689688H1 (LIVRTUT12), 4895996H1 (LIVRTUT12), 4896461F6 (LIVRTUT12), 4984527F6 (LIVRTUT10), 4992946H1 (LIVRTUT11)
29	66	3490990	EPIGNOT01	1235428F1 (LUNGFET03), 1662973T6 (BRSTNOT09), 2362021H1 (LUNGFET05), 2362021R6 (LUNGFET05), 3490990H1 (EPIGNOT01)
30	67	3635154	LIVRNOT03	027592H1 (SPLNFET01), 3635154H1 (LIVRNOT03), g1012932
31	68	4374347	CONFNOT03	860875X11 (BRAITUT03), 898143R6 (BRSTNOT05), 4374347H1 (CONFNOT03)
32	69	4596747	COLSTUT01	137213R1 (SYNORAB01), 545568R6 (OVARNOT02), 1235402F1 (LUNGFET03), 1268010F1 (BRAINOT09), 1271078F1 (TESTTUT02), 1301951F6 (BRSTNOT07), 1994442R6 (BRSTTUT03), 2343102H1 (TESTTUT02), 3274538F6 (PROSBPT06), 4596747H1 (COLSTUT01)
33	70	5052680	BRSTNOT33	1973688H1 (UCMCL5T01), 3926410F6 (KIDNNOT19), 4501839F6 (BRAVXT02), 5052680F6 (BRSTNOT33), 5052680H1 (BRSTNOT33), 5186780F6 (LUNGWT04)
34	71	5373575	BRAINOT22	262776T6 (HNT2AGT01), 1234057F1 (LUNGFET03), 1741526R6 (HIPONON01), 1871204F6 (SKINBIT01), 2192479F6 (THYRTUT03), 2556849H1 (THYMNOT03), 2722451T6 (LUNGTUT10), 4114985H1 (UTRSTUT07), 5373575H1 (BRAINOT22)

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
35	72	5524468	LIVRDIR01	4024068F6 (BRAXNOT02), 5524468H1 (LIVRDIR01), SXBC01952V1
36	73	5944279	COLADIT05	1662182H1 (BRSTNOT09), 1698677F6 (BLADTUT05), 1916639R6 (PROSNOT06), 1916639T6 (PROSNOT06), 2298565R6 (BRSTNOT05), 2298565T6 (BRSTNOT05), 2583019F6 (KIDNTUT13), 2870903F6 (THYRNOT10), 3970715H1 (PROSTUT10), 3971695H1 (PROSTUT10), 5944279H1 (COLADIT05)
37	74	6114480	SINITMT04	1579843F6 (DUODNOT01), 1579843T6 (DUODNOT01), 4181024T6 (SINITUT03), 6114480H1 (SINITMT04), SXBC00007V1, SXBC00504V1, SCSA05104V1

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
1	351	S31 T116 S169 T229 T2 S209 T306	N128	Signal peptide: M1-A33	Paraneoplastic neural antigen MA1 [Homo sapiens] g4104634	BLAST-GenBank MOTIFS SPSCAN
2	458	T198 S27 S37 T87 S251 S257 S325 S373 S405 S422 T454 T210 S228 S401 Y93	N75 N159 N279 N445	Signal peptide: M1-T24 Glycoprotein signature: C199-L448	Pancortin-3 [Mus musculus] g3218528	BLAST-GenBank MOTIFS SPSCAN HMNER BLAST-PRODOM
3	219	T51 S120 S163 T175 T181 S3 T12 T45 S75 S104 S128	N2 N62 N107	Signal peptide: M1-C42 Transmembrane domain: L32-F49 C-type lectin domain: C80-E206	Murine macrophage C- type lectin [Mus musculus] g5821286	BLAST-GenBank MOTIFS SPSCAN HMNER HMNER-PFAM BLIMPS-BLOCKS PROFILESAN BLAST-DOMO
4	276	S213 S91 S113 S35 S70 S76 S147 T163 S206		Signal peptide: M1-G31 Transmembrane domain: I184-F201 Cell attachment sequence: R149-D151		BLAST-GenBank MOTIFS HMNER

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
5	375	S18 S205 T286 S3 S120 S197 T260 Y85		Transmembrane domains: W139-R158; F173-H191 P232-Q254 Transmembrane protein signature: I95-C369	Transmembrane protein [S. pombe] g1065898	BLAST-GenBank MOTIFS HMMER BLAST-DOMO BLAST-PRODOM
6	249	T7 T135 T170 S204 Y154	N18 N92 N147		Phospholipid scramblase [Homo sapiens] g4092081	BLAST-GenBank MOTIFS
7	353	T162 T4 S97 T115 S165 S194 T225 S242 S17 S47 S205	N299	Signal peptide: M1-A33	Paraneoplastic neuronal antigen MA1 [Homo sapiens] g4104634	BLAST-GenBank MOTIFS SPSCAN
8	194	T12 S115 S29 S99 S187	N95 N147	Signal peptide: M1-C50 Transmembrane domain: L38-L56 C-type lectin domain: C75-E194	Lectin-like NK cell receptor LLT1 [Homo sapiens] g6651065	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLIMPS-BLOCKS BLAST-DOMO
9	322	S304 S48 S146 S72 T133 S255 S280	N20 N60 N70	Signal peptide: M1-A50		BLAST-GenBank MOTIFS SPSCAN

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
10	335	S125 S140 S183 S222 T252		Transmembrane domains: G71-L94; A255-I283 Gufa transmembrane protein domain: L12-H101; G180-G335 Glycosaminoglycan attachment site: S310-G313	Gufa protein [Thermotoga maritima] g4982315	BLAST-GenBank MOTIFS HMMER BLAST-PRODOM BLAST-DOMO
11	620	S49 S108 T146 S300 T348 T349 S607 S4 S128 S183 S234 T420 S460 S467 S543 Y597	N144 N202 N264 N274 N293 N341 N492 N505 N526 N542	Transmembrane domain: M563-W582 Immunoglobulin domain: G439-A499 Leucine-rich repeat signature: L337-L350 Glycoprotein hormone receptor domain: T40-L198	Slit2 [Rattus norvegicus] g4585574	BLAST-GenBank MOTIFS HMMER HMMER-PFAM BLIMPS-PRINTS BLAST-DOMO

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
12	491	T231 T232 S253 T482 S185 S276	N56 N220 N229	Transmembrane domains: I115-I142; I184-V201 F422-F441 Transmembrane protein domain: L8-Y215; I396-F471	Selectively expressed in embryonic epithelia protein-1 [Mus musculus] g6715148 PB39 [Homo sapiens] g3462515	BLAST-GenBank MOTIFS HMMER BLAST-PRODOM
13	580	S557 S10 T34 S51 T92 T210 S343 T12 S217 T222 S268 S296 T417 T523 S550	N159 N179 N220 N230	Transmembrane domains: F297-F313; I356-I373 L496-I514 Lipases serine active site: L104-A113		MOTIFS HMMER
14	455	T53 T182 S239 S69 S135 S202 T280 S355 S372 Y38	N67 N180 N243	Transmembrane domains: V81-V99; I343-I361 S375-V392; W425-Y442 Glycosaminoglycan attachment site: S162-G165	putative G- protein coupled receptor [Homo sapiens] g6649579	BLAST-GenBank MOTIFS HMMER

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
15	277	S265 T66 T225 S268 S273 S30 S49 S61 S152 S193 Y242	N29 N38 N47 N48 N92 N160 N210	Transmembrane domain: K9-F27 Brush border protein domain: Y8-R277 RGD cell attachment sequence: R113-D115	AdRab-A brush border membrane protein (Oryctolagus cuniculus) g1762	BLAST-GenBank MOTIFS HMMER BLAST-PRODOR
16	647	S490 T50 S67 S105 T110 S121 T220 S249 S264 S272 S322 T389 S469 T501 S639 S132 T155 S242 S324 T381 T400 S522 S554	N261	Signal peptide: M1-A22 Transmembrane domains: L328-L347; M406-L424 L559-A578; W618-L638 Gufa transmembrane protein domain: E485-L640 Glycosaminoglycan attachment site: S34-G37	LIV-1 protein [Homo sapiens] g1256001	BLAST-GenBank MOTIFS SPSCAN HMMER BLAST-PRODOR
17	406	S29 S215 S236 T69	N23	Transmembrane domains: Q76-V95; W286-S313 M367-I384		MOTIFS HMMER

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
18	290	T221 S44 S69 S71 S81 T94 T101 T113 T131 S216 Y284	N88	Signal peptide: M1-A19 Transmembrane domains: P160-M181 Immunoglobulin domain: R33-I110 Transmembrane glycoprotein domain: I22-D116	NK inhibitory receptor [Homo sapiens] g6707799 CMRF-35-H9 leukocyte antigen [Homo sapiens] g4103066	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLAST-PRODOM BLAST-DOMO
19	390	S7 T68 S153 T23 T166 T281 Y20 Y37	N5 N88 N330 N367	Immunoglobulins and MHC proteins signature: T90-P112; F242-V259 Glycoprotein antigen signature: L61-V72; V92-I113		MOTIFS BLIMPS-BLOCKS BLIMPS-PRODOM
20	427	S13 S41 S65 S66 S99 T150 S323 S324 S101 S275 S353 S367 T399 Y71	N106 N148 N171 N233 N312	Mucin glycoprotein precursor domain: V136-P142	Gastric mucin [Sus scrofa] g915208	BLAST-GenBank MOTIFS BLIMPS-PRODOM

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
21	459	T4 S60 S66 S116 T176 S16 T235	N14 N158 N323	Transmembrane domains: F202-V219; I246-L268 W343-L367; P417-P440	six transmembrane epithelial antigen of prostate [Homo sapiens] g6572948	BLAST-GenBank MOTIFS HMMER
22	229	S13 S118 T155 Y24		Transmembrane domains: I93-V111; V132-L150 F164-V182 Transmembrane protein domain: S156-V182		MOTIFS HMMER BLIMPS-PRODOM
23	311	S85 S234 S236 S269 S80 S119 S186 T294	N22	Transmembrane domains: W58-I76; P152-K177 A216-Y232		MOTIFS HMMER
24	92	S47 T54 T12 S70	N62		HERV-E envelope glycoprotein [Homo sapiens] g2587024	BLAST-GenBank MOTIFS
25	258	S34 T33 S148 S243		Transmembrane domains: I39-I57; F86-L106 V122-I140; L190-S210		MOTIFS HMMER

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
26	226	S56 S128 T196 T167 Y194	N54 N187 N198	Signal peptide: M1-P50 Transmembrane domains: T23-L43; M72-A89 I101-I124; I158-N178 Transmembrane 4 family signature: A70-I120 Lysosomal-associated transmembrane protein domain: C15-Y223	MTP (mouse transporter protein) [Mus musculus] g1276631	BLAST-GenBank MOTIFS SPSCAN HMMER PROFILES SCAN BLAST-PRODOM
27	136	S3 S132		Signal peptide: M1-R53 Transmembrane domains: I10-L28; T26-I50 F70-L89 Transmembrane protein domain: D31-V104		MOTIFS SPSCAN HMMER BLAST-PRODOM

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
28	458	T408 T98 S126 S170 T334	N96 N151 N293 N332	Signal peptide: M1-A20 Transmembrane domain: L10-N30 Membrane glycoprotein signature: L9-V101; L64-Q457 Olfactory ligand binding domain: T67-S452	Potential ligand (odorant) binding protein [Rattus rattus] g57732	BLAST-GenBank MOTIFS SPSCAN HMMER BLAST-PRODOM BLAST-DOMO
29	368	S24 T166 T302 S12 S134 Y307	N17		Fuzzy (TM protein involved in tissue polarity) [Drosophila melanogaster] g2564657	BLAST-GenBank MOTIFS
30	91	T44 S84		Signal peptide: M1-A19 Transmembrane domain: P58-S82 Glycophorin A proteins signature: T22-S32; I63-G91 Glycophorin domain: M1-R86	Preglycophorin B [Homo sapiens] g4803699	BLAST-GenBank MOTIFS SPSCAN HMMER BLIMPS-BLOCKS PROFILESKAN BLAST-PRODOM BLAST-DOMO

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
31	295	S96 T113 S129 T155 T125 T157 T187 S222 T231 T263 Y212	N111 N169 N223	Signal peptide: M1-G48 Transmembrane domain: L241-L259 Immunoglobulin domain: K159-V216 Carcinoembryonic antigen domain: I38-P147 Glycoprotein antigen domain: M1-V140; Y141-Y234 G239-S295	Biliary glycoprotein [Mus musculus] g312590	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLAST-PRODOM BLAST-DOMO
32	724	T39 S47 T171 S205 T224 S225 T241 S285 S301 T323 S352 T353 S439 S509 S517 S537 T659 T707 S8 S18 S49 S72 T85 T159 S173 S271 S367 S560 S588 Y499	N279 N348	Transmembrane domain: I611-F630 Membrane protein domain: T4-L209		MOTIFS HMMER BLAST-DOMO

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
33	331	S117 S147 S149 T320 S138 S174 T274 T319 S328 Y198	N222	Signal peptide: M1-S16 Transmembrane domains: A67-N87; I118-C134 W240-V269; L294-Y310 Transmembrane protein domain: A6-T311	Putative Golgi UDP-GlcNAc transporter (S. pombe) g3738167	BLAST-GenBank MOTIFS SPSCAN HMMER BLAST-PRODOM
34	398	T42 T158 S271 S28 S285 T334 S375		Transmembrane domain: I59-L79 Band 7 family domain: F64-A231, A78-V90; R116-L154 Stomatin signature: T84-L106; L131-P152 T166-L183; I186-G209 L54-Q227	Stomatin-like protein UNC24 [Homo sapiens] g5326747	BLAST-GenBank MOTIFS HMMER HMMER-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO
35	220	S199 T120 S192	N107	Signal peptide: M1-G19 Leucine rich repeats: A62-F85; Q86-S109 G110-G133; A134-R157 A158-S181; H184-P207	Similar to Leucine-rich transmembrane proteins [Homo sapiens] g2781386	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLIMPS-PRINTS

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
36	706	T564 T74 T113 S291 S452 S632 S14 T42 S66 T115 T142 S286 T551 T575 S701	N101	Transmembrane domains: F158-M178; L344-V368 L425-L442; M478-F498 A581-I604; L641-V665 Glycosaminoglycan attachment site: S223-G226	LAK-4p [Homo sapiens] g7209574	BLAST-GenBank MOTIFS HMMER
37	466	T326 S10 T46 T105 S187 S98 T164 T310 S321 Y388	N368	Signal peptide: M1-G23 Transmembrane domain: A236-I255 SPRY domain: A338-S464; E123-S136 E322-W343; V407-F420 Butyrophilin domain: W19-C114	Butyrophilin like receptor [Homo sapiens] g4587209	BLAST-GenBank MOTIFS SPSCAN HMMER HMMER-PFAM BLIMPS-PFAM BLAST-PRODOM BLAST-DOMO

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
38	844-888	Nervous (0.377) Reproductive (0.180) Cardiovascular (0.115) Gastrointestinal (0.115)	Cancer (0.410) Inflammation/Trauma (0.296) Cell Proliferation (0.131)	PBLUESCRIPT
39	579-623	Developmental (0.400) Musculoskeletal (0.200) Nervous (0.200) Urologic (0.200)	Cancer (0.400) Cell Proliferation (0.400)	PSPORT1
40	336-380	Cardiovascular (0.267) Hematopoietic/Immune (0.200) Endocrine (0.133) Reproductive (0.133)	Cancer (0.400) Inflammation/Trauma (0.400) Cell Proliferation (0.133)	pINCY
41	596-640	Nervous (0.588) Gastrointestinal (0.118) Reproductive (0.118)	Inflammation/Trauma (0.470) Cancer (0.235) Cell Proliferation (0.176)	pINCY
42	1281-1325	Reproductive (0.237) Hematopoietic/Immune (0.145) Nervous (0.145)	Cancer (0.441) Inflammation/Trauma (0.323) Cell Proliferation (0.178)	pINCY
43	227-271	Reproductive (0.444) Dermatologic (0.222) Endocrine (0.111) Gastrointestinal (0.111) Nervous (0.111)	Cancer (0.333) Cell Proliferation (0.222) Inflammation/Trauma (0.222)	pINCY

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
44	1368-1412	Nervous (0.339) Reproductive (0.278) Gastrointestinal (0.104)	Cancer (0.478) Inflammation/Trauma (0.278) Cell Proliferation (0.165)	pINCY
45	543-587	Hematopoietic/Immune (0.500) Gastrointestinal (0.188)	Inflammation/Trauma (0.500) Cancer (0.250) Cell Proliferation (0.188)	pINCY
46	280-324	Reproductive (0.267) Nervous (0.233) Gastrointestinal (0.112)	Cancer (0.483) Inflammation/Trauma (0.345) Cell Proliferation (0.155)	pINCY
47	380-424 875-919	Reproductive (0.412) Gastrointestinal (0.176) Cardiovascular (0.118)	Cancer (0.647) Inflammation/Trauma (0.178)	PSPORT1
48	272-316 1514-1558	Nervous (0.645) Developmental (0.129)	Cancer (0.355) Cell Proliferation (0.258) Neurological (0.194)	PSPORT1
49	282-326 768-812	Hematopoietic/Immune (0.238) Gastrointestinal (0.155) Reproductive (0.143)	Cancer (0.381) Inflammation/Trauma (0.381) Cell Proliferation (0.202)	PSPORT1
50	597-641 1074-1118	Reproductive (0.214) Nervous (0.196) Hematopoietic/Immune (0.143)	Cancer (0.464) Inflammation/Trauma (0.304) Cell Proliferation (0.196)	PSPORT1
51	973-1017	Reproductive (0.266) Nervous (0.234) Hematopoietic/Immune (0.125)	Cancer (0.516) Inflammation/Trauma (0.359) Cell Proliferation (0.109)	PSPORT1

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
52	299-343	Gastrointestinal (1.000)	Cancer (0.500) Inflammation/Trauma (0.500)	pINCY
53	380-424 1199-1243	Gastrointestinal (0.289) Reproductive (0.244) Cardiovascular (0.111) Hematopoietic/Immune (0.111)	Cancer (0.578) Inflammation/Trauma (0.311) Cell Proliferation (0.178)	pINCY
54	1135-1179	Nervous (0.195) Reproductive (0.186) Gastrointestinal (0.144)	Cancer (0.449) Inflammation/Trauma (0.305) Cell Proliferation (0.144)	pINCY
55	325-369 820-864	Hematopoietic/Immune (0.750)	Inflammation/Trauma (0.625) Cancer (0.125)	pINCY
56	487-531 1090-1134	Nervous (0.583)	Cancer (0.458) Inflammation/Trauma (0.250)	pINCY
57	569-613 1360-1405	Reproductive (0.429) Hematopoietic/Immune (0.286) Musculoskeletal (0.143) Urologic (0.143)	Cancer (0.571) Inflammation/Trauma (0.286) Cell Proliferation (0.143)	pINCY
58	272-472 551-595 812-1012 1523-1567	Reproductive (0.350) Nervous (0.150) Cardiovascular (0.100) Gastrointestinal (0.100) Hematopoietic/Immune (0.100) Urologic (0.100)	Cancer (0.500) Inflammation/Trauma (0.500)	pINCY

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
59	217-261	Nervous (0.286) Developmental (0.143) Gastrointestinal (0.143) Hematopoietic/Immune (0.143) Reproductive (0.143)	Inflammation/Trauma (0.428) Cancer (0.357) Cell Proliferation (0.143)	PBLUESCRIPT
60	444-488	Nervous (0.207) Reproductive (0.207) Gastrointestinal (0.130) Hematopoietic/Immune (0.130)	Cancer (0.467) Inflammation/Trauma (0.359) Cell Proliferation (0.163)	PSPORT1
61	643-687	Reproductive (0.464) Endocrine (0.143) Cardiovascular (0.107) Gastrointestinal (0.107)	Cancer (0.500) Inflammation/Trauma (0.321)	PSPORT1
62	146-344 390-434 506-704 786-830	Gastrointestinal (0.500) Hematopoietic/Immune (0.250) Reproductive (0.250)	Cancer (0.750) Inflammation/Trauma (0.250)	pINCY
63	163-207	Reproductive (0.315) Gastrointestinal (0.161) Cardiovascular (0.147)	Cancer (0.594) Cell Proliferation (0.231) Inflammation/Trauma (0.210)	pINCY
64	201-506 525-569 606-912 975-1280 1362-1406	Gastrointestinal (0.455) Cardiovascular (0.273) Reproductive (0.189)	Cancer (0.455) Inflammation/Trauma (0.367) Cell Proliferation (0.189)	PSPORT1

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
65	703-747	Gastrointestinal (0.667) Cardiovascular (0.167) Reproductive (0.167)	Cancer (1.000)	pINCY
66	271-315 319-363	Nervous (0.314) Reproductive (0.314) Developmental (0.114) Urologic (0.114)	Cancer (0.429) Cell Proliferation (0.171) Inflammation/Trauma (0.143)	pINCY
67	319-363	Developmental (0.364) Hematopoietic/Immune (0.364) Gastrointestinal (0.182)	Cell Proliferation (0.727) Cancer (0.273) Inflammation/Trauma (0.182)	pINCY
68	812-856	Reproductive (0.444) Nervous (0.222) Endocrine (0.111) Hematopoietic/Immune (0.111) Musculoskeletal (0.111)	Cancer (0.556) Inflammation/Trauma (0.333)	pINCY
69	596-640 1577-1621	Reproductive (0.255) Nervous (0.184) Developmental (0.122) Gastrointestinal (0.122)	Cancer (0.429) Inflammation/Trauma (0.337) Cell Proliferation (0.255)	pINCY
70	379-675 703-747 766-1062 1081-1347	Nervous (0.467) Hematopoietic/Immune (0.200) Reproductive (0.133) Urologic (0.133)	Cancer (0.467) Cell Proliferation (0.267) Inflammation/Trauma (0.267)	pINCY

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
71	18-62	Nervous (0.265) Reproductive (0.206) Musculoskeletal (0.147)	Cancer (0.500) Inflammation/Trauma (0.264) Cell Proliferation (0.147)	pINCY
72	290-488 507-704 759-803	Gastrointestinal (0.333) Hematopoietic/Immune (0.333) Nervous (0.333)	Inflammation/Trauma (0.667) Cancer (0.333)	pINCY
73	649-693 1711-1755	Reproductive (0.392) Gastrointestinal (0.294) Cardiovascular (0.118)	Cancer (0.686) Inflammation/Trauma (0.294)	pINCY
74	704-748	Gastrointestinal (0.923)	Cancer (0.462) Inflammation/Trauma (0.385)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
38	PITUNOT01	This library was constructed using RNA obtained from Clontech (CLON 6584-2, lot 35278). The RNA was isolated from pituitary glands removed from a pool of 18 male and female Caucasian donors, 16 to 70 years old, who died from trauma.
39	KIDNTUT01	This library was constructed using RNA isolated from kidney tumor tissue removed from an 8-month-old female during nephroureterectomy. Pathology indicated Wilms' tumor (nephroblastoma), which involved 90 percent of the renal parenchyma. Prior to surgery, the patient was receiving heparin anticoagulant therapy.
40	BLADTUT04	This library was constructed using RNA isolated from bladder tumor tissue removed from a 60-year-old Caucasian male during a radical cystectomy, prostatectomy, and vasectomy. Pathology indicated grade 3 transitional cell carcinoma in the left bladder wall. Carcinoma in-situ was identified in the dome and trigone. Patient history included tobacco use. Family history included type I diabetes, malignant neoplasm of the stomach, atherosclerotic coronary artery disease, and acute myocardial infarction.
41	PROSNOT19	This library was constructed using RNA isolated from diseased prostate tissue removed from a 59-year-old Caucasian male during a radical prostatectomy with regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+3). The patient presented with elevated prostate-specific antigen (PSA). Patient history included colon diverticuli, asbestosis, and thrombophlebitis. Family history included benign hypertension, multiple myeloma, hyperlipidemia and rheumatoid arthritis.
42	ISLTNOT01	This library was constructed using RNA isolated from a pooled collection of pancreatic islet cells.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
43	ENDCNOT03	This library was constructed using RNA isolated from dermal microvascular endothelial cells removed from a neonatal Caucasian male.
44	SMCANOT01	This library was constructed using RNA isolated from an aortic smooth muscle cell line derived from the explanted heart of a male obtained during a heart transplant.
45	THYMNOT04	This library was constructed using RNA isolated from thymus tissue removed from a 3-year-old Caucasian male, who died from anoxia.
46	FIBAUNT02	This library was constructed using RNA isolated from untreated aortic adventitial fibroblasts removed from a 65-year-old Caucasian female.
47	BRSTTUT01	This library was constructed using RNA isolated from breast tumor tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated invasive grade 4 mammary adenocarcinoma. Patient history included atrial tachycardia and a benign breast neoplasm. Family history included cardiovascular and cerebrovascular disease and depressive disorder.
48	BRAITUT01	This library was constructed using RNA isolated from brain tumor tissue removed from a 50-year-old Caucasian female during a frontal lobectomy. Pathology indicated recurrent grade 3 oligoastrocytoma with focal necrosis and extensive calcification. Patient history included a speech disturbance and epilepsy. The patient's brain had also been irradiated with a total dose of 5,082 cGy (Fraction 8). Family history included a brain tumor.
49	RATRNOT02	This library was constructed using RNA isolated from the right atrium tissue of a 39-year-old Caucasian male, who died from a gunshot wound.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
50	BRSTNOT02	This library was constructed using RNA isolated from diseased breast tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated proliferative fibrocystic changes characterized by apocrine metaplasia, sclerosing adenosis, cyst formation, and ductal hyperplasia without atypia. Pathology for the associated tumor tissue indicated an invasive grade 4 mammary adenocarcinoma. Patient history included atrial tachycardia and a benign neoplasm. Family history included cardiovascular and cerebrovascular disease.
51	BRSTNOT03	This library was constructed using RNA isolated from diseased breast tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy. Pathology for the associated tumor tissue indicated residual invasive grade 3 mammary ductal adenocarcinoma. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia and a malignant neoplasm of the colon.
52	COLNNOT13	This library was constructed using RNA isolated from ascending colon tissue of a 28-year-old Caucasian male with moderate chronic ulcerative colitis.
53	COLNNOT13	This library was constructed using RNA isolated from ascending colon tissue of a 28-year-old Caucasian male with moderate chronic ulcerative colitis.
54	PENITUT01	This library was constructed using RNA isolated from tumor tissue removed from the penis of a 64-year-old Caucasian male during penile amputation. Pathology indicated a fungating invasive grade 4 squamous cell carcinoma involving the inner wall of the foreskin and extending onto the glans penis. Patient history included benign neoplasm of the large bowel, atherosclerotic coronary artery disease, angina pectoris, gout, and obesity. Family history included malignant pharyngeal neoplasm, chronic lymphocytic leukemia, and chronic liver disease.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
55	SPLNNOT04	This library was constructed using RNA isolated from the spleen tissue of a 2-year-old Hispanic male, who died from cerebral anoxia.
56	BRAITUT12	This library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 40-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated grade 4 gemistocytic astrocytoma.
57	URETTUT01	This library was constructed using RNA isolated from right ureter tumor tissue of a 69-year-old Caucasian male during ureterectomy and lymph node excision. Pathology indicated invasive grade 3 transitional cell carcinoma. Patient history included benign colon neoplasm, tobacco use, asthma, emphysema, acute duodenal ulcer, and hyperplasia of the prostate. Family history included atherosclerotic coronary artery disease, congestive heart failure, and malignant lung neoplasm.
58	PROSNOT18	This library was constructed using RNA isolated from diseased prostate tissue removed from a 58-year-old Caucasian male during a radical cystectomy, radical prostatectomy, and gastrectomy. Pathology indicated adenofibromatous hyperplasia; this tissue was associated with a grade 3 transitional cell carcinoma. Patient history included angina and emphysema. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
59	PITUNOT01	This library was constructed using RNA obtained from Clontech (CLON 6584-2, lot 35278). The RNA was isolated from the pituitary glands removed from a pool of 18 male and female Caucasian donors, 16 to 70 years old, who died from trauma.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
60	BRSTTUT03	This library was constructed using RNA isolated from breast tumor tissue removed from a 58-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated multicentric invasive grade 4 lobular carcinoma. The mass was identified in the upper outer quadrant, and three separate nodules were found in the lower outer quadrant of the left breast. Patient history included skin cancer, rheumatic heart disease, osteoarthritis, and tuberculosis. Family history included cerebrovascular disease, coronary artery aneurysm, breast cancer, prostate cancer, atherosclerotic coronary artery disease, and type I diabetes.
61	OVARNOT03	This library was constructed using RNA isolated from ovarian tissue removed from a 43-year-old Caucasian female during removal of the fallopian tubes and ovaries. Pathology for the associated tumor tissue indicated grade 2 mucinous cystadenocarcinoma. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, pancreatic cancer, stress reaction, cerebrovascular disease, breast cancer, and uterine cancer.
62	TESTTUT02	This library was constructed using RNA isolated from testicular tumor tissue removed from a 31-year-old Caucasian male during unilateral orchiectomy. Pathology indicated embryonal carcinoma.
63	NPOLNOT01	This library was constructed using RNA isolated from nasal polyp tissue removed from a 78-year-old Caucasian male during a nasal polypectomy. Pathology indicated a nasal polyp and striking eosinophilia. Patient history included asthma and nasal polyps.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
64	OVARTUN01	This normalized library was constructed from 5.36 million independent clones obtained from an ovarian tumor library. RNA was isolated from tumor tissue removed from the left ovary of a 58-year-old Caucasian female during a total abdominal hysterectomy, removal of a single ovary, and inguinal hernia repair. Pathology indicated a metastatic grade 3 adenocarcinoma of colonic origin, forming a partially cystic and necrotic tumor mass in the left ovary, and a nodule in the left mesovarium. A single intramural leiomyoma was identified in the myometrium. The cervix showed mild chronic cystic cervicitis. Patient history included benign hypertension, follicular ovarian cyst, colon cancer, benign colon neoplasm, and osteoarthritis. Family history included emphysema, myocardial infarction, atherosclerotic coronary artery disease, benign hypertension, hyperlipidemia, and primary tuberculous complex. The normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9228) and Bonaldo et al. (Genome Research (1996) 6:791).
65	LUNGNOT28	This library was constructed using RNA isolated from lung tissue removed from a 53-year-old male. Pathology for the associated tumor tissue indicated grade 4 adenocarcinoma.
66	EPIGNOT01	This library was constructed using RNA isolated from epiglottic tissue removed from a 71-year-old male during laryngectomy with right parathyroid biopsy. Pathology for the associated tumor tissue indicated recurrent grade 1 papillary thyroid carcinoma.
67	LIVRNOT03	This library was constructed using RNA isolated from liver tissue removed from a Caucasian male fetus, who died from Patau's syndrome (trisomy 13) at 20 weeks' gestation.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
68	CONFNOT03	This library was constructed using RNA isolated from mesenteric fat tissue removed from a 71-year-old Caucasian male during a partial colectomy and permanent colostomy. Pathology indicated mesenteric fat tissue associated with diverticulosis and diverticulitis with abscess formation. Approximately 50 diverticula were noted, one of which was perforated and associated with abscess formation in adjacent mesenteric fat. The patient presented with atrial fibrillation. Patient history included viral hepatitis, a hemangioma, and diverticulitis of colon. Family history included extrinsic asthma, atherosclerotic coronary artery disease, and myocardial infarction.
69	COLSTUT01	This library was constructed using RNA isolated from colon tumor tissue removed from the sigmoid colon of a 62-year-old Caucasian male during a sigmoidectomy and permanent colostomy. Pathology indicated invasive grade 2 adenocarcinoma, with invasion through the muscularis. Patient history included hyperlipidemia, cataract disorder and dermatitis. Family history included benign hypertension, atherosclerotic coronary artery disease, hyperlipidemia, breast cancer, and prostate cancer.
70	BRSTNOT33	This library was constructed using RNA isolated from right breast tissue removed from a 46-year-old Caucasian female during a unilateral extended simple mastectomy with breast reconstruction. Pathology for the associated tumor tissue indicated invasive grade 3 adenocarcinoma, ductal type, with apocrine features, nuclear grade 3 forming a mass in the outer quadrant. There was greater than 50% intraductal component. Patient history included breast cancer.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
71	BRAINOT22	This library was constructed using RNA isolated from right temporal lobe tissue removed from a 45-year-old Black male during a brain lobectomy. Pathology for the associated tumor tissue indicated dysembryoplastic neuroepithelial tumor of the right temporal lobe. The right temporal region dura was consistent with calcifying pseudotumor of the neuraxis. Patient history included obesity, meningitis, backache, unspecified sleep apnea, acute stress reaction, acquired knee deformity, and chronic sinusitis. Family history included obesity, benign hypertension, cirrhosis of the liver, obesity, hyperlipidemia, cerebrovascular disease, and type II diabetes.
72	LIVRDIR01	This library was constructed using RNA isolated from diseased liver tissue removed from a 63-year-old Caucasian female during a liver transplant. Patient history included primary biliary cirrhosis. Serology was positive for anti-mitochondrial antibody.
73	COLADIT05	This library was constructed using RNA isolated from diseased ascending colon tissue removed from a 32-year-old Caucasian male during a total intra-abdominal colectomy, abdominal-perineal rectal resection, and temporary ileostomy. Pathology indicated chronic ulcerative colitis extending in a continuous fashion from the mid-portion of the ascending colon distally to the rectum. This was characterized microscopically by crypt abscess formation and inflammation confined to the mucosa and submucosa. The terminal ileum exhibited ileitis and the rectal mucosa showed crypt abscess formation. Patient history included tobacco use. Family history included ulcerative colitis, malignant neoplasm of the breast and acute myocardial infarction.

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
74	SINITWT04	Library was constructed using RNA isolated from ileum tissue removed from a 70-year-old Caucasian female during right hemicolectomy, open liver biopsy, flexible sigmoidoscopy, colonoscopy, and permanent colostomy. Pathology indicated a non-tumorous margin of ileum. Pathology for the associated tumor indicated invasive grade 2 adenocarcinoma forming an ulcerated mass, situated 2 cm distal to the ileocecal valve. The tumor invaded through the muscularis propria just into the serosal adipose tissue. One (of 16) regional lymph node was positive for a microfocus of metastatic adenocarcinoma. Patient history included a malignant breast neoplasm, type II diabetes, hyperlipidemia, viral hepatitis, an unspecified thyroid disorder, osteoarthritis, and a malignant skin neoplasm. Family history included breast cancer, atherosclerotic coronary artery disease, benign hypertension, cerebrovascular disease, ovarian cancer, and hyperlipidemia.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	PE Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	PE Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	PE Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value= 1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families; sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37,

15 b) a naturally occurring amino acid sequence having at least 70% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37,

20 c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, and

25 d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37.

35 2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1,

SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37.

3. An isolated polynucleotide encoding a polypeptide of claim 1.

10 4. An isolated polynucleotide encoding a polypeptide of claim 2.

5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74.

20 6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

7. A cell transformed with a recombinant polynucleotide of claim 6.

25 8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

9. A method for producing a polypeptide of claim 1, the method comprising:

a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

35

11. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74,
- b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74,
- c) a polynucleotide sequence complementary to a),
- d) a polynucleotide sequence complementary to b), and
- e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37.

18. A method for treating a disease or condition associated with decreased expression of functional MEMAP, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of functional MEMAP, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide

of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

5 23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

 24. A method for treating a disease or condition associated with overexpression of functional MEMAP, comprising administering to a patient in need of such treatment a composition of claim 23.

10

 25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- 15 b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

 26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- 20 a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound
- 25 with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

 27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- 30 a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

35 28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- c) quantifying the amount of hybridization complex; and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

SEQUENCE LISTING

<110> INCYTE GENOMICS, INC.

LAL, Preeti

YUE, Henry

TANG, Y. Tom

BANDMAN, Olga

BURFORD, Neil

AZIMZAI, Yalda

BAUGHN, Mariah R.

LU, Dyung Aina M.

PATTERSON, Chandra

<120> MEMBRANE ASSOCIATED PROTEINS

<130> PF-0731 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/149,641; 60/164,203

<151> 1999-08-17; 1999-11-09

<160> 74

<170> PERL Program

<210> 1

<211> 351

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 112301CD1

<400> 1

Met	Thr	Leu	Arg	Leu	Leu	Glu	Asp	Trp	Cys	Arg	Gly	Met	Asp	Met
1				5					10					15
Asn	Pro	Arg	Lys	Ala	Leu	Leu	Ile	Ala	Gly	Ile	Ser	Gln	Ser	Cys
				20					25					30
Ser	Val	Ala	Glu	Ile	Glu	Glu	Ala	Leu	Gln	Ala	Gly	Leu	Ala	Pro
				35					40					45
Leu	Gly	Glu	Tyr	Arg	Leu	Leu	Gly	Arg	Met	Phe	Arg	Arg	Asp	Glu
				50					55					60
Asn	Arg	Lys	Val	Ala	Leu	Val	Gly	Leu	Thr	Ala	Glu	Thr	Ser	His
				65					70					75
Ala	Leu	Val	Pro	Lys	Glu	Ile	Pro	Gly	Lys	Gly	Gly	Ile	Trp	Arg
				80					85					90
Val	Ile	Phe	Lys	Pro	Pro	Asp	Pro	Asp	Asn	Thr	Phe	Leu	Ser	Arg
				95					100					105
Leu	Asn	Glu	Phe	Leu	Ala	Gly	Glu	Gly	Met	Thr	Val	Gly	Glu	Leu
				110					115					120
Ser	Arg	Ala	Leu	Gly	His	Glu	Asn	Gly	Ser	Leu	Asp	Pro	Glu	Gln
				125					130					135
Gly	Met	Ile	Pro	Glu	Met	Trp	Ala	Pro	Met	Leu	Ala	Gln	Ala	Leu
				140					145					150
Glu	Ala	Leu	Gln	Pro	Ala	Leu	Gln	Cys	Leu	Lys	Tyr	Lys	Lys	Leu
				155					160					165
Arg	Val	Phe	Ser	Gly	Arg	Glu	Ser	Pro	Glu	Pro	Gly	Glu	Glu	Glu
				170					175					180
Phe	Gly	Arg	Trp	Met	Phe	His	Thr	Thr	Gln	Met	Ile	Lys	Ala	Trp

Gln Val Pro Asp	185	Val Glu Lys Arg Arg	190	Arg Leu Leu Glu Ser	195
Arg Gly Pro Ala	200	Leu Asp Val Ile Arg	205	Val Leu Lys Ile Asn	210
Pro Leu Ile Thr	215	Val Asp Glu Cys Leu	220	Gln Ala Leu Glu Glu	225
Phe Gly Val Thr	230	Asp Asn Pro Arg Glu	235	Leu Gln Val Lys Tyr	240
Thr Thr Tyr Gln	245	Lys Asp Glu Glu Lys	250	Leu Ser Ala Tyr Val	255
Arg Leu Glu Pro	260	Leu Leu Gln Lys Leu	265	Val Gln Arg Gly Ala	270
Glu Arg Asp Ala	275	Val Asn Gln Ala Arg	280	Leu Asp Gln Val Ile	285
Gly Ala Val His	290	Lys Thr Ile Arg Arg	295	Glu Leu Asn Leu Pro	300
Asp Gly Pro Ala	305	Pro Gly Phe Leu Gln	310	Leu Leu Val Leu Ile	315
Asp Tyr Glu Ala	320	Ala Glu Glu Glu Glu	325	Ala Leu Leu Gln Ala	330
Leu Glu Gly Asn	335	Phe Thr	340		345
	350				

<210> 2

<211> 458

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 997947CD1

<400> 2

Met Gln Ala Thr	Ser Asn Leu Leu Asn	Leu Leu Leu Ser	Leu
1	5	10	15
Phe Ala Gly Leu	Asp Pro Ser Lys Thr	Gln Ile Ser Pro	Lys Glu
	20	25	30
Gly Trp Gln Val	Tyr Ser Ser Ala Gln	Asp Pro Asp Gly	Arg Cys
	35	40	45
Ile Cys Thr Val	Val Ala Pro Glu Gln	Asn Leu Cys Ser	Arg Asp
	50	55	60
Ala Lys Ser Arg	Gln Leu Arg Gln Leu	Glu Lys Val Gln	Asn
	65	70	75
Met Ser Gln Ser	Ile Glu Val Leu Asn	Leu Arg Thr Gln	Arg Asp
	80	85	90
Phe Gln Tyr Val	Leu Lys Met Glu Thr	Gln Met Lys Gly	Leu Lys
	95	100	105
Ala Lys Phe Arg	Gln Ile Glu Asp Asp	Arg Lys Thr Leu	Met Thr
	110	115	120
Lys His Phe Gln	Glu Leu Lys Glu Lys	Met Asp Glu Leu	Leu Pro
	125	130	135
Leu Ile Pro Val	Leu Glu Gln Tyr Lys	Thr Asp Ala Lys	Leu Ile
	140	145	150
Thr Gln Phe Lys	Glu Glu Ile Arg Asn	Leu Ser Ala Val	Leu Thr
	155	160	165
Gly Ile Gln Glu	Glu Ile Gly Ala Tyr	Asp Tyr Glu Glu	Leu His
	170	175	180
Gln Arg Val Leu	Ser Leu Glu Thr Arg	Leu Arg Asp Cys	Met Lys
	185	190	195
Lys Leu Thr Cys	Gly Lys Leu Met Lys	Ile Thr Gly Pro	Val Thr
	200	205	210
Val Lys Thr Ser	Gly Thr Arg Phe Gly	Ala Trp Met Thr	Asp Pro
	215	220	225

Leu Ala Ser Glu Lys Asn Asn Arg Val Trp Tyr Met Asp Ser Tyr
 230 235 240
 Thr Asn Asn Lys Ile Val Arg Glu Tyr Lys Ser Ile Ala Asp Phe
 245 250 255
 Val Ser Gly Ala Glu Ser Arg Thr Tyr Asn Leu Pro Phe Lys Trp
 260 265 270
 Ala Gly Thr Asn His Val Val Tyr Asn Gly Ser Leu Tyr Phe Asn
 275 280 285
 Lys Tyr Gln Ser Asn Ile Ile Ile Lys Tyr Ser Phe Asp Met Gly
 290 295 300
 Arg Val Leu Ala Gln Arg Ser Leu Glu Tyr Ala Gly Phe His Asn
 305 310 315
 Val Tyr Pro Tyr Thr Trp Gly Gly Phe Ser Asp Ile Asp Leu Met
 320 325 330
 Ala Asp Glu Ile Gly Leu Trp Ala Val Tyr Ala Thr Asn Gln Asn
 335 340 345
 Ala Gly Asn Ile Val Ile Ser Gln Leu Asn Gln Asp Thr Leu Glu
 350 355 360
 Val Met Lys Ser Trp Ser Thr Gly Tyr Pro Lys Arg Ser Ala Gly
 365 370 375
 Glu Ser Phe Met Ile Cys Gly Thr Leu Tyr Val Thr Asn Ser His
 380 385 390
 Leu Thr Gly Ala Lys Val Tyr Tyr Ser Tyr Ser Thr Lys Thr Ser
 395 400 405
 Thr Tyr Glu Tyr Thr Asp Ile Pro Phe His Asn Gln Tyr Phe His
 410 415 420
 Ile Ser Met Leu Asp Tyr Asn Ala Arg Asp Arg Ala Leu Tyr Ala
 425 430 435
 Trp Asn Asn Gly His Gln Val Leu Phe Asn Val Thr Leu Phe His
 440 445 450
 Ile Ile Lys Thr Glu Asp Asp Thr
 455

<210> 3

<211> 219

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1521513CD1

<400> 3

Met Asn Ser Ser Lys Ser Ser Glu Thr Gln Cys Thr Glu Arg Gly
 1 5 10 15
 Cys Phe Ser Ser Gln Met Phe Leu Trp Thr Val Ala Gly Ile Pro
 20 25 30
 Ile Leu Phe Leu Ser Ala Cys Phe Ile Thr Arg Cys Val Val Thr
 35 40 45
 Phe Arg Ile Phe Gln Thr Cys Asp Glu Lys Lys Phe Gln Leu Pro
 50 55 60
 Glu Asn Phe Thr Glu Leu Ser Cys Tyr Asn Tyr Gly Ser Gly Ser
 65 70 75
 Val Lys Asn Cys Cys Pro Leu Asn Trp Glu Tyr Phe Gln Ser Ser
 80 85 90
 Cys Tyr Phe Phe Ser Thr Asp Thr Ile Ser Trp Ala Leu Ser Leu
 95 100 105
 Lys Asn Cys Ser Ala Met Gly Ala His Leu Val Val Ile Asn Ser
 110 115 120
 Gln Glu Glu Gln Glu Phe Leu Ser Tyr Lys Lys Pro Lys Met Arg
 125 130 135
 Glu Phe Phe Ile Gly Leu Ser Asp Gln Val Val Glu Gly Gln Trp
 140 145 150
 Gln Trp Val Asp Gly Thr Pro Leu Thr Lys Ser Leu Ser Phe Trp

Asp Val Gly Glu	155	Pro Asn Asn Ile Ala	160	Thr Leu Glu Asp Cys	165
	170		175		180
Thr Met Arg Asp	185	Ser Ser Asn Pro Arg	190	Gln Asn Trp Asn Asp	195
Thr Cys Phe Leu	200	Asn Tyr Phe Arg Ile	205	Cys Glu Met Val Gly	210
Asn Pro Leu Asn	215	Lys Gly Lys Ser Leu			

<210> 4
 <211> 276
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1863994CD1

Met Glu Ser Arg	Met	Trp	Pro	Ala	Leu	Leu	Leu	Ser	His	Leu	Leu
1	5				10						15
Pro Leu Trp	Pro	Leu	Leu	Leu	Pro	Leu	Pro	Pro	Pro	Ala	Gln
	20				25						30
Gly Ser Ser	Ser	Pro	Arg	Thr	Pro	Pro	Ala	Pro	Ala	Arg	Pro
	35				40						45
Pro Cys Ala	Arg	Gly	Gly	Pro	Ser	Ala	Pro	Arg	His	Val	Cys
	50				55						60
Trp Glu Arg	Ala	Pro	Pro	Ser	Arg	Ser	Pro	Arg	Val	Pro	Arg
	65				70						75
Ser Arg Arg	Gln	Val	Leu	Pro	Gly	Thr	Ala	Pro	Pro	Ala	Thr
	80				85						90
Ser Gly Phe	Glu	Gly	Pro	Pro	Ser	Ser	Gln	Tyr	Pro	Trp	Ala
	95				100						105
Ile Val Trp	Gly	Pro	Thr	Val	Ser	Arg	Glu	Asp	Gly	Gly	Asp
	110				115						120
Asn Ser Ala	Asn	Pro	Gly	Phe	Leu	Asp	Tyr	Gly	Phe	Ala	Ala
	125				130						135
His Gly Leu	Ala	Thr	Pro	His	Pro	Asn	Ser	Asp	Ser	Met	Arg
	140				145						150
Asp Gly Asp	Gly	Leu	Ile	Leu	Gly	Glu	Ala	Pro	Ala	Thr	Leu
	155				160						165
Pro Phe Leu	Phe	Gly	Arg	Gly	Glu	Gly	Val	Asp	Pro	Gln	Leu
	170				175						180
Tyr Val Thr	Ile	Thr	Ile	Ser	Ile	Ile	Ile	Val	Leu	Val	Ala
	185				190						195
Gly Ile Ile	Phe	Lys	Phe	Cys	Trp	Asp	Arg	Ser	Gln	Lys	Arg
	200				205						210
Arg Pro Ser	Gly	Gln	Gln	Gly	Ala	Leu	Arg	Gln	Glu	Glu	Ser
	215				220						225
Gln Pro Leu	Thr	Asp	Leu	Ser	Pro	Ala	Gly	Val	Thr	Val	Leu
	230				235						240
Ala Phe Gly	Asp	Ser	Pro	Thr	Pro	Thr	Pro	Asp	His	Glu	Glu
	245				250						255
Arg Gly Gly	Pro	Arg	Pro	Gly	Met	Pro	His	Pro	Lys	Gly	Ala
	260				265						270
Ala Phe Gln	Leu	Asn	Arg								
	275										

<210> 5
 <211> 375
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2071941CD1

<400> 5

```

Met Ser Ser His Lys Gly Ser Val Val Ala Gln Gly Asn Gly Ala
 1      5      10      15
Pro Ala Ser Asn Arg Glu Ala Asp Thr Val Glu Leu Ala Glu Leu
 20      25      30
Gly Pro Leu Leu Glu Glu Lys Gly Lys Arg Val Ile Ala Asn Pro
 35      40      45
Pro Lys Ala Glu Glu Glu Gln Thr Cys Pro Val Pro Gln Glu Glu
 50      55      60
Glu Glu Glu Val Arg Val Leu Thr Leu Pro Leu Gln Ala His His
 65      70      75
Ala Met Glu Lys Met Glu Glu Phe Val Tyr Lys Val Trp Glu Gly
 80      85      90
Arg Trp Arg Val Ile Pro Tyr Asp Val Leu Pro Asp Trp Leu Lys
 95      100     105
Asp Asn Asp Tyr Leu Leu His Gly His Arg Pro Pro Met Pro Ser
110     115     120
Phe Arg Ala Cys Phe Lys Ser Ile Phe Arg Ile His Thr Glu Thr
125     130     135
Gly Asn Ile Trp Thr His Leu Leu Gly Phe Val Leu Phe Leu Phe
140     145     150
Leu Gly Ile Leu Thr Met Leu Arg Pro Asn Met Tyr Phe Met Ala
155     160     165
Pro Leu Gln Glu Lys Val Val Phe Gly Met Phe Phe Leu Gly Ala
170     175     180
Val Leu Cys Leu Ser Phe Ser Trp Leu Phe His Thr Val Tyr Cys
185     190     195
His Ser Glu Lys Val Ser Arg Thr Phe Ser Lys Leu Asp Tyr Ser
200     205     210
Gly Ile Ala Leu Leu Ile Met Gly Ser Phe Val Pro Trp Leu Tyr
215     220     225
Tyr Ser Phe Tyr Cys Ser Pro Gln Pro Arg Leu Ile Tyr Leu Ser
230     235     240
Ile Val Cys Val Leu Gly Ile Ser Ala Ile Ile Val Ala Gln Trp
245     250     255
Asp Arg Phe Ala Thr Pro Lys His Arg Gln Thr Arg Ala Gly Val
260     265     270
Phe Leu Gly Leu Gly Leu Ser Gly Val Val Pro Thr Met His Phe
275     280     285
Thr Ile Ala Glu Gly Phe Val Lys Ala Thr Thr Val Gly Gln Met
290     295     300
Gly Trp Phe Phe Leu Met Ala Val Met Tyr Ile Thr Gly Ala Gly
305     310     315
Leu Tyr Ala Ala Arg Ile Pro Glu Arg Phe Phe Pro Gly Lys Phe
320     325     330
Asp Ile Trp Phe Gln Ser His Gln Ile Phe His Val Leu Val Val
335     340     345
Ala Ala Ala Phe Val His Phe Tyr Gly Val Ser Asn Leu Gln Glu
350     355     360
Phe Arg Tyr Gly Leu Glu Gly Gly Cys Thr Asp Asp Thr Leu Leu
365     370     375

```

<210> 6

<211> 249

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2172512CD1

```

<400> 6
Met Ser Gly Val Val Pro Thr Ala Pro Glu Gln Pro Ala Gly Glu
1      5      10      15
Met Glu Asn Gln Thr Lys Pro Pro Asp Pro Arg Pro Asp Ala Pro
20     25     30
Pro Glu Tyr Ser Ser His Phe Leu Pro Gly Pro Pro Gly Thr Ala
35     40     45
Val Pro Pro Pro Thr Gly Tyr Pro Gly Gly Leu Pro Met Gly Tyr
50     55     60
Tyr Ser Pro Gln Gln Pro Ser Thr Phe Pro Leu Tyr Gln Pro Val
65     70     75
Gly Gly Ile His Pro Val Arg Tyr Gln Pro Gly Lys Tyr Pro Met
80     85     90
Pro Asn Gln Ser Val Pro Ile Thr Trp Met Pro Gly Pro Thr Pro
95     100    105
Met Ala Asn Cys Pro Pro Gly Leu Glu Tyr Leu Val Gln Leu Asp
110    115    120
Asn Ile His Val Leu Gln His Phe Glu Pro Leu Glu Met Met Thr
125    130    135
Cys Phe Glu Thr Asn Asn Arg Tyr Asp Ile Lys Asn Asn Ser Asp
140    145    150
Gln Met Val Tyr Ile Val Thr Glu Asp Thr Asp Asp Phe Thr Arg
155    160    165
Asn Ala Tyr Arg Thr Leu Arg Pro Phe Val Leu Arg Val Thr Asp
170    175    180
Cys Met Gly Arg Glu Ile Met Thr Met Gln Arg Pro Phe Arg Cys
185    190    195
Thr Cys Cys Cys Phe Cys Cys Pro Ser Ala Arg Gln Glu Leu Glu
200    205    210
Val Gln Cys Pro Pro Gly Val Thr Ile Gly Phe Val Ala Glu His
215    220    225
Trp Asn Leu Cys Arg Ala Val Tyr Ser Ile Gln Lys Lys Lys Lys
230    235    240
Lys Ile Ala Ala Gln Ala Tyr Ser Leu
245

```

```

<210> 7
<211> 353
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2483172CD1

```

```

<400> 7
Met Ala Met Thr Leu Leu Glu Asp Trp Cys Arg Gly Met Asp Val
1      5      10      15
Asn Ser Gln Arg Ala Leu Leu Val Trp Gly Ile Pro Val Asn Cys
20     25     30
Asp Glu Ala Glu Ile Glu Glu Thr Leu Gln Ala Ala Met Pro Gln
35     40     45
Val Ser Tyr Arg Met Leu Gly Arg Met Phe Trp Arg Glu Glu Asn
50     55     60
Ala Lys Ala Ala Leu Leu Glu Leu Thr Gly Ala Val Asp Tyr Ala
65     70     75
Ala Ile Pro Arg Glu Met Pro Gly Lys Gly Gly Val Trp Lys Val
80     85     90
Leu Phe Lys Pro Pro Thr Ser Asp Ala Glu Phe Leu Glu Arg Leu
95     100    105
His Leu Phe Leu Ala Arg Glu Gly Trp Thr Val Gln Asp Val Ala
110    115    120
Arg Val Leu Gly Phe Gln Asn Pro Thr Pro Thr Pro Gly Pro Glu
125    130    135

```

```

Met Pro Ala Glu Met Leu Asn Tyr Ile Leu Asp Asn Val Ile Gln
140 145 150
Pro Leu Val Glu Ser Ile Trp Tyr Lys Arg Leu Thr Leu Phe Ser
155 160 165
Gly Arg Asp Ile Pro Gly Pro Gly Glu Glu Thr Phe Asp Pro Trp
170 175 180
Leu Glu His Thr Asn Glu Val Leu Glu Glu Trp Gln Val Ser Asp
185 190 195
Val Glu Lys Arg Arg Arg Leu Met Glu Ser Leu Arg Gly Pro Ala
200 205 210
Ala Asp Val Ile Arg Ile Leu Lys Ser Asn Asn Pro Ala Ile Thr
215 220 225
Thr Ala Glu Cys Leu Lys Ala Leu Glu Gln Val Phe Gly Ser Val
230 235 240
Glu Ser Ser Arg Asp Ala Gln Ile Lys Phe Leu Asn Thr Tyr Gln
245 250 255
Asn Pro Gly Glu Lys Leu Ser Ala Tyr Val Ile Arg Leu Glu Pro
260 265 270
Leu Leu Gln Lys Val Val Glu Lys Gly Ala Ile Asp Lys Asp Asn
275 280 285
Val Asn Gln Ala Arg Leu Glu Gln Val Ile Ala Gly Ala Asn His
290 295 300
Ser Gly Ala Ile Arg Arg Gln Leu Trp Leu Thr Gly Ala Gly Glu
305 310 315
Gly Pro Ala Pro Asn Leu Phe Gln Leu Leu Val Gln Ile Arg Glu
320 325 330
Glu Glu Ala Lys Glu Glu Glu Glu Glu Ala Glu Ala Thr Leu Leu
335 340 345
Gln Leu Gly Leu Glu Gly His Phe
350

```

<210> 8
 <211> 194
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2656128CD1

```

<400> 8
Met His Asp Ser Asn Asn Val Glu Lys Asp Ile Thr Pro Ser Glu
1 5 10 15
Leu Pro Ala Asn Pro Gly Cys Leu His Ser Lys Glu His Ser Ile
20 25 30
Lys Ala Thr Leu Ile Trp Arg Leu Phe Phe Leu Ile Met Phe Leu
35 40 45
Thr Ile Ile Val Cys Gly Met Val Ala Ala Leu Ser Ala Ile Arg
50 55 60
Ala Asn Cys His Gln Glu Pro Ser Val Cys Leu Gln Ala Ala Cys
65 70 75
Pro Glu Ser Trp Ile Gly Phe Gln Arg Lys Cys Phe Tyr Phe Ser
80 85 90
Asp Asp Thr Lys Asn Trp Thr Ser Ser Gln Arg Phe Cys Asp Ser
95 100 105
Gln Asp Ala Asp Leu Ala Gln Val Glu Ser Phe Gln Glu Leu Asn
110 115 120
Phe Leu Leu Arg Tyr Lys Gly Pro Ser Asp His Trp Ile Gly Leu
125 130 135
Ser Arg Glu Gln Gly Gln Pro Trp Lys Trp Ile Asn Gly Thr Glu
140 145 150
Trp Thr Arg Gln Leu Val Met Lys Glu Asp Gly Ala Asn Leu Tyr
155 160 165
Val Ala Lys Val Ser Gln Val Pro Arg Met Asn Pro Arg Pro Val

```


Met Val Ser Tyr 170
 185 Pro Gly Ser Arg Arg 175 Val Cys Leu Phe Glu 180
 190

<210> 9
 <211> 322
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5855841CD1

<400> 9
 Met Ser Ser Leu Gly Gly Gly Ser Gln Asp Ala Gly Gly Ser Ser
 1 5 10 15
 Ser Ser Ser Thr Asn Gly Ser Gly Gly Ser Gly Ser Ser Gly Pro
 20 25 30
 Lys Ala Gly Ala Ala Asp Lys Ser Ala Val Val Ala Ala Ala Ala
 35 40 45
 Pro Ala Ser Val Ala Asp Asp Thr Pro Pro Glu Arg Arg Asn
 50 55 60
 Lys Ser Gly Ile Ile Ser Glu Pro Leu Asn Lys Ser Leu Arg Arg
 65 70 75
 Ser Arg Pro Leu Ser His Tyr Ser Ser Phe Gly Ser Ser Gly Gly
 80 85 90
 Ser Gly Gly Gly Ser Met Met Gly Gly Glu Ser Ala Asp Lys Ala
 95 100 105
 Thr Ala Ala Ala Ala Ala Ser Leu Leu Ala Asn Gly His Asp
 110 115 120
 Leu Ala Ala Ala Met Ala Val Asp Lys Ser Asn Pro Thr Ser Lys
 125 130 135
 His Lys Ser Gly Ala Val Ala Ser Leu Leu Ser Lys Ala Glu Arg
 140 145 150
 Ala Thr Glu Leu Ala Ala Glu Gly Gln Leu Thr Leu Gln Gln Phe
 155 160 165
 Ala Gln Ser Thr Glu Met Leu Lys Arg Val Val Gln Glu His Leu
 170 175 180
 Pro Leu Met Ser Glu Ala Gly Ala Gly Leu Pro Asp Met Glu Ala
 185 190 195
 Val Ala Gly Ala Glu Ala Leu Asn Gly Gln Ser Asp Phe Pro Tyr
 200 205 210
 Leu Gly Ala Phe Pro Ile Asn Pro Gly Leu Phe Ile Met Thr Pro
 215 220 225
 Ala Gly Val Phe Leu Ala Glu Ser Ala Leu His Met Ala Gly Leu
 230 235 240
 Ala Glu Tyr Pro Met Gln Gly Glu Leu Ala Ser Ala Ile Ser Ser
 245 250 255
 Gly Lys Lys Lys Arg Lys Arg Cys Gly Met Cys Ala Pro Cys Arg
 260 265 270
 Arg Arg Ile Asn Cys Glu Gln Cys Ser Ser Cys Arg Asn Arg Lys
 275 280 285
 Thr Gly His Gln Ile Cys Lys Phe Arg Lys Cys Glu Glu Leu Lys
 290 295 300
 Lys Lys Pro Ser Ala Ala Leu Glu Lys Val Met Leu Pro Thr Gly
 305 310 315
 Ala Ala Phe Arg Trp Phe Gln
 320

<210> 10
 <211> 335
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 603462CD1

<400> 10

```

Met Leu Gln Gly His Ser Ser Val Phe Gln Ala Leu Leu Gly Thr
 1      5      10      15
Phe Phe Thr Trp Gly Met Thr Ala Ala Gly Ala Ala Leu Val Phe
 20      25      30
Val Phe Ser Ser Gly Gln Arg Arg Ile Leu Asp Gly Ser Leu Gly
 35      40      45
Phe Ala Ala Gly Val Met Leu Ala Ala Ser Tyr Trp Ser Leu Leu
 50      55      60
Ala Pro Ala Val Glu Met Ala Thr Ser Ser Gly Gly Phe Gly Ala
 65      70      75
Phe Ala Phe Phe Pro Val Ala Val Gly Phe Thr Leu Gly Ala Ala
 80      85      90
Phe Val Tyr Leu Ala Asp Leu Leu Met Pro His Leu Gly Ala Ala
 95      100     105
Glu Asp Pro Gln Thr Ala Leu Ala Leu Asn Phe Gly Ser Thr Leu
110     115     120
Met Lys Lys Lys Ser Asp Pro Glu Gly Pro Ala Leu Leu Phe Pro
125     130     135
Glu Ser Glu Leu Ser Ile Arg Ile Asp Lys Ser Glu Asn Gly Glu
140     145     150
Ala Tyr Gln Arg Lys Lys Ala Ala Ala Thr Gly Leu Pro Glu Gly
155     160     165
Pro Ala Val Pro Val Pro Ser Arg Gly Asn Leu Ala Gln Pro Gly
170     175     180
Gly Ser Ser Trp Arg Arg Ile Ala Leu Leu Ile Leu Ala Ile Thr
185     190     195
Ile His Asn Val Pro Glu Gly Leu Ala Val Gly Val Gly Phe Gly
200     205     210
Ala Ile Glu Lys Thr Ala Ser Ala Thr Phe Glu Ser Ala Arg Asn
215     220     225
Leu Ala Ile Gly Ile Gly Ile Gln Asn Phe Pro Glu Gly Leu Ala
230     235     240
Val Ser Leu Pro Leu Arg Gly Ala Gly Phe Ser Thr Trp Arg Ala
245     250     255
Phe Trp Tyr Gly Gln Leu Ser Gly Met Val Glu Pro Leu Ala Gly
260     265     270
Val Phe Gly Ala Phe Ala Val Val Leu Ala Glu Pro Ile Leu Pro
275     280     285
Tyr Ala Leu Ala Phe Ala Ala Gly Ala Met Val Tyr Val Val Met
290     295     300
Asp Asp Ile Ile Pro Glu Ala Gln Ile Ser Gly Asn Gly Lys Leu
305     310     315
Ala Ser Trp Ala Ser Ile Leu Gly Phe Val Val Met Met Ser Leu
320     325     330
Asp Val Gly Leu Gly
335

```

<210> 11

<211> 620

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 747681CD1

<400> 11

```

Met Gln Val Ser Lys Arg Met Leu Ala Gly Gly Val Arg Ser Met
 1      5      10      15
Pro Ser Pro Leu Leu Ala Cys Trp Gln Pro Ile Leu Leu Leu Val

```

	20		25		30
Leu Gly Ser Val	Leu Ser Gly Ser Ala Thr Gly Cys Pro Pro Arg				
	35		40		45
Cys Glu Cys Ser	Ala Gln Asp Arg Ala Val Leu Cys His Arg Lys				
	50		55		60
Arg Phe Val Ala	Val Pro Glu Gly Ile Pro Thr Glu Thr Arg Leu				
	65		70		75
Leu Asp Leu Gly	Lys Asn Arg Ile Lys Thr Leu Asn Gln Asp Glu				
	80		85		90
Phe Ala Ser Phe	Pro His Leu Glu Glu Leu Glu Leu Asn Glu Asn				
	95		100		105
Ile Val Ser Ala	Val Glu Pro Gly Ala Phe Asn Asn Leu Phe Asn				
	110		115		120
Leu Arg Thr Leu	Gly Leu Arg Ser Asn Arg Leu Lys Leu Ile Pro				
	125		130		135
Leu Gly Val Phe	Thr Gly Leu Ser Asn Leu Thr Lys Leu Asp Ile				
	140		145		150
Ser Glu Asn Lys	Ile Val Ile Leu Leu Asp Tyr Met Phe Gln Asp				
	155		160		165
Leu Tyr Asn Leu	Lys Ser Leu Glu Val Gly Asp Asn Asp Leu Val				
	170		175		180
Tyr Ile Ser His	Arg Ala Phe Ser Gly Leu Asn Ser Leu Glu Gln				
	185		190		195
Leu Thr Leu Glu	Lys Cys Asn Leu Thr Ser Ile Pro Thr Glu Ala				
	200		205		210
Leu Ser His Leu	His Gly Leu Ile Val Leu Arg Leu Arg His Leu				
	215		220		225
Asn Ile Asn Ala	Ile Arg Asp Tyr Ser Phe Lys Arg Leu Tyr Arg				
	230		235		240
Leu Lys Val Leu	Glu Ile Ser His Trp Pro Tyr Leu Asp Thr Met				
	245		250		255
Thr Pro Asn Cys	Leu Tyr Gly Leu Asn Leu Thr Ser Leu Ser Ile				
	260		265		270
Thr His Cys Asn	Leu Thr Ala Val Pro Tyr Leu Ala Val Arg His				
	275		280		285
Leu Val Tyr Leu	Arg Phe Leu Asn Leu Ser Tyr Asn Pro Ile Ser				
	290		295		300
Thr Ile Glu Gly	Ser Met Leu His Glu Leu Leu Arg Leu Gln Glu				
	305		310		315
Ile Gln Leu Val	Gly Gly Gln Leu Ala Val Val Glu Pro Tyr Ala				
	320		325		330
Phe Arg Gly Leu	Asn Tyr Leu Arg Val Leu Asn Val Ser Gly Asn				
	335		340		345
Gln Leu Thr Thr	Leu Glu Glu Ser Val Phe His Ser Val Gly Asn				
	350		355		360
Leu Glu Thr Leu	Ile Leu Asp Ser Asn Pro Leu Ala Cys Asp Cys				
	365		370		375
Arg Leu Leu Trp	Val Phe Arg Arg Arg Trp Arg Leu Asn Phe Asn				
	380		385		390
Arg Gln Gln Pro	Thr Cys Ala Thr Pro Glu Phe Val Gln Gly Lys				
	395		400		405
Glu Phe Lys Asp	Phe Pro Asp Val Leu Leu Pro Asn Tyr Phe Thr				
	410		415		420
Cys Arg Arg Ala	Arg Ile Arg Asp Arg Lys Ala Gln Gln Val Phe				
	425		430		435
Val Asp Glu Gly	His Thr Val Gln Phe Val Cys Arg Ala Asp Gly				
	440		445		450
Asp Pro Pro Pro	Ala Ile Leu Trp Leu Ser Pro Arg Lys His Leu				
	455		460		465
Val Ser Ala Lys	Ser Asn Gly Arg Leu Thr Val Phe Pro Asp Gly				
	470		475		480
Thr Leu Glu Val	Arg Tyr Ala Gln Val Gln Asp Asn Gly Thr Tyr				
	485		490		495

```

Leu Cys Ile Ala Ala Asn Ala Gly Gly Asn Asp Ser Met Pro Ala
500 505 510
His Leu His Val Arg Ser Tyr Ser Pro Asp Trp Pro His Gln Pro
515 520 525
Asn Lys Thr Phe Ala Phe Ile Ser Asn Gln Pro Gly Glu Gly Glu
530 535 540
Ala Asn Ser Thr Arg Ala Thr Val Pro Phe Pro Phe Asp Ile Lys
545 550 555
Thr Leu Ile Ile Ala Thr Thr Met Gly Phe Ile Ser Phe Leu Gly
560 565 570
Val Val Leu Phe Cys Leu Val Leu Leu Phe Leu Trp Ser Arg Gly
575 580 585
Lys Gly Asn Thr Lys His Asn Ile Glu Ile Glu Tyr Val Pro Arg
590 595 600
Lys Ser Asp Ala Gly Ile Ser Ser Ala Asp Ala Pro Arg Lys Phe
605 610 615
Asn Met Lys Met Ile
620

```

<210> 12

<211> 491

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 919469CD1

<400> 12

```

Met Ala Gly Gln Gly Leu Pro Leu His Val Ala Thr Leu Leu Thr
1 5 10 15
Gly Leu Leu Glu Cys Leu Gly Phe Ala Gly Val Leu Phe Gly Trp
20 25 30
Pro Ser Leu Val Phe Val Phe Lys Asn Glu Asp Tyr Phe Lys Asp
35 40 45
Leu Cys Gly Pro Asp Ala Gly Pro Ile Gly Asn Ala Thr Gly Gln
50 55 60
Ala Asp Cys Lys Ala Gln Asp Glu Arg Phe Ser Leu Ile Phe Thr
65 70 75
Leu Gly Ser Phe Met Asn Asn Phe Met Thr Phe Pro Thr Gly Tyr
80 85 90
Ile Phe Asp Arg Phe Lys Thr Thr Val Ala Arg Leu Ile Ala Ile
95 100 105
Phe Phe Tyr Thr Thr Ala Thr Leu Ile Ile Ala Phe Thr Ser Ala
110 115 120
Gly Ser Ala Val Leu Leu Phe Leu Ala Met Pro Met Leu Thr Ile
125 130 135
Gly Gly Ile Leu Phe Leu Ile Thr Asn Leu Gln Ile Gly Asn Leu
140 145 150
Phe Gly Gln His Arg Ser Thr Ile Ile Thr Leu Tyr Asn Gly Ala
155 160 165
Phe Asp Ser Ser Ser Ala Val Phe Leu Ile Ile Lys Leu Leu Tyr
170 175 180
Glu Lys Gly Ile Ser Leu Arg Ala Ser Phe Ile Phe Ile Ser Val
185 190 195
Cys Ser Thr Trp His Val Ala Arg Thr Phe Leu Leu Met Pro Arg
200 205 210
Gly His Ile Pro Tyr Pro Leu Pro Pro Asn Tyr Ser Tyr Gly Leu
215 220 225
Cys Pro Gly Asn Gly Thr Thr Lys Glu Glu Lys Glu Thr Ala Glu
230 235 240
His Glu Asn Arg Glu Leu Gln Ser Lys Glu Phe Leu Ser Ala Lys
245 250 255
Glu Glu Thr Pro Gly Ala Gly Gln Lys Gln Glu Leu Arg Ser Phe

```

Trp Ser Tyr Ala	260	Phe Ser Arg Arg Phe	265	Ala Trp His Leu Val	270
	275		280		285
Leu Ser Val Ile	290	Gln Leu Trp His Tyr	295	Leu Phe Ile Gly Thr	300
Asn Ser Leu Leu	305	Thr Asn Met Ala Gly	310	Gly Asp Met Ala Arg	315
Ser Thr Tyr Thr	320	Asn Ala Phe Ala Phe	325	Thr Gln Phe Gly Val	330
Cys Ala Pro Trp	335	Asn Gly Leu Leu Met	340	Asp Arg Leu Lys Gln	345
Tyr Gln Lys Glu	350	Ala Arg Lys Thr Gly	355	Ser Ser Thr Leu Ala	360
Ala Leu Cys Ser	365	Thr Val Pro Ser Leu	370	Ala Leu Thr Ser Leu	375
Cys Leu Gly Phe	380	Ala Leu Cys Ala Ser	385	Val Pro Ile Leu Pro	390
Gln Tyr Leu Thr	395	Phe Ile Leu Gln Val	400	Ile Ser Arg Ser Phe	405
Tyr Gly Ser Asn	410	Ala Phe Leu Thr	415	Leu Ala Phe Pro Ser	420
His Phe Gly Lys	425	Leu Phe Gly Leu Val	430	Met Ala Leu Ser Ala	435
Val Ser Leu Leu	440	Gln Phe Pro Ile Phe	445	Thr Leu Ile Lys Gly	450
Leu Gln Asn Asp	455	Pro Phe Tyr Val Asn	460	Val Met Phe Met Leu	465
Ile Leu Leu Thr	470	Phe Phe His Pro Phe	475	Leu Val Tyr Arg Glu	480
Arg Thr Trp Lys	485	Glu Ser Pro Ser Ala	490	Ile Ala	

<210> 13

<211> 580

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 977658CD1

<400> 13

Met Thr Ala Pro	1	Ala Gly Pro Arg Gly	5	Ser Glu Thr Glu Arg	10	Leu	15
Leu Thr Pro Asn	20	Pro Gly Tyr Gly Thr	25	Gln Ala Gly Pro Ser	30	Pro	35
Ala Pro Pro Thr	35	Pro Pro Glu Glu Glu	40	Asp Leu Arg Arg Arg	45	Leu	50
Lys Tyr Phe Phe	50	Met Ser Pro Cys Asp	55	Lys Phe Arg Ala Lys	60	Gly	65
Arg Lys Pro Cys	65	Lys Leu Met Leu Gln	70	Val Val Lys Ile Leu	75	Val	80
Val Thr Val Gln	80	Leu Ile Leu Phe Gly	85	Leu Ser Asn Gln Leu	90	Ala	95
Val Thr Phe Arg	95	Glu Glu Asn Thr Ile	100	Ala Phe Arg His Leu	105	Phe	110
Leu Leu Gly Tyr	110	Ser Asp Gly Ala Asp	115	Thr Phe Ala Ala Tyr	120		125
Thr Arg Glu Gln	125	Leu Tyr Gln Ala Ile	130	Phe His Ala Val Asp	135	Gln	140
Tyr Leu Ala Leu	140	Pro Asp Val Ser Leu	145	Gly Arg Tyr Ala Tyr	150	Val	155
Arg Gly Gly Gly	155	Asp Pro Trp Thr Asn	160	Gly Ser Gly Leu Ala	165	Leu	

Cys	Gln	Arg	Tyr	Tyr	His	Arg	Gly	His	Val	Asp	Pro	Ala	Asn	Asp
				170					175					180
Thr	Phe	Asp	Ile	Asp	Pro	Met	Val	Val	Thr	Asp	Cys	Ile	Gln	Val
				185					190					195
Asp	Pro	Pro	Glu	Arg	Pro	Pro	Pro	Pro	Pro	Ser	Asp	Asp	Leu	Thr
				200					205					210
Leu	Leu	Glu	Ser	Ser	Ser	Tyr	Lys		Asn	Leu	Thr	Leu	Lys	Phe
				215					220					225
His	Lys	Leu	Val	Asn	Val	Thr	Ile	His	Phe	Arg	Leu	Lys	Thr	Ile
				230					235					240
Asn	Leu	Gln	Ser	Leu	Ile	Asn	Asn	Glu	Ile	Pro	Asp	Cys	Tyr	Thr
				245					250					255
Phe	Ser	Val	Leu	Ile	Thr	Phe	Asp	Asn	Lys	Ala	His	Ser	Gly	Arg
				260					265					270
Ile	Pro	Ile	Ser	Leu	Glu	Thr	Gln	Ala	His	Ile	Gln	Glu	Cys	Lys
				275					280					285
His	Pro	Ser	Val	Phe	Gln	His	Gly	Asp	Asn	Ser	Phe	Arg	Leu	Leu
				290					295					300
Phe	Asp	Val	Val	Val	Ile	Leu	Thr	Cys	Ser	Leu	Ser	Phe	Leu	Leu
				305					310					315
Cys	Ala	Arg	Ser	Leu	Leu	Arg	Gly	Phe	Leu	Leu	Gln	Asn	Glu	Phe
				320					325					330
Val	Gly	Phe	Met	Trp	Arg	Gln	Arg	Gly	Arg	Val	Ile	Ser	Leu	Trp
				335					340					345
Glu	Arg	Leu	Glu	Phe	Val	Asn	Gly	Trp	Tyr	Ile	Leu	Leu	Val	Thr
				350					355					360
Ser	Asp	Val	Leu	Thr	Ile	Ser	Gly	Thr	Ile	Met	Lys	Ile	Gly	Ile
				365					370					375
Glu	Ala	Lys	Asn	Leu	Ala	Ser	Tyr	Asp	Val	Cys	Ser	Ile	Leu	Leu
				380					385					390
Gly	Thr	Ser	Thr	Leu	Leu	Val	Trp	Val	Gly	Val	Ile	Arg	Tyr	Leu
				395					400					405
Thr	Phe	Phe	His	Asn	Tyr	Asn	Ile	Leu	Ile	Ala	Thr	Leu	Arg	Val
				410					415					420
Ala	Leu	Pro	Ser	Val	Met	Arg	Phe	Cys	Cys	Cys	Val	Ala	Val	Ile
				425					430					435
Tyr	Leu	Gly	Tyr	Cys	Phe	Cys	Gly	Trp	Ile	Val	Leu	Gly	Pro	Tyr
				440					445					450
His	Val	Lys	Phe	Arg	Ser	Leu	Ser	Met	Val	Ser	Glu	Cys	Leu	Phe
				455					460					465
Ser	Leu	Ile	Asn	Gly	Asp	Asp	Met	Phe	Val	Thr	Phe	Ala	Ala	Met
				470					475					480
Gln	Ala	Gln	Gln	Gly	Arg	Ser	Ser	Leu	Val	Trp	Leu	Phe	Ser	Gln
				485					490					495
Leu	Tyr	Leu	Tyr	Ser	Phe	Ile	Ser	Leu	Phe	Ile	Tyr	Met	Val	Leu
				500					505					510
Ser	Leu	Phe	Ile	Ala	Leu	Ile	Thr	Gly	Ala	Tyr	Asp	Thr	Ile	Lys
				515					520					525
His	Pro	Gly	Gly	Ala	Gly	Ala	Glu	Glu	Ser	Glu	Leu	Gln	Ala	Tyr
				530					535					540
Ile	Ala	Gln	Cys	Gln	Asp	Ser	Pro	Thr	Ser	Gly	Lys	Phe	Arg	Arg
				545					550					555
Gly	Ser	Gly	Ser	Ala	Cys	Ser	Leu	Leu	Cys	Cys	Cys	Gly	Arg	Asp
				560					565					570
Pro	Ser	Glu	Glu	His	Ser	Leu	Leu	Val	Asn					
				575					580					

<210> 14
 <211> 455
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature

<223> Incyte ID No: 1004703CD1

<400> 14

Met	Ser	Phe	Leu	Ile	Asp	Ser	Ser	Ile	Met	Ile	Thr	Ser	Gln	Ile	
1				5					10					15	
Leu	Phe	Phe	Gly	Phe	Gly	Trp	Leu	Phe	Phe	Met	Arg	Gln	Leu	Phe	
				20					25					30	
Lys	Asp	Tyr	Glu	Ile	Arg	Gln	Tyr	Val	Val	Gln	Val	Ile	Phe	Ser	
				35					40					45	
Val	Thr	Phe	Ala	Phe	Ser	Cys	Thr	Met	Phe	Glu	Leu	Ile	Ile	Phe	
				50					55					60	
Glu	Ile	Leu	Gly	Val	Leu	Asn	Ser	Ser	Ser	Arg	Tyr	Phe	His	Trp	
				65					70					75	
Lys	Met	Asn	Leu	Cys	Val	Ile	Leu	Leu	Ile	Leu	Val	Phe	Met	Val	
				80					85					90	
Pro	Phe	Tyr	Ile	Gly	Tyr	Phe	Ile	Val	Ser	Asn	Ile	Arg	Leu	Leu	
				95					100					105	
His	Lys	Gln	Arg	Leu	Leu	Phe	Ser	Cys	Leu	Leu	Trp	Leu	Thr	Phe	
				110					115					120	
Met	Tyr	Phe	Phe	Trp	Lys	Leu	Gly	Asp	Leu	Phe	Pro	Ile	Leu	Ser	
				125					130					135	
Pro	Lys	His	Gly	Ile	Leu	Ser	Ile	Glu	Gln	Leu	Ile	Ser	Arg	Val	
				140					145					150	
Gly	Val	Ile	Gly	Val	Thr	Leu	Met	Ala	Leu	Leu	Ser	Gly	Phe	Gly	
				155					160					165	
Ala	Val	Asn	Cys	Pro	Tyr	Thr	Tyr	Met	Ser	Tyr	Phe	Leu	Arg	Asn	
				170					175					180	
Val	Thr	Asp	Thr	Asp	Ile	Leu	Ala	Leu	Glu	Arg	Arg	Leu	Leu	Gln	
				185					190					195	
Thr	Met	Asp	Met	Ile	Ile	Ser	Lys	Lys	Lys	Arg	Met	Ala	Met	Ala	
				200					205					210	
Arg	Arg	Thr	Met	Phe	Gln	Lys	Gly	Glu	Val	His	Asn	Lys	Pro	Ser	
				215					220					225	
Gly	Phe	Trp	Gly	Met	Ile	Lys	Ser	Val	Thr	Thr	Ser	Ala	Ser	Gly	
				230					235					240	
Ser	Glu	Asn	Leu	Thr	Leu	Ile	Gln	Gln	Glu	Val	Asp	Ala	Leu	Glu	
				245					250					255	
Glu	Leu	Ser	Arg	Gln	Leu	Phe	Leu	Glu	Thr	Ala	Asp	Leu	Tyr	Ala	
				260					265					270	
Thr	Lys	Glu	Arg	Ile	Glu	Tyr	Ser	Lys	Thr	Phe	Lys	Gly	Lys	Tyr	
				275					280					285	
Phe	Asn	Phe	Leu	Gly	Tyr	Phe	Phe	Ser	Ile	Tyr	Cys	Val	Trp	Lys	
				290					295					300	
Ile	Phe	Met	Ala	Thr	Ile	Asn	Ile	Val	Phe	Asp	Arg	Val	Gly	Lys	
				305					310					315	
Thr	Asp	Pro	Val	Thr	Arg	Gly	Ile	Glu	Ile	Thr	Val	Asn	Tyr	Leu	
				320					325					330	
Gly	Ile	Gln	Phe	Asp	Val	Lys	Phe	Trp	Ser	Gln	His	Ile	Ser	Phe	
				335					340					345	
Ile	Leu	Val	Gly	Ile	Ile	Ile	Val	Thr	Ser	Ile	Arg	Gly	Leu	Leu	
				350					355					360	
Ile	Thr	Leu	Thr	Lys	Phe	Phe	Tyr	Ala	Ile	Ser	Ser	Ser	Lys	Ser	
				365					370					375	
Ser	Asn	Val	Ile	Val	Leu	Leu	Leu	Ala	Gln	Ile	Met	Gly	Met	Tyr	
				380					385					390	
Phe	Val	Ser	Ser	Val	Leu	Leu	Ile	Arg	Met	Ser	Met	Pro	Leu	Glu	
				395					400					405	
Tyr	Arg	Thr	Ile	Ile	Thr	Glu	Val	Leu	Gly	Glu	Leu	Gln	Phe	Asn	
				410					415					420	
Phe	Tyr	His	Arg	Trp	Phe	Asp	Val	Ile	Phe	Leu	Val	Ser	Ala	Leu	
				425					430					435	
Ser	Ser	Ile	Leu	Phe	Leu	Tyr	Leu	Ala	His	Lys	Gln	Ala	Pro	Glu	
				440					445					450	

Lys Gln Met Ala Pro
455

<210> 15
<211> 277
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1334051CD1

<400> 15
Met Lys Ile Ser Met Ile Asn Tyr Lys Ser Leu Leu Ala Leu Leu
1 5 10 15
Phe Ile Leu Ala Ser Trp Ile Ile Phe Thr Val Phe Gln Asn Ser
20 25 30
Thr Lys Val Trp Ser Ala Leu Asn Leu Ser Ile Ser Leu His Tyr
35 40 45
Trp Asn Asn Ser Thr Lys Ser Leu Phe Pro Lys Thr Pro Leu Ile
50 55 60
Ser Leu Lys Pro Leu Thr Glu Thr Glu Leu Arg Ile Lys Glu Ile
65 70 75
Ile Glu Lys Leu Asp Gln Gln Ile Pro Pro Arg Pro Phe Thr His
80 85 90
Val Asn Thr Thr Thr Ser Ala Thr His Ser Thr Ala Thr Ile Leu
95 100 105
Asn Pro Arg Asp Thr Tyr Cys Arg Gly Asp Gln Leu His Ile Leu
110 115 120
Leu Glu Val Arg Asp His Leu Gly Arg Arg Lys Gln Tyr Gly Gly
125 130 135
Asp Phe Leu Arg Ala Arg Met Ser Ser Pro Ala Leu Met Ala Gly
140 145 150
Ala Ser Gly Lys Val Thr Asp Phe Asn Asn Gly Thr Tyr Leu Val
155 160 165
Ser Phe Thr Leu Phe Trp Glu Gly Gln Val Ser Leu Ser Leu Leu
170 175 180
Leu Ile His Pro Ser Glu Gly Val Ser Ala Leu Trp Ser Ala Arg
185 190 195
Asn Gln Gly Tyr Asp Arg Val Ile Phe Thr Gly Gln Phe Val Asn
200 205 210
Gly Thr Ser Gln Val His Ser Glu Cys Gly Leu Ile Leu Asn Thr
215 220 225
Asn Ala Glu Leu Cys Gln Tyr Leu Asp Asn Arg Asp Gln Glu Gly
230 235 240
Phe Tyr Cys Val Arg Pro Gln His Met Pro Cys Ala Ala Leu Thr
245 250 255
His Met Tyr Ser Lys Asn Lys Lys Val Ser Tyr Leu Ser Lys Gln
260 265 270
Glu Lys Ser Leu Phe Glu Arg
275

<210> 16
<211> 647
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1336728CD1

<400> 16
Met Ala Ser Leu Val Ser Leu Glu Leu Gly Leu Leu Leu Ala Val
1 5 10 15
Leu Val Val Thr Ala Thr Ala Ser Pro Pro Ala Gly Leu Leu Ser

	20	25	30
Leu Leu Thr Ser Gly Gln Gly Ala Leu Asp Gln Glu Ala Leu Gly			
	35	40	45
Gly Leu Leu Asn Thr Leu Ala Asp Arg Val His Cys Thr Asn Gly			
	50	55	60
Pro Cys Gly Lys Cys Leu Ser Val Glu Asp Ala Leu Gly Leu Gly			
	65	70	75
Glu Pro Glu Gly Ser Gly Leu Pro Pro Gly Pro Val Leu Glu Ala			
	80	85	90
Arg Tyr Val Ala Arg Leu Ser Ala Ala Ala Val Leu Tyr Leu Ser			
	95	100	105
Asn Pro Glu Gly Thr Cys Glu Asp Thr Arg Ala Gly Leu Trp Ala			
	110	115	120
Ser His Ala Asp His Leu Leu Ala Leu Leu Glu Ser Pro Lys Ala			
	125	130	135
Leu Thr Pro Gly Leu Ser Trp Leu Leu Gln Arg Met Gln Ala Arg			
	140	145	150
Ala Ala Gly Gln Thr Pro Lys Thr Ala Cys Val Asp Ile Pro Gln			
	155	160	165
Leu Leu Glu Glu Ala Val Gly Ala Gly Ala Pro Gly Ser Ala Gly			
	170	175	180
Gly Val Leu Ala Ala Leu Leu Asp His Val Arg Ser Gly Ser Cys			
	185	190	195
Phe His Ala Leu Pro Ser Pro Gln Tyr Phe Val Asp Phe Val Phe			
	200	205	210
Gln Gln His Ser Ser Glu Val Pro Met Thr Leu Ala Glu Leu Ser			
	215	220	225
Ala Leu Met Gln Arg Leu Gly Val Gly Arg Glu Ala His Ser Asp			
	230	235	240
His Ser His Arg His Arg Gly Ala Ser Ser Arg Asp Pro Val Pro			
	245	250	255
Leu Ile Ser Ser Ser Asn Ser Ser Ser Val Trp Asp Thr Val Cys			
	260	265	270
Leu Ser Ala Arg Asp Val Met Ala Ala Tyr Gly Leu Ser Glu Gln			
	275	280	285
Ala Gly Val Thr Pro Glu Ala Trp Ala Gln Leu Ser Pro Ala Leu			
	290	295	300
Leu Gln Gln Gln Leu Ser Gly Ala Cys Thr Ser Gln Ser Arg Pro			
	305	310	315
Pro Val Gln Asp Gln Leu Ser Gln Ser Glu Arg Tyr Leu Tyr Gly			
	320	325	330
Ser Leu Ala Thr Leu Leu Ile Cys Leu Cys Ala Val Phe Gly Leu			
	335	340	345
Leu Leu Leu Thr Cys Thr Gly Cys Arg Gly Val Thr His Tyr Ile			
	350	355	360
Leu Gln Thr Phe Leu Ser Leu Ala Val Gly Ala Leu Thr Gly Asp			
	365	370	375
Ala Val Leu His Leu Thr Pro Lys Val Leu Gly Leu His Thr His			
	380	385	390
Ser Glu Glu Gly Leu Ser Pro Gln Pro Thr Trp Arg Leu Leu Ala			
	395	400	405
Met Leu Ala Gly Leu Tyr Ala Phe Phe Leu Phe Glu Asn Leu Phe			
	410	415	420
Asn Leu Leu Leu Pro Arg Asp Pro Glu Asp Leu Glu Asp Gly Pro			
	425	430	435
Cys Gly His Ser Ser His Ser His Gly Gly His Ser His Gly Val			
	440	445	450
Ser Leu Gln Leu Ala Pro Ser Glu Leu Arg Gln Pro Lys Pro Pro			
	455	460	465
His Glu Gly Ser Arg Ala Asp Leu Val Ala Glu Glu Ser Pro Glu			
	470	475	480
Leu Leu Asn Pro Glu Pro Arg Arg Leu Ser Pro Glu Leu Arg Leu			
	485	490	495

```

Leu Pro Tyr Met Ile Thr Leu Gly Asp Ala Val His Asn Phe Ala
500 505 510
Asp Gly Leu Ala Val Gly Ala Ala Phe Ala Ser Ser Trp Lys Thr
515 520 525
Gly Leu Ala Thr Ser Leu Ala Val Phe Cys His Glu Leu Pro His
530 535 540
Glu Leu Gly Asp Phe Ala Ala Leu Leu His Ala Gly Leu Ser Val
545 550 555
Arg Gln Ala Leu Leu Leu Asn Leu Ala Ser Ala Leu Thr Ala Phe
560 565 570
Ala Gly Leu Tyr Val Ala Leu Ala Val Gly Val Ser Glu Glu Ser
575 580 585
Glu Ala Trp Ile Leu Ala Val Ala Thr Gly Leu Phe Leu Tyr Val
590 595 600
Ala Leu Cys Asp Met Leu Pro Ala Met Leu Lys Val Arg Asp Pro
605 610 615
Arg Pro Trp Leu Leu Phe Leu Leu His Asn Val Gly Leu Leu Gly
620 625 630
Gly Trp Thr Val Leu Leu Leu Leu Ser Leu Tyr Glu Asp Asp Ile
635 640 645
Thr Phe

```

```

<210> 17
<211> 406
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 1452856CD1

```

```

<400> 17
Met Ala Glu Asn Gly Lys Asn Cys Asp Gln Arg Arg Val Ala Met
1 5 10 15
Asn Lys Glu His His Asn Gly Asn Phe Thr Asp Pro Ser Ser Val
20 25 30
Asn Glu Lys Lys Arg Arg Glu Arg Glu Glu Arg Gln Asn Ile Val
35 40 45
Leu Trp Arg Gln Pro Leu Ile Thr Leu Gln Tyr Phe Ser Leu Glu
50 55 60
Ile Leu Val Ile Leu Lys Glu Trp Thr Ser Lys Leu Trp His Arg
65 70 75
Gln Ser Ile Val Val Ser Phe Leu Leu Leu Leu Ala Val Leu Ile
80 85 90
Ala Thr Tyr Tyr Val Glu Gly Val His Gln Gln Tyr Val Gln Arg
95 100 105
Ile Glu Lys Gln Phe Leu Leu Tyr Ala Tyr Trp Ile Gly Leu Gly
110 115 120
Ile Leu Ser Ser Val Gly Leu Gly Thr Gly Leu His Thr Phe Leu
125 130 135
Leu Tyr Leu Gly Pro His Ile Ala Ser Val Thr Leu Ala Ala Tyr
140 145 150
Glu Cys Asn Ser Val Asn Phe Pro Glu Pro Pro Tyr Pro Asp Gln
155 160 165
Ile Ile Cys Pro Asp Glu Glu Gly Thr Glu Gly Thr Ile Ser Leu
170 175 180
Trp Ser Ile Ile Ser Lys Val Arg Ile Glu Ala Cys Met Trp Gly
185 190 195
Ile Gly Thr Ala Ile Gly Glu Leu Pro Pro Tyr Phe Met Ala Arg
200 205 210
Ala Ala Arg Leu Ser Gly Ala Glu Pro Asp Asp Glu Glu Tyr Gln
215 220 225
Glu Phe Glu Glu Met Leu Glu His Ala Glu Ser Ala Gln Asp Phe

```

Ala Ser Arg Ala	230	Lys Leu Ala Val Gln	235	Lys Leu Val Gln Lys	240
	245		250		255
Gly Phe Phe Gly	260	Ile Leu Ala Cys Ala	265	Ser Ile Pro Asn Pro	270
	275		280		285
Phe Asp Leu Ala	290	Gly Ile Thr Cys Gly	295	His Phe Leu Val Pro	300
	305		310		315
Trp Thr Phe Phe	320	Gly Ala Thr Leu Ile	325	Gly Lys Ala Ile Ile	330
	335		340		345
Met His Ile Gln	350	Lys Ile Phe Val Ile	355	Ile Thr Phe Ser Lys	360
	365		370		375
Ile Val Glu Gln	380	Met Val Ala Phe Ile	385	Gly Ala Val Pro Gly	390
	395		400		405
Gly Pro Ser Leu		Gln Lys Pro Phe Gln		Glu Tyr Leu Glu Ala	
Arg Gln Lys Leu		His His Lys Ser Glu		Met Gly Thr Pro Gln	
Glu Asn Trp Leu		Ser Trp Met Phe Glu		Lys Leu Val Val Val	
Val Cys Tyr Phe		Ile Leu Ser Ile Ile		Asn Ser Met Ala Gln	
Tyr Ala Lys Arg		Ile Gln Gln Arg Leu		Asn Ser Glu Glu Lys	
Lys					

<210> 18
 <211> 290
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1562471CD1

<400> 18	
Met Pro Leu Leu Thr	Leu Tyr Leu Leu Leu Phe Trp Leu Ser Gly
1	5
Tyr Ser Ile Ala Thr	Gln Ile Thr Gly Pro Thr Thr Val Asn Gly
20	25
Leu Glu Arg Gly Ser	Leu Thr Val Gln Cys Val Tyr Arg Ser Gly
35	40
Trp Glu Thr Tyr Leu	Lys Trp Trp Cys Arg Gly Ala Ile Trp Arg
50	55
Asp Cys Lys Ile Leu	Val Lys Thr Ser Gly Ser Glu Gln Glu Val
65	70
Lys Arg Asp Arg Val	Ser Ile Lys Asp Asn Gln Lys Asn Arg Thr
80	85
Phe Thr Val Thr Met	Glu Asp Leu Met Lys Thr Asp Ala Asp Thr
95	100
Tyr Trp Cys Gly Ile	Glu Lys Thr Gly Asn Asp Leu Gly Val Thr
110	115
Val Gln Val Thr Ile	Asp Pro Ala Pro Val Thr Gln Glu Glu Thr
125	130
Ser Ser Ser Pro Thr	Leu Thr Gly His His Leu Asp Asn Arg His
140	145
Lys Leu Leu Lys Leu	Ser Val Leu Leu Pro Leu Ile Phe Thr Ile
155	160
Leu Leu Leu Leu Leu	Val Ala Ala Ser Leu Leu Ala Trp Arg Met
170	175
Met Lys Tyr Gln Gln	Lys Ala Ala Gly Met Ser Pro Glu Gln Val
185	190
Leu Gln Pro Leu Glu	Gly Asp Leu Cys Tyr Ala Asp Leu Thr Leu
200	205
	210

Gln	Leu	Ala	Gly	Thr	Ser	Pro	Arg	Lys	Ala	Thr	Thr	Lys	Leu	Ser
				215					220					225
Ser	Ala	Gln	Val	Asp	Gln	Val	Glu	Val	Glu	Tyr	Val	Thr	Met	Ala
				230					235					240
Ser	Leu	Pro	Lys	Glu	Asp	Ile	Ser	Tyr	Ala	Ser	Leu	Thr	Leu	Gly
				245					250					255
Ala	Glu	Asp	Gln	Glu	Pro	Thr	Tyr	Cys	Asn	Met	Gly	His	Leu	Ser
				260					265					270
Ser	His	Leu	Pro	Gly	Arg	Gly	Pro	Glu	Glu	Pro	Thr	Glu	Tyr	Ser
				275					280					285
Thr	Ile	Ser	Arg	Pro										
				290										

<210> 19

<211> 390

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1618158CD1

<400> 19

Met	Phe	Ser	Thr	Asn	Tyr	Ser	His	Met	Glu	Asn	Tyr	Arg	Lys	Arg
1				5					10					15
Glu	Asp	Leu	Val	Tyr	Gln	Ser	Thr	Val	Arg	Leu	Pro	Glu	Val	Arg
				20					25					30
Ile	Ser	Asp	Asn	Gly	Pro	Tyr	Glu	Cys	His	Val	Gly	Ile	Tyr	Asp
				35					40					45
Arg	Ala	Thr	Arg	Glu	Lys	Val	Val	Leu	Ala	Ser	Gly	Asn	Ile	Phe
				50					55					60
Leu	Asn	Val	Met	Ala	Pro	Pro	Thr	Ser	Ile	Glu	Val	Val	Ala	Ala
				65					70					75
Asp	Thr	Pro	Ala	Pro	Phe	Ser	Arg	Tyr	Gln	Ala	Gln	Asn	Phe	Thr
				80					85					90
Leu	Val	Cys	Ile	Val	Ser	Gly	Gly	Lys	Pro	Ala	Pro	Met	Val	Tyr
				95					100					105
Phe	Lys	Arg	Asp	Gly	Glu	Pro	Ile	Asp	Ala	Val	Pro	Leu	Ser	Glu
				110					115					120
Pro	Pro	Ala	Ala	Ser	Ser	Gly	Pro	Leu	Gln	Asp	Ser	Arg	Pro	Phe
				125					130					135
Arg	Ser	Leu	Leu	His	Arg	Asp	Leu	Asp	Asp	Thr	Lys	Met	Gln	Lys
				140					145					150
Ser	Leu	Ser	Leu	Leu	Asp	Ala	Glu	Asn	Arg	Gly	Gly	Arg	Pro	Tyr
				155					160					165
Thr	Glu	Arg	Pro	Ser	Arg	Gly	Leu	Thr	Pro	Asp	Pro	Asn	Ile	Leu
				170					175					180
Leu	Gln	Pro	Thr	Thr	Glu	Asn	Ile	Pro	Glu	Thr	Val	Val	Ser	Arg
				185					190					195
Glu	Phe	Pro	Arg	Trp	Val	His	Ser	Ala	Glu	Pro	Thr	Tyr	Phe	Leu
				200					205					210
Arg	His	Ser	Arg	Thr	Pro	Ser	Ser	Asp	Gly	Thr	Val	Glu	Val	Arg
				215					220					225
Ala	Leu	Leu	Thr	Trp	Thr	Leu	Asn	Pro	Gln	Ile	Asp	Asn	Glu	Ala
				230					235					240
Leu	Phe	Ser	Cys	Glu	Val	Lys	His	Pro	Ala	Leu	Ser	Met	Pro	Met
				245					250					255
Gln	Ala	Glu	Val	Thr	Leu	Val	Ala	Pro	Lys	Gly	Pro	Lys	Ile	Val
				260					265					270
Met	Thr	Pro	Ser	Arg	Ala	Arg	Val	Gly	Asp	Thr	Val	Arg	Ile	Leu
				275					280					285
Val	His	Gly	Phe	Gln	Asn	Glu	Val	Phe	Pro	Glu	Pro	Met	Phe	Thr
				290					295					300
Trp	Thr	Arg	Val	Gly	Ser	Arg	Leu	Leu	Asp	Gly	Ser	Ala	Glu	Phe

305	310	315
Asp Gly Lys Glu Leu Val Leu Glu Arg Val Pro Ala Glu Leu Asn		
320	325	330
Gly Ser Met Tyr Arg Cys Thr Ala Gln Asn Pro Leu Gly Ser Thr		
335	340	345
Asp Thr His Thr Arg Leu Ile Val Phe Glu Asn Pro Asn Ile Pro		
350	355	360
Arg Gly Thr Glu Asp Ser Asn Gly Ser Ile Gly Pro Thr Gly Ala		
365	370	375
Arg Leu Thr Leu Val Leu Ala Leu Thr Val Ile Leu Glu Leu Thr		
380	385	390

<210> 20
 <211> 427
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1656935CD1

<400> 20

Met Asn Val Asn Ser Met Asp Met Thr Gly Gly Leu Ser Val Lys	
1 5 10 15	
Asp Pro Ser Gln Ser Gln Ser Arg Leu Pro Gln Trp Thr His Pro	
20 25 30	
Asn Ser Met Asp Asn Leu Pro Ser Ala Ala Ser Pro Leu Glu Gln	
35 40 45	
Asn Pro Ser Lys His Gly Ala Ile Pro Gly Gly Leu Ser Ile Gly	
50 55 60	
Pro Pro Gly Lys Ser Ser Ile Asp Asp Ser Tyr Gly Arg Tyr Asp	
65 70 75	
Leu Ile Gln Asn Ser Glu Ser Pro Ala Ser Pro Pro Val Ala Val	
80 85 90	
Pro His Ser Trp Ser Arg Ala Lys Ser Asp Ser Asp Lys Ile Ser	
95 100 105	
Asn Gly Ser Ser Ile Asn Trp Pro Pro Glu Phe His Pro Gly Val	
110 115 120	
Pro Trp Lys Gly Leu Gln Asn Ile Asp Pro Glu Asn Asp Pro Asp	
125 130 135	
Val Thr Pro Gly Ser Val Pro Thr Gly Pro Thr Ile Asn Thr Thr	
140 145 150	
Ile Gln Asp Val Asn Arg Tyr Leu Leu Lys Ser Gly Gly Ser Ser	
155 160 165	
Pro Pro Ser Ser Gln Asn Ala Thr Leu Pro Ser Ser Ser Ala Trp	
170 175 180	
Pro Leu Ser Ala Ser Gly Tyr Ser Ser Ser Phe Ser Ser Ile Ala	
185 190 195	
Ser Ala Pro Ser Val Ala Gly Lys Leu Ser Asp Ile Lys Ser Thr	
200 205 210	
Trp Ser Ser Gly Pro Thr Ser His Thr Gln Ala Ser Leu Ser His	
215 220 225	
Glu Leu Trp Lys Val Pro Arg Asn Ser Thr Ala Pro Thr Arg Pro	
230 235 240	
Pro Pro Gly Leu Thr Asn Pro Lys Pro Ser Ser Thr Trp Gly Ala	
245 250 255	
Ser Pro Leu Gly Trp Thr Ser Ser Tyr Ser Ser Gly Ser Ala Trp	
260 265 270	
Ser Thr Asp Thr Ser Gly Arg Thr Ser Ser Trp Leu Val Leu Arg	
275 280 285	
Asn Leu Thr Pro Gln Ile Asp Gly Ser Lys Leu Arg Thr Leu Cys	
290 295 300	
Leu Gln His Gly Pro Leu Ile Thr Phe His Leu Asn Leu Thr Gln	

	305		310		315
Gly Asn Ala Val	Val Arg Tyr Ser Ser	Lys Glu Glu Gly Leu	Pro		
	320		325		330
Lys Ala Gln Glu	Val Leu Cys Thr Ile	Val Arg Pro Trp Glu	Thr		
	335		340		345
Leu Ser His Ser	Leu Gly Pro Ser Phe	Arg Leu Val Gly Thr	Lys		
	350		355		360
Glu Val Gly Ile	Arg Val Ser Phe Lys	Pro Pro Glu Gly Pro	Gly		
	365		370		375
Arg Ile Gly Gln	Ser Thr Ile Phe Gln	Gly Leu Ala Gln Phe	His		
	380		385		390
Asp Gln Arg Gly	Val Ser Lys Leu Thr	Gly Arg Gly Gly Ile	His		
	395		400		405
Arg Pro Arg Gly	Arg Gly Lys Ala Ser	His Gln Leu Ala His	Met		
	410		415		420
Arg His Cys Glu	Leu Thr Phe				
	425				

<210> 21
 <211> 459
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1859305CD1

<400> 21

Met Glu Lys Thr Cys	Ile Asp Ala Leu Pro	Leu Thr Met Asn Ser
1	5	10
Ser Glu Lys Gln Glu	Thr Val Cys Ile Phe	Gly Thr Gly Asp Phe
	20	25
Gly Arg Ser Leu Gly	Leu Lys Met Leu Gln	Cys Gly Tyr Ser Val
	35	40
Val Phe Gly Ser Arg	Asn Pro Gln Lys Thr	Thr Leu Leu Pro Ser
	50	55
Gly Ala Glu Val Leu	Ser Tyr Ser Glu Ala	Ala Lys Lys Ser Asp
	65	70
Ile Ile Ile Ile Ala	Ile His Arg Glu His	Tyr Asp Phe Leu Thr
	80	85
Glu Leu Thr Glu Val	Leu Asn Gly Lys Ile	Leu Val Asp Ile Ser
	95	100
Asn Asn Leu Lys Ile	Asn Gln Tyr Pro Glu	Ser Asn Ala Glu Tyr
	110	115
Leu Ala His Leu Val	Pro Gly Ala His Val	Val Lys Ala Phe Asn
	125	130
Thr Ile Ser Ala Trp	Ala Leu Gln Ser Gly	Ala Leu Asp Ala Ser
	140	145
Arg Gln Val Phe Val	Cys Gly Asn Asp Ser	Lys Ala Lys Gln Arg
	155	160
Val Met Asp Ile Val	Arg Asn Leu Gly Leu	Thr Pro Met Asp Gln
	170	175
Gly Ser Leu Met Ala	Ala Lys Glu Ile Glu	Lys Tyr Pro Leu Gln
	185	190
Leu Phe Pro Met Trp	Arg Phe Pro Phe Tyr	Leu Ser Ala Val Leu
	200	205
Cys Val Phe Leu Phe	Phe Tyr Cys Val Ile	Arg Asp Val Ile Tyr
	215	220
Pro Tyr Val Tyr Glu	Lys Lys Asp Asn Thr	Phe Arg Met Ala Ile
	230	235
Ser Ile Pro Asn Arg	Ile Phe Pro Ile Thr	Ala Leu Thr Leu Leu
	245	250
Ala Leu Val Tyr Leu	Pro Gly Val Ile Ala	Ala Ile Leu Gln Leu
	260	265

Tyr Arg Gly Thr Lys Tyr Arg Arg Phe Pro Asp Trp Leu Asp His
 275 280
 Trp Met Leu Cys Arg Lys Gln Leu Gly Leu Val Ala Leu Gly Phe
 290 295 300
 Ala Phe Leu His Val Leu Tyr Thr Leu Val Ile Pro Ile Arg Tyr
 305 310 315
 Tyr Val Arg Trp Arg Leu Gly Asn Leu Thr Val Thr Gln Ala Ile
 320 325 330
 Leu Lys Lys Glu Asn Pro Phe Ser Thr Ser Ser Ala Trp Leu Ser
 335 340 345
 Asp Ser Tyr Val Ala Leu Gly Ile Leu Gly Phe Phe Leu Phe Val
 350 355 360
 Leu Leu Gly Ile Thr Ser Leu Pro Ser Val Ser Asn Ala Val Asn
 365 370 375
 Trp Arg Glu Phe Arg Phe Val Gln Ser Lys Leu Gly Tyr Leu Thr
 380 385 390
 Leu Ile Leu Cys Thr Ala His Thr Leu Val Tyr Gly Gly Lys Arg
 395 400 405
 Phe Leu Ser Pro Ser Asn Leu Arg Trp Tyr Leu Pro Ala Ala Tyr
 410 415 420
 Val Leu Gly Leu Ile Ile Pro Cys Thr Val Leu Val Ile Lys Phe
 425 430 435
 Val Leu Ile Met Pro Cys Val Asp Asn Thr Leu Thr Arg Ile Arg
 440 445 450
 Gln Gly Trp Glu Arg Asn Ser Lys His
 455

<210> 22
 <211> 229
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1949083CD1

<400> 22
 Met Leu Pro Val Ser Arg Thr Cys Leu Leu Glu Ser Ser Thr Arg
 1 5 10 15
 Leu Lys Pro His Glu Ala Gln Asn Tyr Arg Lys Lys Ala Leu Trp
 20 25 30
 Val Ser Trp Phe Ser Ile Ile Val Thr Leu Ala Leu Ala Val Ala
 35 40 45
 Ala Phe Thr Val Ser Val Met Arg Tyr Ser Ala Ser Ala Phe Gly
 50 55 60
 Phe Ala Phe Asp Ala Ile Leu Asp Val Leu Ser Ser Ala Ile Val
 65 70 75
 Leu Trp Arg Tyr Ser Asn Ala Ala Ala Val His Ser Ala His Arg
 80 85 90
 Glu Tyr Ile Ala Cys Val Ile Leu Gly Val Ile Phe Leu Leu Ser
 95 100 105
 Ser Ile Cys Ile Val Val Lys Ala Ile His Asp Leu Ser Thr Arg
 110 115 120
 Leu Leu Pro Glu Val Asp Asp Phe Leu Phe Ser Val Ser Ile Leu
 125 130 135
 Ser Gly Ile Leu Cys Ser Ile Leu Ala Val Leu Lys Phe Met Leu
 140 145 150
 Gly Lys Val Leu Thr Ser Arg Ala Leu Ile Thr Asp Gly Phe Asn
 155 160 165
 Ser Leu Val Gly Gly Val Met Gly Phe Ser Ile Leu Leu Ser Ala
 170 175 180
 Glu Val Phe Lys His Asp Ser Ala Val Trp Tyr Leu Asp Gly Ser
 185 190 195
 Ile Gly Val Leu Ile Gly Leu Thr Ile Phe Ala Tyr Gly Val Lys

	200		205		210
Leu Leu Ile Asp Met Val Pro Lys Val Arg Gln Thr Arg His Tyr					
	215		220		225
Glu Met Phe Glu					

<210> 23
 <211> 311
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1996357CD1

<400> 23
 Met Ala Val Asp Ile Gln Pro Ala Cys Leu Gly Leu Tyr Cys Gly
 1 5 10 15
 Lys Thr Leu Leu Phe Lys Asn Gly Ser Thr Glu Ile Tyr Gly Glu
 20 25 30
 Cys Gly Val Cys Pro Arg Gly Gln Arg Thr Asn Ala Gln Lys Tyr
 35 40 45
 Cys Gln Pro Cys Thr Glu Ser Pro Glu Leu Tyr Asp Trp Leu Tyr
 50 55 60
 Leu Gly Phe Met Ala Met Leu Pro Leu Val Leu His Trp Phe Phe
 65 70 75
 Ile Glu Trp Tyr Ser Gly Lys Lys Ser Ser Ser Ala Leu Phe Gln
 80 85 90
 His Ile Thr Ala Leu Phe Glu Cys Ser Met Ala Ala Ile Ile Thr
 95 100 105
 Leu Leu Val Ser Asp Pro Val Gly Val Leu Tyr Ile Arg Ser Cys
 110 115 120
 Arg Val Leu Met Leu Ser Asp Trp Tyr Thr Met Leu Tyr Asn Pro
 125 130 135
 Ser Pro Asp Tyr Val Thr Thr Val His Cys Thr His Glu Ala Val
 140 145 150
 Tyr Pro Leu Tyr Thr Ile Val Phe Ile Tyr Tyr Ala Phe Cys Leu
 155 160 165
 Val Leu Met Met Leu Leu Arg Pro Leu Leu Val Lys Lys Ile Ala
 170 175 180
 Cys Gly Leu Gly Lys Ser Asp Arg Phe Lys Ser Ile Tyr Ala Ala
 185 190 195
 Leu Tyr Phe Phe Pro Ile Leu Thr Val Leu Gln Ala Val Gly Gly
 200 205 210
 Gly Leu Leu Tyr Tyr Ala Phe Pro Tyr Ile Ile Leu Val Leu Ser
 215 220 225
 Leu Val Thr Leu Ala Val Tyr Met Ser Ala Ser Glu Ile Glu Asn
 230 235 240
 Cys Tyr Asp Leu Leu Val Arg Lys Lys Arg Leu Ile Val Leu Phe
 245 250 255
 Ser His Trp Leu Leu His Ala Tyr Gly Ile Ile Ser Ile Ser Arg
 260 265 270
 Val Asp Lys Leu Glu Gln Asp Leu Pro Leu Leu Ala Leu Val Pro
 275 280 285
 Thr Pro Ala Leu Phe Tyr Leu Phe Thr Ala Lys Phe Thr Glu Pro
 290 295 300
 Ser Arg Ile Leu Ser Glu Gly Ala Asn Gly His
 305 310

<210> 24
 <211> 92
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2061330CD1

<400> 24

```

Met Arg Phe Ile Phe Leu Lys Phe Trp Thr Tyr Thr Val Arg Ala
 1          5          10          15
Ser Thr Asp Leu Thr Gln Thr Gly Asp Cys Ser Gln Cys Thr His
          20          25          30
Gln Val Thr Glu Val Gly Gln Gln Ile Lys Thr Ile Phe Leu Phe
          35          40          45
Tyr Ser Tyr Tyr Glu Cys Met Glu Thr Ile Lys Glu Thr Cys Leu
          50          55          60
Tyr Asn Ala Thr Gln Tyr Lys Val Cys Ser Pro Arg Asn Asp Arg
          65          70          75
Pro Asp Val Cys Tyr Asn Pro Ser Glu Pro Pro Ala Pro Pro Phe
          80          85          90
Leu Lys

```

<210> 25

<211> 258

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2346947CD1

<400> 25

```

Met Ala Glu Ser Pro Gly Cys Cys Ser Val Trp Ala Arg Cys Leu
 1          5          10          15
His Cys Leu Tyr Ser Cys His Trp Arg Lys Cys Pro Arg Glu Arg
          20          25          30
Met Gln Thr Ser Lys Cys Asp Cys Ile Trp Phe Gly Leu Leu Phe
          35          40          45
Leu Thr Phe Leu Leu Ser Leu Ser Trp Leu Tyr Ile Gly Leu Val
          50          55          60
Leu Leu Asn Asp Leu His Asn Phe Asn Glu Phe Leu Phe Arg Arg
          65          70          75
Trp Gly His Trp Met Asp Trp Ser Leu Ala Phe Leu Leu Val Ile
          80          85          90
Ser Leu Leu Val Thr Tyr Ala Ser Leu Leu Leu Val Leu Ala Leu
          95          100          105
Leu Leu Arg Leu Cys Arg Gln Pro Leu His Leu His Ser Leu His
          110          115          120
Lys Val Leu Leu Leu Leu Ile Met Leu Leu Val Ala Ala Gly Leu
          125          130          135
Val Gly Leu Asp Ile Gln Trp Gln Gln Glu Trp His Ser Leu Arg
          140          145          150
Val Ser Leu Gln Ala Thr Ala Pro Phe Leu His Ile Gly Ala Ala
          155          160          165
Ala Gly Ile Ala Leu Leu Ala Trp Pro Val Ala Asp Thr Phe Tyr
          170          175          180
Arg Ile His Arg Arg Gly Pro Lys Ile Leu Leu Leu Leu Leu Phe
          185          190          195
Phe Gly Val Val Leu Val Ile Tyr Leu Ala Pro Leu Cys Ile Ser
          200          205          210
Ser Pro Cys Ile Met Glu Pro Arg Asp Leu Pro Pro Lys Pro Gly
          215          220          225
Leu Val Gly His Arg Gly Ala Pro Met Leu Ala Pro Glu Asn Thr
          230          235          240
Leu Met Ser Leu Arg Lys Thr Ala Glu Cys Gly Leu Leu Cys Leu
          245          250          255
Arg Leu Met

```

```
<220>  
<221> misc_feature  
<223> Incyte ID No: 2795577CD1
```

```
<210> 27
<211> 136
<212> PRT
<213> Homo sapiens
```

```
<220>  
<221> misc_feature  
<223> Incyte ID No: 3255825CD1
```

<400> 27														
Met	Ile	Ser	Ile	Thr	Glu	Trp	Gln	Lys	Ile	Gly	Val	Gly	Ile	Thr
1				5					10					15
Gly	Phe	Gly	Ile	Phe	Phe	Ile	Leu	Phe	Gly	Thr	Leu	Leu	Tyr	Phe
				20					25					30
Asp	Ser	Val	Leu	Leu	Ala	Phe	Gly	Asn	Leu	Leu	Phe	Leu	Thr	Gly
				35					40					45
Leu	Ser	Leu	Ile	Ile	Gly	Leu	Arg	Lys	Thr	Phe	Trp	Phe	Phe	Phe
				50					55					60
Gln	Arg	His	Lys	Leu	Lys	Gly	Thr	Ser	Phe	Leu	Leu	Gly	Gly	Val
				65					70					75

Val	Ile	Val	Leu	Leu	Arg	Trp	Pro	Leu	Leu	Gly	Met	Phe	Leu	Glu	
				80						85				90	
Thr	Tyr	Gly	Phe	Phe	Ser	Leu	Phe	Lys	Gly	Phe	Phe	Pro	Val	Ala	
				95					100					105	
Phe	Gly	Ser	Trp	Ala	Met	Ser	Ala	Thr	Ser	Pro	Ser	Trp	Val	Arg	
				110					115					120	
Cys	Ser	Gly	Asp	Phe	Lys	Ala	Leu	Ala	Arg	Trp	Ser	Glu	Lys	Gln	
				125					130					135	

Arg

<210> 28

<211> 458

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3393430CD1

<400> 28

Met	Ala	Trp	Ala	Ser	Arg	Leu	Gly	Leu	Leu	Leu	Ala	Leu	Leu	Leu	
1				5					10					15	
Pro	Val	Val	Gly	Ala	Ser	Thr	Pro	Gly	Thr	Val	Val	Arg	Leu	Asn	
				20					25					30	
Lys	Ala	Ala	Leu	Ser	Tyr	Val	Ser	Glu	Ile	Gly	Lys	Ala	Pro	Leu	
				35					40					45	
Gln	Arg	Ala	Leu	Gln	Val	Thr	Val	Pro	His	Phe	Leu	Asp	Trp	Ser	
				50					55					60	
Gly	Glu	Ala	Leu	Gln	Pro	Thr	Arg	Ile	Arg	Ile	Leu	Asn	Val	His	
				65					70					75	
Val	Pro	Arg	Leu	His	Leu	Lys	Phe	Ile	Ala	Gly	Phe	Gly	Val	Arg	
				80					85					90	
Leu	Leu	Ala	Ala	Ala	Asn	Phe	Thr	Phe	Lys	Val	Phe	Arg	Ala	Pro	
				95					100					105	
Glu	Pro	Leu	Glu	Leu	Thr	Leu	Pro	Val	Glu	Leu	Leu	Ala	Asp	Thr	
				110					115					120	
Arg	Val	Thr	Gln	Ser	Ser	Ile	Arg	Thr	Pro	Val	Val	Ser	Ile	Ser	
				125					130					135	
Ala	Cys	Ser	Leu	Phe	Ser	Gly	His	Ala	Asn	Glu	Phe	Asp	Gly	Ser	
				140					145					150	
Asn	Ser	Thr	Ser	His	Ala	Leu	Leu	Val	Leu	Val	Gln	Lys	His	Ile	
				155					160					165	
Lys	Ala	Val	Leu	Ser	Asn	Lys	Leu	Cys	Leu	Ser	Ile	Ser	Asn	Leu	
				170					175					180	
Val	Gln	Gly	Val	Asn	Val	His	Leu	Gly	Thr	Leu	Ile	Gly	Leu	Asn	
				185					190					195	
Pro	Val	Gly	Pro	Glu	Ser	Gln	Ile	Arg	Tyr	Ser	Met	Val	Ser	Val	
				200					205					210	
Pro	Thr	Val	Thr	Ser	Asp	Tyr	Ile	Ser	Leu	Glu	Val	Asn	Ala	Val	
				215					220					225	
Leu	Phe	Leu	Leu	Gly	Lys	Pro	Ile	Ile	Leu	Pro	Thr	Asp	Ala	Thr	
				230					235					240	
Pro	Phe	Val	Leu	Pro	Arg	His	Val	Gly	Thr	Glu	Gly	Ser	Met	Ala	
				245					250					255	
Thr	Val	Gly	Leu	Ser	Gln	Gln	Leu	Phe	Asp	Ser	Ala	Leu	Leu	Leu	
				260					265					270	
Leu	Gln	Lys	Ala	Gly	Ala	Leu	Asn	Leu	Asp	Ile	Thr	Gly	Gln	Leu	
				275					280					285	
Arg	Ser	Asp	Asp	Asn	Leu	Leu	Asn	Thr	Ser	Ala	Leu	Gly	Arg	Leu	
				290					295					300	
Ile	Pro	Glu	Val	Ala	Arg	Gln	Phe	Pro	Glu	Pro	Met	Pro	Val	Val	
				305					310					315	
Leu	Lys	Val	Arg	Leu	Gly	Ala	Thr	Pro	Val	Ala	Met	Leu	His	Thr	

Asn Asn Ala Thr	320	Asn Asn Ala Thr	325	Asn Asn Ala Thr	330
Leu Arg Leu Gln Pro	335	Phe Val Glu Val Leu Ala	340	Leu Arg Leu Gln Pro	345
Thr Ala Ser Asn Ser	350	Leu Phe Ser Leu Asp Val	355	Thr Ala Ser Asn Ser	360
Val Val Asn Leu Arg	365	Val Ser Lys Val Lys Leu	370	Val Val Asn Leu Arg	375
Gln Gly Thr Thr Ser	380	Val Gln Leu Thr Val Ala	385	Gln Gly Thr Thr Ser	390
Ser Ser Asn Val Gly	395	Gln Val Arg Thr Leu	400	Ser Ser Asn Val Gly	405
Met Gly Thr Val Phe	410	Leu Asp His Leu Asn Ala	415	Met Gly Thr Val Phe	420
Leu Leu Ala Met Gly	425	Val Val Asn Leu His	430	Leu Leu Ala Met Gly	435
Tyr Val Ala Pro Glu	440	Gly Tyr Val Val Ile	445	Tyr Val Ala Pro Glu	450
Ser Ser Gly Leu Phe	455			Ser Ser Gly Leu Phe	

<210> 29
 <211> 368
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3490990CD1

<400> 29

Met Phe Gly Gln Asn	Leu Glu Val Gln	Leu Ser Ser Ala Arg	Thr
1	5	10	15
Glu Asn Thr Thr Val	Val Trp Lys Ser	Phe His Asp Ser Ile	Thr
20	25	30	
Leu Ile Val Leu Ser	Ser Glu Val Gly	Ile Ser Glu Leu Arg	Leu
35	40	45	
Glu Arg Leu Leu Gln	Met Val Phe Gly	Ala Met Val Leu Leu	Val
50	55	60	
Gly Leu Glu Glu Leu	Thr Asn Ile Arg	Asn Val Glu Arg Leu	Lys
65	70	75	
Lys Asp Leu Arg Ala	Ser Tyr Cys Leu	Ile Asp Ser Phe Leu	Gly
80	85	90	
Asp Ser Glu Leu Ile	Gly Asp Leu Thr	Gln Cys Val Asp Cys	Val
95	100	105	
Ile Pro Pro Glu Gly	Ser Leu Leu Gln	Glu Ala Leu Ser Gly	Phe
110	115	120	
Ala Glu Ala Ala Gly	Thr Thr Phe Val	Ser Leu Val Val Ser	Gly
125	130	135	
Arg Val Val Ala Ala	Thr Glu Gly Trp	Trp Arg Leu Gly Thr	Pro
140	145	150	
Glu Ala Val Leu Leu	Pro Trp Leu Val	Gly Ser Leu Pro Pro	Gln
155	160	165	
Thr Ala Arg Asp Tyr	Pro Val Tyr Leu	Pro His Gly Ser Pro	Thr
170	175	180	
Val Pro His Arg Leu	Leu Thr Leu Thr	Leu Leu Pro Ser Leu	Glu
185	190	195	
Leu Cys Leu Leu Cys	Gly Pro Ser Pro	Pro Leu Ser Gln Leu	Tyr
200	205	210	
Pro Gln Leu Leu Glu	Arg Trp Trp Gln	Pro Leu Leu Asp Pro	Leu
215	220	225	
Arg Ala Cys Leu Pro	Leu Gly Pro Arg	Ala Leu Pro Ser Gly	Phe
230	235	240	
Pro Leu His Thr Asp	Ile Leu Gly Leu	Leu Leu Leu His Leu	Glu
245	250	255	

```

Leu Lys Arg Cys Leu Phe Thr Val Glu Pro Leu Gly Asp Lys Glu
260 265 270
Pro Ser Pro Glu Gln Arg Arg Arg Leu Leu Arg Asn Phe Tyr Thr
275 280 285
Leu Val Thr Ser Thr His Phe Pro Pro Glu Pro Gly Pro Pro Glu
290 295 300
Lys Thr Glu Asp Glu Val Tyr Gln Ala Gln Leu Pro Arg Ala Cys
305 310 315
Tyr Leu Val Leu Gly Thr Glu Glu Pro Gly Thr Gly Val Arg Leu
320 325 330
Val Ala Leu Gln Leu Gly Leu Arg Arg Leu Leu Leu Leu Leu Ser
335 340 345
Pro Gln Ser Pro Thr His Gly Leu Arg Ser Leu Ala Thr His Thr
350 355 360
Leu His Ala Leu Thr Pro Leu Leu
365

```

<210> 30
 <211> 91
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3635154CD1

```

<400> 30
Met Tyr Gly Lys Ile Ile Phe Val Leu Leu Leu Ser Glu Ile Val
1 5 10 15
Ser Ile Ser Ala Ser Ser Thr Thr Gly Val Ala Met His Thr Ser
20 25 30
Thr Ser Ser Ser Val Thr Lys Ser Tyr Ile Ser Ser Gln Thr Asn
35 40 45
Gly Glu Thr Gly Gln Leu Val His Arg Phe Thr Val Pro Ala Pro
50 55 60
Val Val Ile Ile Leu Ile Ile Leu Cys Val Met Ala Gly Ile Ile
65 70 75
Gly Thr Ile Leu Leu Phe Ser Tyr Ser Phe Arg Arg Leu Ile Lys
80 85 90
Gly

```

<210> 31
 <211> 295
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4374347CD1

```

<400> 31
Met Gly Pro Pro Ser Ala Cys Pro His Arg Glu Cys Ile Pro Trp
1 5 10 15
Gln Gly Leu Leu Leu Thr Ala Ser Leu Leu Thr Phe Trp Asn Ala
20 25 30
Pro Thr Thr Ala Trp Leu Phe Ile Ala Ser Ala Pro Phe Glu Val
35 40 45
Ala Glu Gly Glu Asn Val His Leu Ser Val Val Tyr Leu Pro Glu
50 55 60
Asn Leu Tyr Ser Tyr Gly Trp Tyr Lys Gly Lys Thr Val Glu Pro
65 70 75
Asn Gln Leu Ile Ala Ala Tyr Val Ile Asp Thr His Val Arg Thr
80 85 90
Pro Gly Pro Ala Tyr Ser Gly Arg Glu Thr Ile Ser Pro Ser Gly

```

	95		100		105
Asp	Leu	His	Phe	Gln	Asn
	110	Val	Thr	Leu	Glu
Asn	Leu	Gln	Val	Thr	Tyr
	125	Arg	Asn	Ser	Gln
His	His	Leu	Arg	Val	Ala
	140	Glu	Ser	Val	Gln
Ala	Ser	Ser	Thr	Thr	Val
	155	Thr	Glu	Lys	Gly
Cys	His	Thr	Asn	Asn	Thr
	170	Gly	Thr	Ser	Phe
Asn	Gln	Arg	Leu	Gln	Val
	185	Thr	Lys	Arg	Met
Asn	His	Val	Leu	Thr	Ile
	200	Asp	Pro	Ile	Arg
Glu	Tyr	Gln	Cys	Glu	Val
	215	Ser	Asn	Pro	Val
Asp	Pro	Leu	Lys	Leu	Thr
	230	Val	Lys	Tyr	Asp
Leu	Ile	Gly	Val	Leu	Val
	245	Gly	Ser	Leu	Leu
Cys	Phe	Leu	Leu	Leu	Arg
	260	Lys	Thr	Gly	Arg
Asp	Phe	Arg	Glu	Gln	Pro
	275	Pro	Pro	Ala	Ser
Pro	Ser	Asp	Ser	Ser	Ile
	290	Ser	Ser	Ser	Ile

<210> 32

<211> 724

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4596747CD1

<400> 32

Met	Phe	Asp	Thr	Thr	Pro	His	Ser	Gly	Arg	Ser	Thr	Pro	Ser	Ser
1				5					10					15
Ser	Pro	Ser	Leu	Arg	Lys	Arg	Leu	Gln	Leu	Leu	Pro	Pro	Ser	Arg
				20					25					30
Pro	Pro	Pro	Glu	Pro	Glu	Pro	Gly	Thr	Met	Val	Glu	Lys	Gly	Ser
				35					40					45
Asp	Ser	Ser	Ser	Glu	Lys	Gly	Gly	Val	Pro	Gly	Thr	Pro	Ser	Thr
				50					55					60
Gln	Ser	Leu	Gly	Ser	Arg	Asn	Phe	Ile	Arg	Asn	Ser	Lys	Lys	Met
				65					70					75
Gln	Ser	Trp	Tyr	Ser	Met	Leu	Ser	Pro	Thr	Tyr	Lys	Gln	Arg	Asn
				80					85					90
Glu	Asp	Phe	Arg	Lys	Leu	Phe	Ser	Lys	Leu	Pro	Glu	Ala	Glu	Arg
				95					100					105
Leu	Ile	Val	Asp	Tyr	Ser	Cys	Ala	Leu	Gln	Arg	Glu	Ile	Leu	Leu
				110					115					120
Gln	Gly	Arg	Leu	Tyr	Leu	Ser	Glu	Asn	Trp	Ile	Cys	Phe	Tyr	Ser
				125					130					135
Asn	Ile	Phe	Arg	Trp	Glu	Thr	Thr	Ile	Ser	Ile	Gln	Leu	Lys	Glu
				140					145					150
Val	Thr	Cys	Leu	Lys	Lys	Glu	Lys	Thr	Ala	Lys	Leu	Ile	Pro	Asn
				155					160					165
Ala	Ile	Gln	Ile	Cys	Thr	Glu	Ser	Glu	Lys	His	Phe	Phe	Thr	Ser
				170					175					180
Phe	Gly	Ala	Arg	Asp	Arg	Cys	Phe	Leu	Leu	Ile	Phe	Arg	Leu	Trp
				185					190					195

Gln	Asn	Ala	Leu	Leu	Glu	Lys	Thr	Leu	Ser	Pro	Arg	Glu	Leu	Trp
				200					205					210
His	Leu	Val	His	Gln	Cys	Tyr	Gly	Ser	Glu	Leu	Gly	Leu	Thr	Ser
				215					220					225
Glu	Asp	Glu	Asp	Tyr	Val	Ser	Pro	Leu	Gln	Leu	Asn	Gly	Leu	Gly
				230					235					240
Thr	Pro	Lys	Glu	Val	Gly	Asp	Val	Ile	Ala	Leu	Ser	Asp	Ile	Thr
				245					250					255
Ser	Ser	Gly	Ala	Ala	Asp	Arg	Ser	Gln	Glu	Pro	Ser	Pro	Val	Gly
				260					265					270
Ser	Arg	Arg	Gly	His	Val	Thr	Pro	Asn	Leu	Ser	Arg	Ala	Ser	Ser
				275					280					285
Asp	Ala	Asp	His	Gly	Ala	Glu	Glu	Asp	Lys	Glu	Glu	Gln	Val	Asp
				290					295					300
Ser	Gln	Pro	Asp	Ala	Ser	Ser	Ser	Gln	Thr	Val	Thr	Pro	Val	Ala
				305					310					315
Glu	Pro	Pro	Ser	Thr	Glu	Pro	Thr	Gln	Pro	Asp	Gly	Pro	Thr	Thr
				320					325					330
Leu	Gly	Pro	Leu	Asp	Leu	Leu	Pro	Ser	Glu	Glu	Leu	Leu	Thr	Asp
				335					340					345
Thr	Ser	Asn	Ser	Ser	Ser	Ser	Thr	Gly	Glu	Glu	Ala	Asp	Leu	Ala
				350					355					360
Ala	Leu	Leu	Pro	Asp	Leu	Ser	Gly	Arg	Leu	Leu	Ile	Asn	Ser	Val
				365					370					375
Phe	His	Val	Gly	Ala	Glu	Arg	Leu	Gln	Gln	Met	Leu	Phe	Ser	Asp
				380					385					390
Ser	Pro	Phe	Leu	Gln	Gly	Phe	Leu	Gln	Gln	Cys	Lys	Phe	Thr	Asp
				395					400					405
Val	Thr	Leu	Ser	Pro	Trp	Ser	Gly	Asp	Ser	Lys	Cys	His	Gln	Arg
				410					415					420
Arg	Val	Leu	Thr	Tyr	Thr	Ile	Pro	Ile	Ser	Asn	Pro	Leu	Gly	Pro
				425					430					435
Lys	Ser	Ala	Ser	Val	Val	Glu	Thr	Gln	Thr	Leu	Phe	Arg	Arg	Gly
				440					445					450
Pro	Gln	Ala	Gly	Gly	Cys	Val	Val	Asp	Ser	Glu	Val	Leu	Thr	Gln
				455					460					465
Gly	Ile	Pro	Tyr	Gln	Asp	Tyr	Phe	Tyr	Thr	Ala	His	Arg	Tyr	Cys
				470					475					480
Ile	Leu	Gly	Leu	Ala	Arg	Asn	Lys	Ala	Arg	Leu	Arg	Val	Ser	Ser
				485					490					495
Glu	Ile	Arg	Tyr	Arg	Lys	Gln	Pro	Trp	Ser	Leu	Val	Lys	Ser	Leu
				500					505					510
Ile	Glu	Lys	Asn	Ser	Trp	Ser	Gly	Ile	Glu	Asp	Tyr	Phe	His	His
				515					520					525
Leu	Glu	Arg	Glu	Leu	Ala	Lys	Ala	Glu	Lys	Leu	Ser	Leu	Glu	Glu
				530					535					540
Gly	Gly	Lys	Asp	Ala	Arg	Gly	Leu	Leu	Ser	Gly	Leu	Arg	Arg	Arg
				545					550					555
Lys	Arg	Pro	Leu	Ser	Trp	Arg	Ala	His	Gly	Asp	Gly	Pro	Gln	His
				560					565					570
Pro	Asp	Pro	Asp	Pro	Cys	Ala	Arg	Ala	Gly	Ile	His	Thr	Ser	Gly
				575					580					585
Ser	Leu	Ser	Ser	Arg	Phe	Ser	Glu	Pro	Ser	Val	Asp	Gln	Gly	Pro
				590					595					600
Gly	Ala	Gly	Ile	Pro	Ser	Ala	Leu	Val	Leu	Ile	Ser	Ile	Val	Ile
				605					610					615
Cys	Val	Ser	Leu	Ile	Ile	Leu	Ile	Ala	Leu	Asn	Val	Leu	Leu	Phe
				620					625					630
Tyr	Arg	Leu	Trp	Ser	Leu	Glu	Arg	Thr	Ala	His	Thr	Phe	Glu	Ser
				635					640					645
Trp	His	Ser	Leu	Ala	Leu	Ala	Lys	Gly	Lys	Phe	Pro	Gln	Thr	Ala
				650					655					660
Thr	Glu	Trp	Ala	Glu	Ile	Leu	Ala	Leu	Gln	Lys	Gln	Phe	His	Ser

Val	Glu	Val	His	Lys	Trp	Arg	Gln	Ile	Leu	Arg	Ala	Ser	Val	Glu	665	670	675
Leu	Leu	Asp	Glu	Met	Lys	Phe	Ser	Leu	Glu	Lys	Leu	His	Gln	Gly	680	685	690
Ile	Thr	Val	Ser	Asp	Pro	Pro	Phe	Asp	Thr	Gln	Pro	Arg	Pro	Asp	695	700	705
Asp	Ser	Phe	Ser												710	715	720

<210> 33
 <211> 331
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5052680CD1

<400> 33

Met	Arg	Pro	Ala	Leu	Ala	Val	Gly	Leu	Val	Phe	Ala	Gly	Cys	Cys	1	5	10	15
Ser	Asn	Val	Ile	Phe	Leu	Glu	Leu	Leu	Ala	Arg	Lys	His	Pro	Gly	20	25	30	35
Cys	Gly	Asn	Ile	Val	Thr	Phe	Ala	Gln	Phe	Leu	Phe	Ile	Ala	Val	40	45	50	55
Glu	Gly	Phe	Leu	Phe	Glu	Ala	Asp	Leu	Gly	Arg	Lys	Pro	Pro	Ala	60	65	70	75
Ile	Pro	Ile	Arg	Tyr	Ala	Ile	Met	Val	Thr	Met	Phe	Phe	Thr		80	85	90	95
Pro	Leu	His	Met	Ile	Phe	Arg	Ser	Gly	Ser	Leu	Ile	Ala	Asn	Met	100	105	110	115
Ile	Leu	Gly	Ile	Ile	Ile	Leu	Lys	Lys	Arg	Tyr	Ser	Ile	Phe	Lys	120	125	130	135
Tyr	Thr	Ser	Ile	Ala	Leu	Val	Ser	Val	Gly	Ile	Phe	Ile	Cys	Thr	140	145	150	155
Phe	Met	Ser	Ala	Lys	Gln	Val	Thr	Ser	Gln	Ser	Ser	Leu	Ser	Glu	160	165	170	175
Asn	Asp	Gly	Phe	Gln	Ala	Phe	Val	Trp	Trp	Leu	Leu	Gly	Ile	Gly	180	185	190	195
Ala	Leu	Thr	Phe	Ala	Leu	Leu	Met	Ser	Ala	Arg	Met	Gly	Ile	Phe	200	205	210	215
Gln	Glu	Thr	Leu	Tyr	Lys	Arg	Phe	Gly	Lys	His	Ser	Lys	Glu	Ala	220	225	230	235
Leu	Phe	Tyr	Asn	His	Ala	Leu	Pro	Leu	Pro	Gly	Phe	Val	Phe	Leu	240	245	250	255
Ala	Ser	Asp	Ile	Tyr	Asp	His	Ala	Val	Leu	Phe	Asn	Lys	Ser	Glu	260	265	270	275
Leu	Tyr	Glu	Ile	Pro	Val	Ile	Gly	Val	Thr	Leu	Pro	Ile	Met	Trp	280	285	290	295
Phe	Tyr	Leu	Leu	Met	Asn	Ile	Ile	Thr	Gln	Tyr	Val	Cys	Ile	Arg	300	305	310	315
Gly	Val	Phe	Ile	Leu	Thr	Thr	Glu	Cys	Ala	Ser	Leu	Thr	Val	Thr	320	325	330	


```
<210> 34
<211> 398
<212> PRT
<213> Homo sapiens
```

<400> 34															
Met	Leu	Gly	Arg	Ser	Gly	Tyr	Arg	Ala	Leu	Pro	Leu	Gly	Asp	Phe	
1				5					10					15	
Asp	Arg	Phe	Gln	Gln	Ser	Ser	Phe	Gly	Phe	Leu	Gly	Ser	Gln	Lys	
				20					25					30	
Gly	Cys	Leu	Ser	Pro	Glu	Arg	Gly	Gly	Val	Gly	Thr	Gly	Ala	Asp	
				35					40					45	
Val	Pro	Gln	Ser	Trp	Pro	Ser	Cys	Leu	Cys	His	Gly	Leu	Ile	Ser	
				50					55					60	
Phe	Leu	Gly	Phe	Leu	Leu	Leu	Leu	Val	Thr	Phe	Pro	Ile	Ser	Gly	
				65					70					75	
Trp	Phe	Ala	Leu	Lys	Ile	Val	Pro	Thr	Tyr	Glu	Arg	Met	Ile	Val	
				80					85					90	
Phe	Arg	Leu	Gly	Arg	Ile	Arg	Thr	Pro	Gln	Gly	Pro	Gly	Met	Val	
				95					100					105	
Leu	Leu	Leu	Pro	Phe	Ile	Asp	Ser	Phe	Gln	Arg	Val	Asp	Leu	Arg	
				110					115					120	
Thr	Arg	Ala	Phe	Asn	Val	Pro	Pro	Cys	Lys	Leu	Ala	Ser	Lys	Asp	
				125					130					135	
Gly	Ala	Val	Leu	Ser	Val	Gly	Ala	Asp	Val	Gln	Phe	Arg	Ile	Trp	
				140					145					150	
Asp	Pro	Val	Leu	Ser	Val	Met	Thr	Val	Lys	Asp	Leu	Asn	Thr	Ala	
				155					160					165	
Thr	Arg	Met	Thr	Ala	Gln	Asn	Ala	Met	Thr	Lys	Ala	Leu	Leu	Lys	
				170					175					180	
Arg	Pro	Leu	Arg	Glu	Ile	Gln	Met	Glu	Lys	Leu	Lys	Ile	Ser	Asp	
				185					190					195	
Gln	Leu	Leu	Leu	Glu	Ile	Asn	Asp	Val	Thr	Arg	Ala	Trp	Gly	Leu	
				200					205					210	
Glu	Val	Asp	Arg	Val	Glu	Leu	Ala	Val	Glu	Ala	Val	Leu	Gln	Pro	
				215					220					225	
Pro	Gln	Asp	Ser	Pro	Ala	Gly	Pro	Asn	Leu	Asp	Ser	Thr	Leu	Gln	
				230					235					240	
Gln	Leu	Ala	Leu	His	Phe	Leu	Gly	Gly	Ser	Met	Asn	Ser	Met	Ala	
				245					250					255	
Gly	Gly	Ala	Pro	Ser	Pro	Gly	Pro	Ala	Asp	Thr	Val	Glu	Met	Val	
				260					265					270	
Ser	Glu	Val	Glu	Pro	Pro	Ala	Pro	Gln	Val	Gly	Ala	Arg	Ser	Ser	
				275					280					285	
Pro	Lys	Gln	Pro	Leu	Ala	Glu	Gly	Leu	Leu	Thr	Ala	Leu	Gln	Pro	
				290					295					300	
Phe	Leu	Ser	Glu	Ala	Leu	Val	Ser	Gln	Val	Gly	Ala	Cys	Tyr	Gln	
				305					310					315	
Phe	Asn	Val	Val	Leu	Pro	Ser	Gly	Thr	Gln	Ser					

380
Glu Ala Val Leu Arg Ala Leu Lys
395

385

390

<210> 35
<211> 220
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5524468CD1

<400> 35
Met Thr Trp Leu Val Leu Leu Gly Thr Leu Leu Cys Met Leu Arg
1 5 10 15
Val Gly Leu Gly Thr Pro Asp Ser Glu Gly Phe Pro Pro Arg Ala
20 25 30
Leu His Asn Cys Pro Tyr Lys Cys Ile Cys Ala Ala Asp Leu Leu
35 40 45
Ser Cys Thr Gly Leu Gly Leu Gln Asp Val Pro Ala Glu Leu Pro
50 55 60
Ala Ala Thr Ala Asp Leu Asp Leu Ser His Asn Ala Leu Gln Arg
65 70 75
Leu Arg Pro Gly Trp Leu Ala Pro Leu Phe Gln Leu Arg Ala Leu
80 85 90
His Leu Asp His Asn Glu Leu Asp Ala Leu Gly Arg Gly Val Phe
95 100 105
Val Asn Ala Ser Gly Leu Arg Leu Leu Asp Leu Ser Ser Asn Thr
110 115 120
Leu Arg Ala Leu Gly Arg His Asp Leu Asp Gly Leu Gly Ala Leu
125 130 135
Glu Lys Leu Leu Leu Phe Asn Asn Arg Leu Val His Leu Asp Glu
140 145 150
His Ala Phe His Gly Leu Arg Ala Leu Ser His Leu Tyr Leu Gly
155 160 165
Cys Asn Glu Leu Ala Ser Phe Ser Phe Asp His Leu His Gly Leu
170 175 180
Ser Ala Thr His Leu Leu Thr Leu Asp Leu Ser Ser Asn Arg Leu
185 190 195
Gly His Ile Ser Val Pro Glu Leu Ala Ala Leu Pro Ala Phe Leu
200 205 210
Lys Asn Gly Leu Tyr Leu His Asp Asn Thr
215 220

<210> 36
<211> 706
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5944279CD1

<400> 36
Met Glu Glu Asn Pro Thr Leu Glu Ser Glu Ala Trp Gly Ser Ser
1 5 10 15
Arg Gly Trp Leu Ala Pro Arg Glu Ala Arg Gly Gly Pro Ser Leu
20 25 30
Ser Ser Val Leu Asn Glu Leu Pro Ser Ala Ala Thr Leu Arg Tyr
35 40 45
Arg Asp Pro Gly Val Leu Pro Trp Gly Ala Leu Glu Glu Glu Glu
50 55 60
Glu Asp Gly Gly Arg Ser Arg Lys Ala Phe Thr Glu Val Thr Gln
65 70 75

Thr	Glu	Leu	Gln	Asp	Pro	His	Pro	Ser	Arg	Glu	Leu	Pro	Trp	Pro
				80					85					90
Met	Gln	Ala	Arg	Arg	Ala	His	Arg	Gln	Arg	Asn	Ala	Ser	Arg	Asp
				95					100					105
Gln	Val	Val	Tyr	Gly	Ser	Gly	Thr	Lys	Thr	Asp	Arg	Trp	Ala	Arg
				110					115					120
Leu	Leu	Arg	Arg	Ser	Lys	Glu	Lys	Thr	Lys	Glu	Gly	Leu	Arg	Ser
				125					130					135
Leu	Gln	Pro	Trp	Ala	Trp	Thr	Leu	Lys	Arg	Ile	Gly	Gly	Gln	Phe
				140					145					150
Gly	Ala	Gly	Thr	Glu	Ser	Tyr	Phe	Ser	Leu	Leu	Arg	Phe	Leu	Leu
				155					160					165
Leu	Leu	Asn	Val	Leu	Ala	Ser	Val	Leu	Met	Ala	Cys	Met	Thr	Leu
				170					175					180
Leu	Pro	Thr	Trp	Leu	Gly	Gly	Ala	Pro	Pro	Gly	Pro	Pro	Gly	Pro
				185					190					195
Asp	Ile	Ser	Ser	Pro	Cys	Gly	Ser	Tyr	Asn	Pro	His	Ser	Gln	Gly
				200					205					210
Leu	Val	Thr	Phe	Ala	Thr	Gln	Leu	Phe	Asn	Leu	Leu	Ser	Gly	Glu
				215					220					225
Gly	Tyr	Leu	Glu	Trp	Ser	Pro	Leu	Phe	Tyr	Gly	Phe	Tyr	Pro	Pro
				230					235					240
Arg	Pro	Arg	Leu	Ala	Val	Thr	Tyr	Leu	Cys	Trp	Ala	Phe	Ala	Val
				245					250					255
Gly	Leu	Ile	Cys	Leu	Leu	Leu	Ile	Leu	His	Arg	Ser	Val	Ser	Gly
				260					265					270
Leu	Lys	Gln	Thr	Leu	Leu	Ala	Glu	Ser	Glu	Ala	Leu	Thr	Ser	Tyr
				275					280					285
Ser	His	Arg	Val	Phe	Ser	Ala	Trp	Asp	Phe	Gly	Leu	Cys	Gly	Asp
				290					295					300
Val	His	Val	Arg	Leu	Arg	Gln	Arg	Ile	Ile	Leu	Tyr	Glu	Leu	Lys
				305					310					315
Val	Glu	Leu	Glu	Glu	Thr	Val	Val	Arg	Arg	Gln	Ala	Ala	Val	Arg
				320					325					330
Thr	Leu	Gly	Gln	Gln	Ala	Arg	Val	Trp	Leu	Val	Arg	Val	Leu	Leu
				335					340					345
Asn	Leu	Leu	Val	Val	Ala	Leu	Leu	Gly	Ala	Ala	Phe	Tyr	Gly	Val
				350					355					360
Tyr	Trp	Ala	Thr	Gly	Cys	Thr	Val	Glu	Leu	Gln	Glu	Met	Pro	Leu
				365					370					375
Val	Gln	Glu	Leu	Pro	Leu	Leu	Lys	Leu	Gly	Val	Asn	Tyr	Leu	Pro
				380					385					390
Ser	Ile	Phe	Ile	Ala	Gly	Val	Asn	Phe	Val	Leu	Pro	Pro	Val	Phe
				395					400					405
Lys	Leu	Ile	Ala	Pro	Leu	Glu	Gly	Tyr	Thr	Arg	Ser	Arg	Gln	Ile
				410					415					420
Val	Phe	Ile	Leu	Leu	Arg	Thr	Val	Phe	Leu	Arg	Leu	Ala	Ser	Leu
				425					430					435
Val	Val	Leu	Leu	Phe	Ser	Leu	Trp	Asn	Gln	Ile	Thr	Cys	Gly	Gly
				440					445					450
Asp	Ser	Glu	Ala	Glu	Asp	Cys	Lys	Thr	Cys	Gly	Tyr	Asn	Tyr	Lys
				455					460					465
Gln	Leu	Pro	Cys	Trp	Glu	Thr	Val	Leu	Gly	Gln	Glu	Met	Tyr	Lys
				470					475					480
Leu	Leu	Leu	Phe	Asp	Leu	Leu	Thr	Val	Leu	Ala	Val	Ala	Leu	Leu
				485					490					495
Ile	Gln	Phe	Pro	Arg	Lys	Leu	Leu	Cys	Gly	Leu	Cys	Pro	Gly	Ala
				500					505					510
Leu	Gly	Arg	Leu	Ala	Gly	Thr	Gln	Glu	Phe	Gln	Val	Pro	Asp	Glu
				515					520					525
Val	Leu	Gly	Leu	Ile	Tyr	Ala	Gln	Thr	Val	Val	Trp	Val	Gly	Ser
				530					535					540
Phe	Phe	Cys	Pro	Leu	Leu	Pro	Leu	Leu	Asn	Thr	Val	Lys	Phe	Leu

Leu	Leu	Phe	Tyr	545	Leu	Lys	Lys	Leu	Thr	550	Leu	Phe	Ser	Thr	Cys	555	Ser
				560	Thr	Phe	Arg	Ala	Ser	565	Ala	Ala	Asn	Phe	Phe	570	Phe
Pro	Ala	Ala	Arg	575						580						585	
Pro	Leu	Val	Leu	590	Leu	Leu	Gly	Leu	Ala	595	Ile	Ser	Ser	Val	Pro	600	Leu
Leu	Tyr	Ser	Ile	605	Phe	Leu	Ile	Pro	Pro	610	Ser	Lys	Leu	Cys	Gly	615	Pro
Phe	Arg	Gly	Gln	620	Ser	Ser	Ile	Trp	Ala	625	Gln	Ile	Pro	Glu	Ser	630	Ile
Ser	Ser	Leu	Pro	635	Glu	Thr	Thr	Gln	Asn	640	Phe	Leu	Phe	Phe	Leu	645	Gly
Thr	Gln	Ala	Phe	650	Ala	Val	Pro	Leu	Leu	655	Ile	Ser	Ser	Ile	Leu	660	Leu
Met	Ala	Tyr	Thr	665	Val	Ala	Leu	Ala	Asn	670	Ser	Tyr	Gly	Arg	Leu	675	Ile
Ser	Glu	Leu	Lys	680	Arg	Gln	Arg	Gln	Thr	685	Glu	Ala	Gln	Asn	Lys	690	Val
Phe	Leu	Ala	Arg	695	Arg	Ala	Val	Ala	Leu	700	Thr	Ser	Thr	Lys	Pro	705	Ala

Leu

<210> 37

<211> 466

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 6114480CD1

<400> 37

Met	Ala	Phe	Val	Leu	Ile	Leu	Val	Leu	Ser	Phe	Tyr	Glu	Leu	Val			
1				5					10					15			
Ser	Gly	Gln	Trp	Gln	Val	Thr	Gly	Pro	Gly	Lys	Phe	Val	Gln	Ala			
				20					25					30			
Leu	Val	Gly	Glu	Asp	Ala	Val	Phe	Ser	Cys	Ser	Leu	Phe	Pro	Glu			
				35					40					45			
Thr	Ser	Ala	Glu	Ala	Met	Glu	Val	Arg	Phe	Phe	Arg	Asn	Gln	Phe			
				50					55					60			
His	Ala	Val	Val	His	Leu	Tyr	Arg	Asp	Gly	Glu	Asp	Trp	Glu	Ser			
				65					70					75			
Lys	Gln	Met	Pro	Gln	Tyr	Arg	Gly	Arg	Thr	Glu	Phe	Val	Lys	Asp			
				80					85					90			
Ser	Ile	Ala	Gly	Gly	Arg	Val	Ser	Leu	Arg	Leu	Lys	Asn	Ile	Thr			
				95					100					105			
Pro	Ser	Asp	Ile	Gly	Leu	Tyr	Gly	Cys	Trp	Phe	Ser	Ser	Gln	Ile			
				110					115					120			
Tyr	Asp	Glu	Glu	Ala	Thr	Trp	Glu	Leu	Arg	Val	Ala	Ala	Leu	Gly			
				125					130					135			
Ser	Leu	Pro	Leu	Ile	Ser	Ile	Val	Gly	Tyr	Val	Asp	Gly	Gly	Ile			
				140					145					150			
Gln	Leu	Leu	Cys	Leu	Ser	Ser	Gly	Trp	Phe	Pro	Gln	Pro	Thr	Ala			
				155					160					165			
Lys	Trp	Lys	Gly	Pro	Gln	Gly	Gln	Asp	Leu	Ser	Ser	Asp	Ser	Arg			
				170					175					180			
Ala	Asn	Ala	Asp	Gly	Tyr	Ser	Leu	Tyr	Asp	Val	Glu	Ile	Ser	Ile			
				185					190					195			
Ile	Val	Gln	Glu	Asn	Ala	Gly	Ser	Ile	Leu	Cys	Ser	Ile	His	Leu			
				200					205					210			
Ala	Glu	Gln	Ser	His	Glu	Val	Glu	Ser	Lys	Val	Leu	Ile	Gly	Glu			
				215					220					225			

Thr	Phe	Phe	Gln	Pro	Ser	Pro	Trp	Arg	Leu	Ala	Ser	Ile	Leu	Leu	
				230					235					240	
Gly	Leu	Leu	Cys	Gly	Ala	Leu	Cys	Gly	Val	Val	Met	Gly	Met	Ile	
				245					250					255	
Ile	Val	Phe	Phe	Lys	Ser	Lys	Gly	Lys	Ile	Gln	Ala	Glu	Leu	Asp	
				260					265					270	
Trp	Arg	Arg	Lys	His	Gly	Gln	Ala	Glu	Leu	Arg	Asp	Ala	Arg	Lys	
				275					280					285	
His	Ala	Val	Glu	Val	Thr	Leu	Asp	Pro	Glu	Thr	Ala	His	Pro	Lys	
				290					295					300	
Leu	Cys	Val	Ser	Asp	Leu	Lys	Thr	Val	Thr	His	Arg	Lys	Ala	Pro	
				305					310					315	
Gln	Glu	Val	Pro	His	Ser	Glu	Lys	Arg	Phe	Thr	Arg	Lys	Ser	Val	
				320					325					330	
Val	Ala	Ser	Gln	Gly	Phe	Gln	Ala	Gly	Arg	His	Tyr	Trp	Glu	Val	
				335					340					345	
Asp	Val	Gly	Gln	Asn	Val	Gly	Trp	Tyr	Val	Gly	Val	Cys	Arg	Asp	
				350					355					360	
Asp	Val	Asp	Arg	Gly	Lys	Asn	Asn	Val	Thr	Leu	Ser	Pro	Asn	Asn	
				365					370					375	
Gly	Tyr	Trp	Val	Leu	Arg	Leu	Thr	Thr	Glu	His	Leu	Tyr	Phe	Thr	
				380					385					390	
Phe	Asn	Pro	His	Phe	Ile	Ser	Leu	Pro	Pro	Ser	Thr	Pro	Pro	Thr	
				395					400					405	
Arg	Val	Gly	Val	Phe	Leu	Asp	Tyr	Glu	Gly	Gly	Thr	Ile	Ser	Phe	
				410					415					420	
Phe	Asn	Thr	Asn	Asp	Gln	Ser	Leu	Ile	Tyr	Thr	Leu	Leu	Thr	Cys	
				425					430					435	
Gln	Phe	Glu	Gly	Leu	Leu	Arg	Pro	Tyr	Ile	Gln	His	Ala	Met	Tyr	
				440					445					450	
Asp	Glu	Glu	Lys	Gly	Thr	Pro	Ile	Phe	Ile	Cys	Pro	Val	Ser	Trp	
				455					460					465	

Gly

<210> 38

<211> 2801

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 112301CB1

<400> 38

```

cgccctcccc gagcgagacc aaaacaggtg gaatccgggc tggagccgga gctccggcgg 60
cgcggtgtggc ggcacgtccc tccagacagt accacaggca cctggagtac cggcatcggt 120
cgctgtggccc cccgagtgtc cgtcagagcc taggggagcc tgccctcccg cgcctcgctcg 180
gggcccggccc aggcaccttg gccgccggcg cacggacgcg ggcacgagca ctagatcacg 240
gctgctggac ctcggcacgt tgacaagatt tctctggggt accgcgagg attactttga 300
atttcggtgg tcgcctgtgg tctggcatat ttagaactta agtctattat ttcgggcacc 360
atgactttga ggcttttaga agactgggtgc agggggatgg acatgaacct tcggaaagcg 420
ctattgattg ccggcatctc ccagagctgc agtgtggcag aaatcgagga ggctctgcag 480
gctggttttag ctcccttggg ggagtacaga ctgcttggaa ggatgttcag gagggatgag 540
aacaggaaag tagccttagt agggcttact gcggagacta gtcacgccct ggtccctaag 600
gagataccgg gaaaaggggg tatctggaga gtgatcttta agccccctga cccagataat 660
acatttttaa gcagattaaa tgaattttta gcgggagagg gcatgacagt ggggtgagttg 720
agcagagctc ttggacatga aaatggctcc ttagaccag agcaggcat gatcccgaa 780
atgtgggccc ctatgttggc acaggcatta gaggtctctc agcctgccct gcaatgcttg 840
aagtataaaa agctgagagt gttctcgggc agggagtctc cagaaccagg agaagaagaa 900
tttggacgct ggatgtttca tactactcag atgataaagg cgtggcagg gccagatgta 960
gagaagagaa ggcgattgct agagagcctt cgaggcccag cacttgatgt tattcgtgc 1020
ctcaagataa acaatccttt aattactgtc gatgaatgtc tgcaggctct tgaggaggta 1080
tttgggggta cagataatcc tagggagttg cagggtcaa atctaaccac ttaccagaag 1140

```

```

gatgaggaaa agttgtcggc ttatgtacta aggctggagc ctttgttaca gaagctggta 1200
cagagaggag caattgagag agatgctgtg aatcaggccc gcctagacca agtcattgct 1260
ggggcagtc acaaaaacaat tcgcagagag cttaatctgc cagaggatgg cccagcccct 1320
ggtttcttgc agttattggt actaataaag gattatgagg cagctgagga ggaggaggcc 1380
cttctccagg caatattgga aggtaatttc acctgagctc cagggaaacca cgaagggata 1440
tggcaatgag tagagcatga aggtagaaca gtctatatac tcttgtaga catacaatcc 1500
ctacctgtg ctgccaagta actcattttt gtgcaattct cagtataagc cctttgtcgt 1560
ttctgtgcct atttaaagtc tcctaaaggt gtaattgact aggaaggatg tagttctaca 1620
ctgccattta cctattttaa ttcctccttg tgaatatctt tgttgttgtt gttgagacag 1680
agtctcgctc tgcacccag gctggagtgc agtgacgtga tcttggtcct ctgcatcctc 1740
cgcttccag gtttaagcta ttctcctgcc tcagttgccc gagtagctgg gactacagga 1800
atgactcacc acaccagct aagttttgca ttatcagtag agacggggtt tctccatact 1860
ggtcaggctg gtctcgaact cctgacctca agtgatccac ctgcttggc ctcccaaagt 1920
gctgggatta caggcgtgag ccaccacacc cagcctcagt tcccttattt taataattgt 1980
cctatttcgt ggggttgcta taaagattac atgagacttt aatgggaaag tatttaggac 2040
agtgtatggg acaaagttag taccagacat gcataaggca tttgaaaacc tgggtgcttt 2100
ctctctcact ctaccattt atagacaggg gtcatataat ttttttgtt gagacgagt 2160
ctcgctctgt ggcccaggct ggagagcagt ggtgtgatct cagctcatgg gttcaagcga 2220
ttctcgtgcc tcagcctccc gagtagctgg gactataggc atgcgccacc aggcccagcc 2280
aatttttata ttttttagtag agatgggggt tctccatgtt agccaggcta cccaggctgg 2340
tctcgatctc ctgacctcag gtgatccacc cacctgggcc tcccaaagtg ctgggattac 2400
aggcgtgagc cactgcccc tgcttcttca cacttttctt ttgagactgg gtcgcacagt 2460
cgcacagtga gactccatct cctatgaaaa aaaagattgg cctagcgtct ggctcaggcc 2520
tgtaatccca gaactttggg aggccaaagg gggcggtatca caaggtcagg agatcgaggc 2580
cagcatggcc aacatggtga caccctgttt ctactaaaaa taaaaaaa attagccggg 2640
catggtggcg catgcctata gtgtcagcta ctctggagaa tcgcttgaa cggggaggcg 2700
gaggttgtag tgagccgaga ttgcgtcact gcactacagc ctgggcaaca gagcgagact 2760
cccgtctcag aaaaaaaaaa ttttggtgcc aaccggacga g 2801

```

<210> 39

<211> 2656

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 997947CB1

<220>

<221> unsure

<222> 2554, 2587, 2604, 2606, 2611, 2644, 2646-2647, 2651

<223> a, t, c, g, or other

<400> 39

```

ccactctcta gagctcccg actcctagct ctgagaatgc ctgcggaatg atcgccccc 60
aggccggctg ccgcgctgc cgctgctgct gtatttgcta ctgctgctgc cgccgctct 120
gcttccactc ggctctgact ggcaggcaga aagtgcact gacgaggga aggtctctgc 180
agtgagtgga gagcctacat aaaagagagt aaagaggggc aaaaaccag atcagaatgc 240
aggcgacgtc caaccttctc aacctcctgc tgctgtctt gtttgccgga ttagatcctt 300
ccaagactca gattagtcc aaagaagggt ggcagggtga cagctcagct caggatcctg 360
atggggcggtg catttgcaca gttgttgctc cagaacaaaa cctgtgttcc cgggatgcca 420
aaagcaggca acttcgcca ctactggaaa aggttcagaa catgtcccag tctattgaag 480
tcttaaaact gagaaactcag agagatttcc aatatgtttt aaaaatggaa acccaaata 540
aagggtgtaa ggcaaaattt cggcagattg aagatgatcg aaagacactt atgaccaagc 600
attttcaggga gttgaaagag aaaatggacg agctcctgcc tttgatcccc gtgctggaac 660
agtacaaaac agatgctaag ttaatcacc agttcaagga ggaaataagg aatctgtctg 720
ctgtcctcac tggatttcag gaggaattg gtgcctatga ctacgaggaa ctacaccaa 780
gagtgtgtag cttggaaca agacttcgtg actgcatgaa aaagctaaca tgtggcaaac 840
tgatgaaaa cacaggccca gttacagtca agacatctgg aaccgattt ggtgcttga 900
tgacagacc tttagcatct gagaaaaaca acagagtctg gtacatggac agttatacta 960
acaataaaat tgttcgtgaa tacaatacaa ttgcagactt tgtcagtggt gctgaatcaa 1020
ggacatacaa ccttctcttc aagtgggag gaactaacca tgttgtctac aatggctcac 1080
tctattttta caagtatcag agtaatatca tcatcaaata cagctttgat atggggagag 1140

```

```

tgcttgccca acgaagcctg gagtatgctg gttttcataa tgtttacccc tacacatggg 1200
gtggattctc tgacatcgac ctaatggctg atgaaatcgg gctgtgggct gtgtatgcaa 1260
ctaaccagaa tgcaggcaat attgtcatca gccaaacttaa ccaagatacc ttggagggtga 1320
tgaagagctg gagcactggc taccccaaga gaagtgcagg ggaatctttc atgatctgtg 1380
ggacactgta tgtcaccaac tcccacttaa ctggagccaa ggtgtattat tcctattcca 1440
ccaaaacctc cacatatgag tacacagaca ttcccttcca taaccaatac ttccacatat 1500
ccatgcttga ctacaatgca agagatcgag ctctctatgc ctggaacaat ggccaccagg 1560
tgctgttcaa tgtcacccctt ttccatatca tcaagacaga ggatgacaca taggcaaatg 1620
tgacatgttt tcattgattt aaacagtggtg atttgtgata aactctataa gaccccttcc 1680
gtttttttct tcactattat ttttcatcat ttctccaaag caaagcattt ttattgtaaa 1740
gttgggtgtt caaaaacata gctgagcttg tctaacttac catgttggaa acacatctta 1800
acttctaaat ttacaaggcc tatcatgtcc ttgtcatgaa aagcactaaa aaaaaaaga 1860
gtttaagtgg ctaaagtcat agttttgcaa gagattaatg atctgcctta tattagagtc 1920
agagactaat ggtggcttaa atgcacgaat gtcttttttt tttaaaactg tcatttttta 1980
ctgtcttttg tcccatatca ggaaatattt tggtaggaat taggagaaca aaaagcactt 2040
ttatcccat tttttcttta aaaaatgtaa ggatttcatt tatattgaaa aataatatta 2100
atcatttttg tgttaacaca attctctgat gcggtgctgt acagtcattt ttaaatctct 2160
tgctaaccatt ttattggcag tatgtatttc taccattgta accaccattg tgctattgta 2220
tctcttctact tctgtgaaag taatattttt tataaaatac actgaaattht aacctcagta 2280
attgagtcct ttttcaagtg tggtaagaaa taatcttctt ggcttacccc ttacataaag 2340
cattataaac taaaatgaaa accaaaccag acacctgaca tagagtcttt attttacccc 2400
aagttttttg ggccactgac attgaatgca acaactgatt tcatacaact gagttactct 2460
gttcaactcca ctgaatgcaa cccatatagt ttcttgacaa aggtgcactc ggattctaaa 2520
tattggattt gagttgactc tactcattta ttanacacta aagaaatttt gttcttcata 2580
gttaaancta ctgacttta attnanattt nacatacaac ttgcaataat gaaattcctt 2640
atgncnngac nctgaa 2656

```

<210> 40
 <211> 968
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1521513CB1

```

<400> 40
tctacatcta tcggagctga acttcctaaa agacaaaagt tttatctttc aagattcatt 60
ctccctgaat cttaaccaaa aaacactcct gaggagaaaag aaagagaggg agggagagaa 120
aaagagagag agagaaacaa aaaaccaaag agagagaaaa aatgaattca tctaaatcat 180
ctgaaacaca atgcacagag agaggatgct tctcttccca aatgttctta tggactgttg 240
ctgggatccc catcctattt ctcaagtgcct gtttcatcac cagatgtgtt gtgacatttc 300
gcatctttca aacctgtgat gagaaaaagt ttcagctacc tgagaatttc acagagctct 360
cctgctacaa ttatggatca ggttcagtca agaattgttg tccattgaac tgggaatatt 420
ttcaatccag ctgctacttc ttttctactg acaccatttc ctgggcgtta agtttaaaga 480
actgctcagc catgggggct cacctgggtg ttatcaactc acaggaggag cagggaattcc 540
tttccataaa gaaacctaaa atgagagagt tttttattgg actgtcagac cagggtgtcg 600
agggtcagtg gcaatgggtg gacggcacac ctttgacaaa gtctctgagc ttctgggatg 660
taggggagcc caacaacata gctacctggg aggactgtgc caccatgaga gactcttcaa 720
acccaaggca aaattggaat gatgtaacct gtttctcaa ttattttcgg atttgtgaaa 780
tggtaggaat aaatcctttg aacaaaggaa aatctcttta agaacagaag gcacaactca 840
aatgtgtaaa gaaggaagag caagaacatg gccacacca ccgccccaca cgagaaattt 900
gtgcgctgaa cttcaaagga cttcataagt atttgttact ctgatataaa taaaaataag 960
tagtttta 968

```

<210> 41
 <211> 1837
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1863994CB1

<400> 41

```
ccccagcccc gctcagtcct gagcgcccgcg agtcgtcgcg ccgcccgcgc aagcatgcag 60
taaaggctga aaatctgggt cacagctgag gaagacctca gacatggagt ccaggatgtg 120
gcctgcgctg ctgctgtccc acctctctcc tctctggcca ctgctgttgc tgcccctccc 180
accgctgtct cagggtctct catcctcccc tcgaacccca ccagcccccag cccgcccccc 240
gtgtgccagg ggaggccctt cggcccccacg tcatgtgtgc gtgtgggagc gagcacctcc 300
accaagccga tctcctcggg tcccaagatc acgtcggcaa gtcctgcctg gcaactgcac 360
ccagccacc ccacaggtt ttgaggaggg gccgcccctca tcccaatacc cctgggctat 420
cgtgtggggg cccaccgtgt ctcgagagga tggaggggag cccaactctg ccaatcccgg 480
atttctggac tatggttttg cagccccctca tgggctcgca accccacacc ccaactcaga 540
ctccatgcga ggtgatggag atgggcttat ccttgagag gcaactgccca ccctgcggcc 600
attcctgttc gggggccgtg ggggaaggtgt ggacccccag ctctatgtca caattaccat 660
ctccatcatc attgttctcg tggccactgg catcatcttc aagttctgtt gggaccgcag 720
ccagaagcga cgcagacctt cagggcagca aggtgccctg aggcaggagg agagccagca 780
gccaactgaca gacctgtccc cggctggagt cactgtgtgt ggggcttctg gggactcacc 840
tacccccacc cctgaccatg aggagccccg agggggagcc cggcctggga tgccccacc 900
caagggggct ccagccttcc agttgaaccg gtgagggcag gggcaatggg atgggagggc 960
aaagagggaa ggcaacttag gtcttcagag ctgggggtgg ggtgccctct ggatgggtag 1020
tgaggaggca ggcgtggcct cccacagccc ctggccctcc caagggggct ggaccagctc 1080
ctctctggga ggcaccttc ctctctccag tctctcagga tctgtgtctt attctctgtt 1140
gcccataact ccaactctgc cctcttttgt tcttctctat gccacctgtt ctaagacaac 1200
tctgccctct taaccttgat tccccctctt tgttctgaac tccccctctt attctggcct 1260
acccttgggt tctgactgtt gccctttccc tcttctctc aggattcccc tgggtaatat 1320
gtgatgcccc caatgttggg gtgcagccaa gcaggaggcc aaggggcccg cacagcccc 1380
atcccactga ggggtggggc gctgtgggga gctggggcca caggggctcc tggctcctgc 1440
cccttgacaca ccacctggaa cactccccag ccccacgggc aatcctatct gctcgcctc 1500
ctgcaggtgg gggcctcaca tatctgtgac ttccgggtccc tgtccccacc cttgtgcact 1560
cacatgaag ccttgacacac tcacctccac cttcacaggc catttgacac cgctcctgca 1620
ccctctcccc gtccataccg ctccgctcag ctgactctca tgttctctcg tctcacattt 1680
gcaactctct cttcccatc tctgtgtctc gctcactcag tggtcagcgt ttcctgcaca 1740
ctttacctct catgtgcgtt tcccggcctg atgttgtgtt ggtgtgcggc gtgctcactc 1800
tctccctcat gaacaccac ccacctcggt tcgcagc 1837
```

<210> 42

<211> 2124

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2071941CB1

<400> 42

```
gtgacgcggc tgccggaggtg acgcggggagg tcgcgcgccc cttccggcgc ggggagggcg 60
ctgaagatcg gggccgctcg gccgcaggcg cctccagcgc cgcgggatgt agcgcggggg 120
accgcggccc ccagcagagc ccgcctgccc ggctgtgcta ccatcagagg gagatctctg 180
ccccctgggg ctgagagacc ccaacctttc cccaagctga agctgcaggg tattgaggta 240
ccagccagat gtcttccac aaaggatctg tgggtggcaca ggggaatggg gctcctgcca 300
gtaacaggga agctgacacg gtggaactgg ctgaactggg acccctgcta gaagagaagg 360
gcaaacgggt aatcgccaac ccacccaaag ctgaagaaga gcaaaccatgc ccagtgcacc 420
aggaagaaga ggaggaggtg cgggtactga cacttcccc gcaagccac cacgccatgg 480
agaagatgga agagtgtgtg tacaaggtct gggagggagc ttggagggtc atcccatatg 540
atgtgtctcc tgactggcta aaggacaacg actatctgct acatggctat agacctccca 600
tgccctcctt tcgggcttgc ttcaagagca tcttccgcat tcatacagaa actggcaaca 660
tctggaccca tctgcttggg ttctgtctgt ttctctttt gggaaatctt accatgctca 720
gaccaaatat gtacttcacg gccctctac aggagaaggt ggtttttggg atgttctttt 780
tggtgtcagt gctctgcctc agcttctctt ggctcttca caccgtctat tgtcattcag 840
agaaagtctc tcggactttt tccaaactgg actattcagg gattgtctt ctaattatgg 900
ggagctttgt cccctggctc tattatctct tctactgctc cccacagcca cggtcatct 960
acctctccat cgtctgtgtc ctgggcattt ctgccaatcat tgtggcgagc tgggaccgtt 1020
ttgccactcc taagcacagg cagacaagag caggcgtgtt cctgggactt ggcttgagt 1080
gcgtcgtgac caccatgcac ttactatctg ctgagggtt tgtcaaggcc accacagtgg 1140
gccagatggg ctggttcttc ctcatggctg tgatgtacat cactggagct ggcctttatg 1200
```



```
ctgctcgaat tccctgagcgc ttctttcctg gaaaaattga catatgggtc cagtctcatc 1260
agattttcca tgccttggtg gtggcagcag cctttgtcca ctctatgga gtctccaacc 1320
ttcaggaatt ccgttacggc ctagaaggcg gctgtactga tgacaccctt ctctgagcct 1380
tcccacctgc ggggtggagg aggaacttcc caagtgtttt taaaaataac ttctttgctg 1440
aagttagagg aagagtctga gttgtctgtt tctagaagaa acctcttaga gaattcagta 1500
ccaaccaagc ttcagcccac ttccacaccc actgggcaat aaactttcca ttccattctt 1560
cctagctggg gatggggcat ggtcaaaact agccatcccc tcttcagcaa ggcatctacc 1620
ggccccctcac agagacagta ctttgaaact catgttgaga ttttaccctc tcctccaacc 1680
attttgggaa aattatggac tgggactctt cagaaattct gtcttttctt ctggaagaaa 1740
atgtccctcc cttaccccca tccttaactt tgtatcctgg cttataacag gccatccatt 1800
ttgttagcac acttttcaaa aacaattata taccctggtc ccacttttct agggcctgga 1860
tctgtttata gagcaggaag aataaagcca ccaactttta cctagcccggt ctaatcatgg 1920
aagtgtgtcc aggtctcaag taacttgagt ttttaatttt ttttttttct tggcagagta 1980
atgtaaaatt taaatgggga aagatattta atatttaata ctaagcttta aaaagaaacc 2040
tgctatcatt gctatgtatc ttgatgcaaa gactatgatg ttaataaaaag aaagtacaga 2100
agagacttgg cattcaaaaa aaaa 2124
```

<210> 43
<211> 993
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2172512CB1

```
<400> 43
ccgcgcgctt actttgttta taacttgaaa aatcctctcc gtctcccttc cctgcctcct 60
ttcccttccc ttctctctgc cagtacaact agaccggcg tctggcgctc ccggtgcccc 120
gcattctcgc gggcaggcgg attaattgga attcttcaaa atgtcagggtg tggtagccac 180
agccccctgaa cagcctgcag gtgaaatgga aaatcaaaca aaaccaccag atccaaggcc 240
tgatgtcctt cctgaataca gttctcattt tttaaccagga ccccttgga cagctgtccc 300
tccacctact gggtacccag gaggttggc tatgggatac tacagtccac agcaaccacg 360
taccttccct ttgtaccagc cagttgggtg tatccatcct gtccgggtatc agcctggcaa 420
atatectatg ccaaatcagt ctgttccaat aacatggatg ccagggccaa ctccatggc 480
aaactgccct cctggtctgg aatacttagt tcagttggac aacatacatg ttcttcagca 540
ttttgagcct ctggaaatga tgacatgttt tgaactaat aatagatatg atattaaaaa 600
caactcagac cagatggttt acattgtaac cgaagacaca gatgacttta ccaggaatgc 660
ctatcggaca ctaaggccct tcgtctccg ggtcactgat tgtatgggcc gagaaatcat 720
gacaatgcag agacccttca gatgcacctg ctgttgcttc tgttgcccct ctgccagaca 780
agagctggag gtgcagtgtc ctctgggtgt caccattggc tttgttgccg aacattggaa 840
cctgtgcagg gcggtgtaca gcatccaaaa aaaaaaaaaa aaaattgcgg cgcaagctta 900
ttccctttag tgaggttaat ttttgggggc atgtgccgtg gcagtgcggg ggtgttggtg 960
gttgtgtggg ggtgggggtg ggttgggtg ggt 993
```

<210> 44
<211> 2214
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2483172CB1

```
<400> 44
ggctctcggc ccagcgcgcc tgccttcgcc gcccgccgtc gtccttcgcc cgctgcacga 60
cgacgcgacg cccctgtctg aggcggcgga cccgaccgga cccagaccca gacgcaagat 120
ggcgacggcc gcgtgactgc ctcagcgtcc ccgagctcgg ctccgagtgc acctacggac 180
tgactgtggg ggcagagaag ggcgagatca ggactctgtc tttgttaatc gtgactgcat 240
gaaggtcgcc tccctcgggc ctacttggtg ggagtgtctg gtattgttct aaggccagga 300
gcacggtgag ccacagtctg ttggtagaat ttggcgtctt gatagttgag aaaatggcga 360
tgacactgtt ggaagactgg tgccggggga tggatgtgaa ctcccagaga gctctgttag 420
tctggggcat cccagtgaac tgtgatgagg ctgaaatcga agagaccctc caggctgcga 480
```

```

tgccccaggt ctcctaccga atgcttggga gaatgttctg gaggggaagaa aatgcgaaaag 540
cagccttatt agagctcact ggcgctgtag attacgccgc gatccccagg gagatgccgg 600
gcaaaggagg ggtctggaaa gtgttattta agcccccac ttctgatgct gaatttttag 660
aaagattgca cctcttccta gctagagagg ggtggaccgt gcaagatgtt gcccggtgtcc 720
ttgggtttca gaaccctact ccgaccccg gcccagagat gccagcagag atgctaaact 780
atattttgga taatgttatt cagcctcttg ttgagtccat atggtacaag aggctgacac 840
ttttctcggg gagggacatc ccagggcctg gagaggaaac ctttgatccc tggctggagc 900
acactaatga ggtccctagag gagtggcagg tgtccgatgt agaaaagagg cggcggttga 960
tgagagagtct tagaggcccc gccgctgatg ttattcgcct ccttaagtcc aacaacccc 1020
cgataaccac tgccgaatgc ctgaaggcgc ttgagcagg gtttgggagc gttgagagct 1080
ctagggtatg ccagatcaaa tttctgaaca cttatcagaa cccgggagaa aaattgtctg 1140
cttatgtcat tcgtctggag cctctgctac agaaggtggt agagaagggg gccattgata 1200
aagataatgt gaaccaggcc cgcctagagc aggtcattgc cggggccaac cacagcggg 1260
ccatccgaag gcagctgtgg cttaccgggg ctggggaagg gccagcccca aacctctttc 1320
agttgctggt gcagatccgt gaggaggaag ccaaggagga ggaggaggag gctgaggcca 1380
cccttctgca gttaggcctg gaagggcact tctgagtgc aggaaaggca gctttagtgc 1440
agacctagat cacagctact tttcttgtcc ctgtggggtc ttacagatgt gtctctgagt 1500
agtaaaaggct tagccttggt ctgttttgtt gttttttgga ggggaagggt agtcaggcct 1560
gagtattcat gtaacattct aaaattgtgc cagcgagcac cgtgaacgac tgcaatgcaa 1620
gcggtcttg ctggctaagg taaagggttg gttggacaca gcgcttagtg cacgctgtca 1680
tcattggacat cataatcagt tgtgaaaaac acgcgaacct atgacacttc ttattccaca 1740
ctgaatgtga aattgcatgt tcagatgttt actacgaggc ctggctcaca ggaagtgttc 1800
agtaaaaagta tgcactgtta gattactgat aacgcggata gatttttgtt taccataaat 1860
tgttccagat ttatattaat ggaaggaaat gtgcatttat tagctattac tcaactttac 1920
aatgcaaaca tcttatttct catctttaaa catgtcgacc agtttaattg aaaagtattc 1980
tgagactgca aaatggggtg ttaaaaaata ctgcagttac ggagctgtgt aaaccagttt 2040
ctcattgcat aagatacaga tgtaaattgc atggagaggt tgatatgcac ctgtacagta 2100
attcactccc ccatttcaca tctttgtcag agaattagttc ttgttcatac tgagtgttct 2160
aaatttgaag ttatatatac aaatttaaat atttttaaaa ttcaaaaaaa aaaa 2214

```

<210> 45

<211> 897

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2656128CB1

<400> 45

```

gcaaaatgca tgacagtaac aatgtggaga aagacattac accatctgaa ttgcctgcaa 60
acccagggtt tctgcattca aaagagcatt ctattaaagc taccttaatt tggcgcttat 120
ttttcttaat catgtttctg acaatcatag tgtgtggaat ggttgctgct ttaagcgcaa 180
taagagctaa ctgccatcaa gagccatcag tatgtcttca agctgcatgc ccagaaagct 240
ggattgggtt tcaaagaaaag tgtttctatt tttctgatga caccaagaac tggacatcaa 300
gtcagagggt ttgtgactca caagatgctg atcttgctca ggttgaaagc ttccaggaaac 360
tgaatttcct gttgagatat aaaggcccat ctgatcactg gattgggctg agcagagaaac 420
aaggccaacc atggaaatgg ataaatggta ctgaatggac aagacagtta gtcatgaaag 480
aagatgggtg caacttgtat gttgcaaagg tttcacaagt tcctcgaatg aatccaagac 540
ctgtcatggt ttccatcctt gggagcagga gagtgtgcct atttgaatga caaagggtgc 600
agtagtgcca ggcactacac agagaggaag tggatttgtt ccaaatcaga tatacatgtc 660
tagatgttac agcaaagccc caactaatct ttagaagcat attggaactg ataactccat 720
tttaaaatga gcaaagaatt tatttcttat accaacaggt atatgaaaat atgctcaata 780
tcactaataa ctgggaaaat acaaatcaaa atcatagtaa aatattacct gttttcatgg 840
tgctaataat acctgttctc ccactgctaa tgacataccc gagactgagt aatttaa 897

```

<210> 46

<211> 2167

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5855841CB1

<400> 46

```
gccccgggccc gagttcgctg caagtcggcg gaaagtttgg ctgcgcgggt tccccgaag 60
ttcagagtga agacatttcc acctggacac ctgaccatgt gcctgcccctg agcagcgagg 120
cccaccaggc atctctgttg tgggcagcag gggcagggtcc tggctctgtgg accctcggca 180
gttggcaggc tccctctgca gtgggggtctg ggctcgggcc ccaccatgtc gagcctcggc 240
ggtggctccc aggatgcccg cggcagtagc agcagcagca ccaatggcag cggtggcagt 300
ggcagcagtg gcccacaggc agggagcagca gacaagagtg cagtgggtggc tgccgcccga 360
ccagcctcag tggcagatga cacaccaccc cccgagcgtc ggaacaagag cggtatcatc 420
agtggagcccc tgaacaagag cctgcgccgc tccccccgc tctccacta ctcttctttt 480
ggcagcagtg gtggtagtgg cgggtggcagc atgatggcg gagagtctgc tgacaaggcc 540
actgcggctg cagccgctgc ctccctgttg gccaatggg atgacctggc ggcgcccatg 600
gcggtggaca aaagcaaccc tacctcaaac cacaacagtg gtgctgtggc cagcctgctg 660
agcaaggcag agcggggccac ggagctggca gccgagggac agctgacgct gcagcagttt 720
gcgcagtgcca cagagatgct gaagcgcgtg gtgcaggagc atctcccgt gatgagcgag 780
gcgggtgctg gcctgcctga catggaggct gtggcagtg cgaagccct caatggccag 840
tccgacttcc cctacctggg cgttttcccc atcaaccag gcctcttcat tatgacccc 900
gcaggtgtgt tcttgccga gagcgcgtg cacatggcg gcctggctga gtaccccatg 960
caggggagagc tggcctctgc catcagctcc ggcaagaaga agcggaaacg ctgcggcatg 1020
tgcgcgccct gccggcggg catcaactgc gagcagtgca gcagttgtag gaatcgaag 1080
actggccatc agatttgcaa attcagaaaa tgtgaggaa tcaaaaaaga gccttccgt 1140
gctctggaga aggtgatgct tccgacggga gccgccttcc ggtggtttca gtgacggcg 1200
cggaacccaa agctgcccctc tccgtgcaat gtcactgctc gtgtggtctc cagcaaggga 1260
ttcggcgcaa gacaaacgga tgcacccgtc tttagaacca aaaatattct ctacagatt 1320
tcattcctgt tttatataat atattttttg ttgtcgtttt aacatctcca cgtccctagc 1380
ataaaaaaga aaagaaaaaa atttaaactg ctttttcgga agaacaacaa caaaaaagag 1440
gtaaaagcga atctataaag taccgagact tccctgggcaa agaattggca atcagtttcc 1500
ttcctgtgtc gatgtcgatg ttgtctgtgc aggagatgca gtttttgtgt agagaatgta 1560
aattttctgt aaccttttga aatctagtta ctaataagca ctactgtaat ttagcacagt 1620
ttaactccac cctcatttaa acttcccttg attctttccg accatgaaat agtgcatagt 1680
ttgcctggag aatccactca cgttcataaa gagaatgttg atggcgccgt gtagaagccg 1740
ctctgtatcc atccacgcgt gcagagctgc cagcagggag ctacagaaag gggaggggagc 1800
accaggccag ctgagctgca cccacagctc cgagactggg atccccacc ccaacagtga 1860
ttttggaaaa aaaaatgaaa gttctgttcg ttatccatt gcgatctggg gagccccatc 1920
tcgatatttc caatcctggc tacttttctt agagaaaata agtccttttt ttctggcctt 1980
gctaattggca acagaagaaa gggcttcttt gcgtgggtccc ctgctgggtgg ggggtgggtcc 2040
ccagggggcc ccctgcggcc tgggcccccc tgcccacggc cagcttctct ctgatgaaca 2100
tgctgtttgt attgttttag gaaaccaggc tgttttgtga ataaaacgaa tgcattgttt 2160
tgtcacg
```

<210> 47

<211> 1235

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 603462CB1

<400> 47

```
gtggagcccg cgagagagtg gcagcggggg ctgatggaag tgcaagtggg gctggagagg 60
gcacctact gtatccagca tgcctcaagg ccacagctct gtgttccagg ccttgctggg 120
gaccttcttc acctggggga tgacagcagc tggggcagct ctctgttctg tattctctag 180
tggaagaggg cggatcttag atggaagtct tggctttgct gcaggggtca tgttggcagc 240
ttcctattgg tctcttcttg ccccagcagt tgagatggcc acgtcctctg ggggcttccg 300
tgctttggcc ttcttccctg tggctgttgg cttcaccctt ggagcggtt ttgtctactt 360
ggctgacctc ctgatgcctc acttgggtgc agcagaagac ccccagacgg ccctggcact 420
gaacttcggc tctacgttga tgaagaagaa gtctgacctt gagggtcccg cgctgctctt 480
ccctgagagt gaactttcca tccgataga caagagtgg aatgggtgagg catatcagag 540
aaagaaggcg gcagccactg gccttccaga gggctctgct gtccctgtgc cttctcgagg 600
gaatctggca cagcccggcg gcagcagctg gaggaggatc gcactgctca tcttggccat 660
cactatacac aacgtttccag agggctctcg tgttggagtt ggatttgggg ctatagaaaa 720
```

```
gacggcatct gctacctttg agagtgccag gaatttggcc attggaatcg ggatccagaa 780
tttccccgag ggccttggtg tcagccttcc cttgcgaggg gcaggcttct ccacctggag 840
agctttcttg tatgggcagc tgagcgccat ggtggagccc ctggccgggg tctttggtgc 900
ctttgccgtg gtgctggtg agcccatcct gccctacgct ctggcctttg ctgccggtgc 960
catggtctac gtggtcatgg acgacatcat ccccgaaagg cagatcagtg gtaatgggaa 1020
actggcatcc tgggcctcca tcctgggatt tgtagtgtg atgtcactgg acgttggcct 1080
gggctagggc tgagacgctt cggaccccg gaaaggccat acgaagaaac agcagtgggt 1140
ggcttctatg ggacaacaag cttctttctt cacattaaaa cttttttcct tcctctcttc 1200
ttcatctcat tctcctgatt gactctaatt ataata 1235
```

<210> 48

<211> 2257

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 747681CB1

<400> 48

```
ccgagccgag cggaccgaag ggcgcgccga gatgcaggtg agcaagagga tgctggcggg 60
gggctgtagg agcatgccc gcccctcctt ggcctgctgg cagcccatcc tcctgctggt 120
gctgggctca gtgctgtcag gctcggccac gggctgcccc ccccgctgcg agtgctccgc 180
ccaggaccgc gctgtgctgt gccaccgcaa gcgctttgtg gcagtcccc agggcatccc 240
caccgagacg cgcctgctgg acctaggcaa gaaccgcatc aaaacgctca accaggacga 300
gttcgccagc tccccgacc tggaggagct ggagctcaac gagaacatcg tgagcgccgt 360
ggagcccgcc gccttcaaca acctcttcaa cctccggacg ctgggtctcc gcagcaaccg 420
cctgaagctc atcccgtag gegtcttcac tggcctcagc aaactgacca agctggacat 480
cagcgagAAC aagattgtta tcctgctgga ctacatgttt caggacctgt acaacctcaa 540
gtcactggag gttggcgaca atgacctcgt ctacatctct caccgcgcct tcagcggcct 600
caacagcctg gagcagctga cgctggagaa atgcaacctg acctccatcc ccaccgaggc 660
gctgtcccac ctgcacggcc tcctcgtcct gaggctccgg cactcaaca tcaatgccat 720
ccgggactac tccttcaaga ggctctaccg actcaaggctc ttggagatct cccactggcc 780
ctacttggac acctagacac ccaactgcct ctacggcctc aaactgacgt ccctgtccat 840
cacacactgc aatctgaccg ctgtgcccta cctggccgtc cgccacctag tctatctccg 900
cttctctaac ctctctaca accccatcag caccattgag ggctccatgt tgcattgagct 960
gctccggctg caggagatcc agctggtggg cgggcagctg gccgtggtgg agccctatgc 1020
cttccgcgcc ctcaactacc tgcgcgtgct caatgtctct ggcaaccagc tgaccacact 1080
ggaggaatca gtcttccact cgggtgggcaa cctggagaca ctcatcctgg actccaacct 1140
gctggcctgc gactgtcggc tcctgtgggt gttccggcgc cgctggcggc tcaacttcaa 1200
ccggcagcag cccactgtcg ccacgcccga gtttgtccag ggcaaggagt tcaaggactt 1260
cctgatgtg ctactgccc actacttcac ctgcccgcgc gcccgcatcc gggaccgcaa 1320
ggcccagcag gtgtttgtgg acgagggcca cacggtgcag tttgtgtgcc gggccgatgg 1380
cgacccgccc cccgccatcc tctggtcttc accccgaaag cacctggtct cagccaagag 1440
caatgggccc ctacagctct tcctgatgg cacgctggag gtgcgctacg cccaggtaca 1500
ggacaacggc acgtacctgt gcatcgcggc caacgcgggc ggcaacgact ccatgcccgc 1560
ccacctgcat gtgcgcagct actcgcccga ctggcccat cagcccaaca agaccttcgc 1620
tttcatctcc aaccagccgg gcgagggaga ggccaacagc acccgcgcca ctgtgccttt 1680
ccccttcgac atcaagacc tcactcatgc caccacctg ggcttcatct ctttctctgg 1740
cgctgtcttc ttctgctggt tgcgtgtgtt tctctggagc cggggcaagg gcaacacaaa 1800
gcacaacatc gagatcgagt atgtgcccc aaagtccgg acaggcatca gctccgcccga 1860
cgcgccccgc aagttcaaca tgaagatgat atgaggccgg ggcggggggc agggaccccc 1920
ggcgcgccgg gcaggggaag gggcctggcc gccacctgct cactctccag tccttcccac 1980
ctctcccta ccttctaca cacgttctct ttctccctcc cgctccgctc cctgtctgcc 2040
ccccgccagg cctcaccacc tgccctcctt ctaccaggac ctcaagaagg cagacctggg 2100
gacccacact acacaggggc attgacagac tggagttgaa agccgacgaa ccgacacgg 2160
gcagagtcaa taattcaata aaaaagttac gaactttctc tgtaacttgg gtttcaataa 2220
ttatggattt ttatgaaaac ttgaaataaa aaaaaaa 2257
```

<210> 49

<211> 2359

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 919469CB1

<400> 49

```
gtgggatttg aagatccact ccacttctgc tcatggcggg ccagggcctg cccctgcacg 60
tgccacact gctgactggg ctgctggaat gcctgggctt tgctggcgct ctctttggct 120
ggccttcaact agtgtttgtc ttcaagaatg aagattactt taaggatctg tgtggaccag 180
atgtggggcc gattggcaat gccacagggc aggttgactg caaagcccag gatgagaggt 240
tctcactcat cttcacocctg ggttccttca tgaacaactt catgacattc cccactggct 300
acatctttga ceggttcaag accacogtgg cacgcctcat agccatattt ttctacacca 360
ccgccacact catcatagcc ttacacctg caggctcagc cgtgctgctc ttccctggcca 420
tgccaatgct caccattggg ggaatcctgt ttctcatcac caacctgcag attgggaacc 480
tatttggcca acacogtctg accatcatca ctctgtacaa tggagcattt gactcttcc 540
cggcagctct ccttattatt aagcttcttt atgaaaaagg catcagcctc agggcctcct 600
tcacttctcat ctctgtctgc agtacctggc atgtagcagc cactttctc ctgatgcccc 660
gggggcacat cccataccca ctgcccccca actacagcta tggcctgtgc cctgggaatg 720
gcaccacaaa ggaagagaag gaaacagctg agcatgaaaa cagggagcta cagtcaagg 780
agttcctttc agcgaaggaa gagaccccag gggcagggca gaagcaggaa ctccgctcct 840
tctggagcta cgctttctct cggcgctttg cctggcacct ggtgtggctg tctgtgatac 900
agtgtggca ctacctcttc attggcactc tcaactcctt gctgaccaac atggccggtg 960
gggacatggc acgagtcagc acctacacaa atgcctttgc cttcactcag ttcggagtgc 1020
tgtgtgcccc ctggaatggc ctgctcatgg accggcttaa acagaagtac cagaaggaa 1080
caagaaagac aggttccctc actttggcgg tggccctctg ctgcagcgtg ccttcgctgg 1140
ccctgacatc cctgctgtgc ctgggcttcg cctctgtgc ctgagtcctc atcctccctc 1200
tccagtcact cacttctcat ctgcaagtga tcagcgcctc ctctctctat gggagcaacg 1260
cggccttctc cacccttgct ttcccttcag agcactttgg caagctcttt gggctggtga 1320
tgcccttgct ggctgtgggt tctctgctcc agttcccat cttcacctc atcaaaggct 1380
cccttcagaa tgacccattt tacgtgaatg tgatgttcat gcttgccatt cttctgacat 1440
tcttccaccc ctttctggtg tatcggaat gccgtacttg gaaagaaagt cctctgcaa 1500
ttgcatagtt cagaagccct cacttttcag ccccgaggat ggttttgttc atcttccacc 1560
acctttgagg acctcgtgtc ccaaaagact ttgcctatcc cagcaaaaaca cacacacaca 1620
cacacacaca caaaaataa agacacacaa ggacgtctgc gcagcaagaa aagaatctca 1680
gttgccaagc agattgatat cacacagact caaagcaag gcagtgtgaa cttctttatt 1740
tcaaaacaga agtgtctcct tgcaacttag cttggcagac ccttgactcc aggggagatg 1800
acctggggga ggaagtgtgt caactatttc tttaggcctg tttggctcgg aagcctatat 1860
gtgcctggat cctctgccac ggggttaaatt tttaggtgaa gagttaggtt gtcagggcct 1920
cagctatgct tccctgctct cctcaagag tgcagccttg gctagagaac tcacagctct 1980
gggaaaaaga ggagcagaca ggggttccctg ggcccagctc cagcccagcc actgatgctg 2040
gatgaccttg gcctgacctt ggtctggtct cagaatcact ttcccatct gtaaaattga 2100
gatgaatttt ggtgttgaaa gttcttctct gagcagatgt cctagaaggt tttaggaata 2160
gtgacagagt cagggcaccc caagggccat gggagccagc tgacctgctt gaccgaagga 2220
ttctgacag actatctttg gggatgtttt caagaaggga tataagttat ttactttggg 2280
catttaaaag aaaatttctc tcgggaataa ttttatagaa aaataaagct tctgtgtcta 2340
aggcaaaaaa aaaaaaaaaa                                     2359
```

<210> 50

<211> 2052

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 977658CB1

<400> 50

```
gggtttgaag ccgcgccgag agggagcgag gtcgcagtga cagcggcggg cgatcggacc 60
caggtgccc cgccgtaccc gcctgcgtcc cgcgtcccg cccagcatg acagccccgg 120
cgggtccgag cggtcagag accgagcgcc ttctgacccc caacccccgg tatgggaccc 180
aggcggggcc ttcacgggcc cctccgacac cccagaaga ggaagacct cgccgtcgtc 240
tcaaatactt ttctatgagt ccctgcgaca agtttcgagc caagggccgc aagccctgca 300
agctgatgct gcaagtgggtc aagatcctgg tggtcacggt gcagctcctc ctgtttgggc 360
tcagtaatac gctggctgtg acattccggg aagagaacac catgccttc cgacacctct 420
```

```

tctctgctggg ctactcggac ggagcgggatg acaccttcgc agcctacacg cgggagcagc 480
tgtaccaggc catcttccat gctgtggacc agtacctggc gttgctgac gtgtcactgg 540
gccggtatgc gtatgtccgt ggtgggggtg acccttggac caatggctca gggcttgctc 600
tctgccagcg gtactaccac cgaggccacg tggaccggc caacgacaca tttgacattg 660
atccgatggg ggttactgac tgcattccag tggatccccc cgagcggccc cctccgcccc 720
ccagcgacga tctcaccctc ttggaaagca gctccagtta caagaacctc acgctcaaat 780
tccacaagct ggtcaatgtc accatccact tccggctgaa gaccattaac ctccagagcc 840
tcatcaataa tgagatcccg gactgctata ccttcagcgt cctgatcacg tttgacaaca 900
aagcacacag tgggcggtac cccatcagcc tggagaccca gggccacatc caggagtgtg 960
agcaccocag tgtcttccag cacggagaca acagcttccg gctcctgttt gacgtggtgg 1020
tcatcctcac ctgctccctg tcttccctcc tctgcgcccg ctactcctt cgaggcttcc 1080
tgctgcagaa tgagtttgtg gggttcatgt ggcggcagcg gggacgggtc atcagcctgt 1140
gggagcgggt ggaatttgtc aatggctggg acatcctgct cgtcaccagc gatgtgctca 1200
ccatctcggg caccatcatg aagatcggca tccaggccaa gaacttggcg agctacgacg 1260
tctgcagcat cctcctgggc acctcgacgc tgctggtgtg ggtgggctgt atccgctacc 1320
tgaccttctt ccacaactac aatatcctca tccccacact gcgggtggcc ctgccacgag 1380
tcatgcgctt ctgctgtgct gtggctgtca tctacctggg ctactgcttc tgtggctgga 1440
tcgtgctggg gcccctatcat gtgaagtctc gctcactctc catggtgtct gactgctgtg 1500
tctcgctcat caatggggac gacatgtttg tgacgttcgc cgccatgcag gcgcagcagg 1560
gccgcagcag cctgggtgtg ctcttctccc agctctacct ttactcctt atcagcctct 1620
tcatctacat ggtgctcagc ctctcatcgc cgctcatcac cggcgccctac gacaccatca 1680
agcatcccg ggcgcagggc gcagaggaga gcgagctgca ggcctacatc gcacagtggc 1740
aggacagccc cactccgggc aagttccgcc gcgggagcgg ctccggctgc agccttctct 1800
gctgctgctg aagggacccc tccgaggagc attcgtgct ggtgaattga ttccagctga 1860
ctgcccgttg accgtaggcc ctggactgca gagaccccc ccccgaccc cgcttattta 1920
ttttaggggt ttgcttttaa ggatcggctc cctgtcgcgc ccgaggaggg cctggacctt 1980
tcgtgtcgga cccttggggg cggggagact ggggtggggag ggtgtgaat aaaagggaaa 2040
ataaaaaaaa aa 2052

```

<210> 51

<211> 1939

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1004703CB1

<400> 51

```

cttgatggcg tgggctgga gagccgcagt cccggctgca gcacctggga gaaggcagac 60
cgtgtgaggg ggcctgtggc ccagcgtgct gtggcctcgg gtagtgggaa gtggaggcag 120
gagccttccct tacacttcgc catgagtctc ctcatcgact ccagcatcat gattacctcc 180
cagatactat tttttggatt tgggtggctt ttcttcattg gccaatgtgt taaagactat 240
gagatacgtc agtatgttgt acaggtgatc ttctccgtga cgtttgcatt ttcttgacc 300
atgtttgagc tcatcatctt tgaatctta ggagtattga atagcagctc ccgttatttt 360
cactggaaaa tgaacctgtg tgtaattctg ctgactctgg tttcatggt gcctttttac 420
attggctatt ttattgtgag caatatccga ctactgcata aacaacgact gcttttttcc 480
tgtctcttat ggctgacctt tatgtatttc ttctggaaac taggagatct ctttcccat 540
ctcagcccaa aacatgggat cttatccata gaacagctca tcagccgggt tgggtgtgatt 600
ggagtgaact tcatggctct tctttctgga tttggtgctg tcaactgcc atacacttac 660
atgtcttact tctcaggaa tgtgactgac acggatattc tagccctgga acggcgactg 720
ctgcaaacca tggatatgat cataagcaaa aagaaaagga tggcaatggc acgggagaaca 780
atgttccaga agggggaagt gcataacaaa ccatcagggt tctggggaat gataaaaagt 840
gttaccactt cagcatcagg aagtgaatat cttactctta ttcaacagga agtggatgct 900
ttggaagaat taagcaggca gctttttctg gaaacagctg atctatatgc tacciaaggag 960
agaatagaat actccaaaac cttcaagggg aaatatttta attttcttgg ttacttttcc 1020
tctatttact gtgtttggaa aattttcatg gctaccatca atattgtttt tgatcgagtt 1080
gggaaaacgg atcctgtcac aagaggcatt gagatcactg tgaattatct gggaatccaa 1140
tttgatgtga agttttggtc ccaacacatt tccttcattc ttgttggaat aatcatcgtc 1200
acatccatca gaggattgct gatcactctt accaagttct tttatgccat ctctagcagt 1260
aagtccctca atgtcattgt cctgctatta gcacagataa tgggcatgta cttgtctcc 1320
tctgtgctgc tgatccgaat gagtatgcct ttagaatacc gcaccataat cactgaagtc 1380
cttgaggaaac tgcagttcaa cttctatcac cgttggtttg atgtgatctt cctggtcagc 1440

```

```
gctctctcta gcatactctt cctctatttg gctcacaaac aggcaccaga gaagcaaagt 1500
gcaccttgaa cttaagccta ctacagactg ttagaggcca gtggtttcaa aatttagata 1560
taagaggggg gaaaaatgga accagggcct gacattttat aaacaaacaa aatgctatgg 1620
tagcattttt caccttcata gcatactcct tccccgtcag gtgatactat gaccatgagt 1680
agcatcagcc agaacatgag agggagaact aactcaagac aatactcagc agagagcatc 1740
ccgtgtggat atgaggctgg tgtagaggcg gagaggagcc aagaaactaa aggtgaaaaa 1800
tacactggaa ctctggggca agacatgtct atggtagctg agccaaacac gtaggatttc 1860
cgttttaagg ttcacatgga aaagggtata gctttgcctt gagattgact cattaaaatc 1920
agagactgta aaaaaaaaaa 1939
```

<210> 52
<211> 1138
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1334051CB1

```
<400> 52
caaactgcaa cttatatctg caatttattt tggatatagac aagaggtagt ccagtagcac 60
actgggtggc tcagaagaaa ttctcaacac ctactctgcc agagagtcta tgtatgggat 120
tgaacaatct gtaaaactaaa ggatcctaata catgaaaata agtatgataa attataagtc 180
actattggca ctggtgttta tattagcctc ctggatcatt ttacagttt tccagaactc 240
cacaaagggt tggctctctc taaacttata catctccctc cattactgga acaactccac 300
aaagtcctta ttccctaaaa caccactgat atcattaaag ccactaacag agactgaact 360
cagaataaag gaaatcatag agaaactaga tcagcagatc ccaccagac ctttcaccca 420
cgtgaacacc accaccagcg ccacacatag cacagccacc atcctcaacc ctcgagatac 480
gtactgcagg ggagaccagc tgcacatcct gctggagggt aggggaccact tgggacgcag 540
gaagcaatat ggcggggatt tcctgagggc caggatgtct tccccagcgc tgatggcagg 600
tgcttcagga aaggtagctg acttcaacaa cggcacctac ctggtcagct tcaactctgt 660
ctgggagggc caggtctctc tgtctctgct gctcatccac cccagtgaag ggggtgcagc 720
tctctggagt gcaaggaacc aaggctatga cagggtgatc ttactggcc agtttgtcaa 780
tggcacttcc caagtccact ctgaatgtgg cctgatccta aacacaaatg ctgaattgtg 840
ccagtacctg gacaacagag accaagaagg cttctactgt gtgaggcctc aacacatgcc 900
ctgtgctgca ctcaactaca tgtattctaa gaacaagaaa gtttcttata ttagcaaaaca 960
agaaaagagc ctctttgaaa ggtaaaaata attactctt gagactacct gtgcaaatat 1020
tgtgatttgg cctatatact gatccaaaga aaggtcttgt gagtgattta attttgggtg 1080
tctttagtaa gagcctttgg ggaaaggatc tgtgaattca tttagagaca gtgcccat 1138
```

<210> 53
<211> 2117
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1336728CB1

```
<400> 53
tggccacagc aacaccact gagcacgctg ggagctgagt atggcgctcc tggctctgct 60
ggagctgggg ctgcttctgg ctgtgctggt ggtgacggcg acggcgctcc cgcctgctgg 120
tctgctgagc ctgctcacct ctggccaggg cgtctggat caagaggctc tgggcgccct 180
gttaaatagc ctggcggacc gtgtgactg caccacggg cgtgtggaa agtgccctgtc 240
tgtggaggac gccctgggcc tgggcgagcc tgagggtgca gggctgcccc cgggcccggg 300
cctggaggcc aggtacgtcg ccgcctcag tgcgcgcgc gtccgtgacc tcagcaaccc 360
cgaggggcacc tgtgaggaca ctgggctgg cctctgggcc tctcatgcag accacctcct 420
ggcctgtctc gagagcccca aggcctgac ccgggctg agctggctgc tgcagaggat 480
gcaggcccg gctgcccggc agacccecaa gacggcctgc gtatagatcc ctgagctgct 540
ggaggaggcg tggggggcgg gggctccggc cagtgtctgg ggcgtcctgg ctgcccctgt 600
ggaccatgtc agtagcgggt cttgcttcca cgccttgcc agccctcagt acttcgtgga 660
ctttgtgttc cagcagcaca gcagcagggt ccctatgacg ctggccgagc tgtcagcctt 720
gatgcagcgc ctgggggtgg gcaggaggcg ccacagtgc cacagtcatc ggcacagggg 780
```

```
agccagcagc cgggaccctg tgccctcat cagctccagc aacagctcca gtgtgtgga 840
cacggtatgc ctgagtgcca gggacgtgat ggctgcatat ggactgtcgg aacaggctgg 900
ggtgaccccg gaggcctggg cccaactgag ccctgccctg ctccaacagc agctgagtgg 960
agcctgcacc tcccagtcga ggcccccggt ccaggaccag ctccagccagt cagagaggta 1020
tctgtacggc tccctggcca cgctgctcat ctgcctctgc gcggtctttg gcctcctgct 1080
gctgacctgc actggctgca ggggggtcac ccactacatc ctgcagacct tccctgagcct 1140
ggcagtgggt gcactcactg gggacgctgt cctgcatctg acgcccagg tgctggggct 1200
gcatacacac agcgaagagg gcctcagccc acagcccacc tggcgccctc tggctatgct 1260
ggccggggtc tacgccttct tccctgttga gaacctcttc aatctcctgc tgcccaggga 1320
cccggaggac ctggaggacg ggccttgagg ccacagcagc catagccacg gggggccacg 1380
ccacggtgtg tccctgcagc tggcaccag cgagctccgg cagcccaagc cccccacga 1440
gggctcccg cagacctggg tggcgaggga gagcccgagg ctgctgaacc ctgagcccg 1500
gagactgagc ccagagttag ggctactgac ctatatgatc actctgggag acgcccgtgca 1560
caacttcgcc gacgggctgg ccgtggcgcg cgccttcgag tccctcctga agaccgggct 1620
ggccacctcg ctggccgtgt tctgccacga gttgccacac gagctggggg acttcgccgc 1680
cttgctgcac gcggggctgt ccgtgcgcca agcactgctg ctgaacctgg cctccgcgct 1740
cacggccttc gctggtctct acgtggcact cgcggttggg gtcagcgagg agagcgaggc 1800
ctggatcctg gcagtgcca ccggcctgtt cctctacgta gcactctgag acatgctccc 1860
ggcgatgttg aaagtacggg acccgcgggc ctggtccttc ttctgctgac acaacgtggg 1920
cctgctgggc ggctggaccg tccctgctgt gctgtccctg tacgaggatg acatcacctt 1980
ctgataccct gccctagtcc cccacctttg acttaagatc ccacacctca caaacctaca 2040
gcccagaaac cagaagcccc tatagaggcc ccagtcccaa ctccagtaaa gacactcttg 2100
tccttgaaaa aaaaaaa 2117
```

<210> 54
<211> 1495
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1452856CB1

```
<400> 54
ctgaaatccc gcgaggatca accgagctcg ccgaaaggag ggaggaacgt atcccttctg 60
gaggctgtct cagggggcag agggaccgga ccggaagtga cgtgagcggg ttccgggtgt 120
ctggagccca gcgggcggtg tgagagtccg taaggagcag cttccaggat cctgagatcc 180
ggagcagccg gggctcgagg ggctcctcaa gagttactga tctatgaaat ggcagagaat 240
ggaaaaaatt gtgaccagag acgtgtagca atgaacaagg aacatcataa tggaaatttc 300
acagaccctt ctccagtga tgaagaagagg agggaggagc ggggaagaaag gcagaatatt 360
gtcctgtgga gacagccgct cattaccttg cagtattttt ctctggaaat ccttgtaatc 420
ttgaaggaat ggacctcaaa attatggcat cgtcaaagca ttgtggtgtc ttttttactg 480
ctgcttgctg tgcttatagc tacgtattat gttgaaggag tgcatacaac gtatgtgcaa 540
cgtatagaga aacagtttct tttgtatgcc tactggatag gcttaggaat tttgtcttct 600
gttgggcttg gaacagggct gcacaccttt ctgctttatc tgggtccaca tatagcctca 660
gttacattag ctgcttatga atgcaattca gtttaatttc ccgaaccacc ctatcctgat 720
cagattatatt gtccagatga agagggcact gaaggaaacca tttctttgtg gagtatcatc 780
tcaaaaagta ggattgaagc ctgcatgtgg ggtatcggtc cagcaatcgg agagctgcct 840
ccataatttca tggccagagc agctcgcttc tcagggtgctg aaccagatga tgaagagtat 900
caggaatttg aagagatgct ggaacatgca gagtctgcac aagactttgc ctcccgggcc 960
aaactggcag ttcaaaaact agtacagaaa gttggatttt ttggaatttt ggcctgtgct 1020
tcaattccaa atcctttatt tgatctggct ggaataacgt gtggacactt tctggtacct 1080
ttttggacct tctttggtgc aaccctaatt ggaaaagcaa taataaaaat gcataatccag 1140
aaaatttttg ttataataac attcagcaag cacatagtgg agcaaatggg ggctttcatt 1200
ggtgctgtcc ccggcatagg tccatctctg cagaagccat ttcaggagta cctggaggct 1260
caacggcaga agcttcacca caaaagcgaa atgggcacac cacagggaga aaactggttg 1320
tcctggatgt ttgaaaagtt ggtcgttgtc atggtgtgtt acttcatcct atctatcatt 1380
aactccatgg cacaaggtta tgccaaacga atccagcagc ggttgaactc agaggagaaa 1440
actaaataag tagagaaagt tttaaactgc agaaattgga gtggatgggt tctgc 1495
```

<210> 55
<211> 1747
<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1562471CB1

<400> 55

```
accagcagaa ggctgggagt ctgtagtttg ttcctgctgc caggctccac tgaggggaac 60
ggggacctgt ctgaagagaa gatgcccctg ctgacactct acctgctcct cttctggctc 120
tcaggctact ccattgccac tcaaatcacc ggtccaacaa cagtgaatgg cttggagcgg 180
ggctccttga cgtgcagtg tgtttacaga tcaggctggg agacctactt gaagtggagg 240
tgtcaggagg ctatttggcg tgactgcaag atccttggtt aaaccagtgg gtcagagcag 300
gaggtgaaga gggaccgggt gtccatcaag gacaatcaga aaaaccgcac gttcactgtg 360
accatggagg atctcatgaa aactgatgct gacacttact ggtgtggaat tgagaaaact 420
ggaaatgacc ttgggggtcac agttcaagtg accattgacc cagcaccagt caccacaaga 480
gaaactagca gctcccaaac tctgaccggc caccacttgg acaacaggca caagctcctg 540
aagctcagtg tcctcctgccc cctcatcttc accatatgtc tgctgctttt ggtggccgcc 600
tcactcttgg cttggaggat gatgaagtac cagcagaaag cagccgggat gtccccagag 660
cagggtactgc agccctgga gggcgacctc tgctatgcag acctgacctt gcagctggcc 720
ggaacctccc cgcgaaaggc taccacgaag ctttcctctg cccagggttg ccagggtgga 780
gtggaatatg tcaccatggc ttccttgccg aaggaggaca tttcctatgc atctctgacc 840
ttgggtgctg aggatcagga accgacctac tgcaacatgg gccacctcag tagccacctc 900
cccggcaggg gccctgagga gccacgggaa tacagcacca tcagcaggcc ttagcctgca 960
ctccaggctc cttcttggac cccaggctgt gagcacactc ctgcctcctc gaccgtctgc 1020
cccctgctcc cctcatcagg accaaccggg ggactgggtg ctctgcctga tcagccagca 1080
ttgcccttag ctctgggttg ggcttggggc caagtctcag ggggcttcta ggagttgggg 1140
ttttctaaac gtcccctcct ctctacata gttgaggagg gggctaggga tatgctctgg 1200
ggctttcatg ggaatgatga agatgataat gagaaaaatg ttatgccta ccacatgtta tgggctgaat 1320
taccattatc ataatacaat gaacctttat ttattgccta ccacatgtta tgggctgaat 1320
aatggccccc aaagatatct gtgtcctaata cctcagaact tgtgactgtt accttctgtg 1380
gcagaaaagg acagtgcaga tgtatgtaag ttaaggactt tgagatagag aggttattct 1440
tgctgattca ggtgggcccc aaatatcacc acaagggtcc tcataagaaa gaggccagaa 1500
ggtcaaagag gtagagacaa agtgatgatg gaagtggacg tgggtgtgac gtgagcaggg 1560
gccatgaatg ccgcagcctt cagatgccag aaaggggaaa gaatggattc ccctgcctgg 1620
agcctccaaa agaaaccagc cctgcccacg ccttgacttg agcccattga aactgatctt 1680
gagctcctgg cctccagaat tgccaggagaa taaatttgtg ttgtttttaa tgaaaaaaa 1740
aaaaaaa
```

<210> 56

<211> 1473

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1618158CB1

<400> 56

```
acagcgggac agcccggcag ccttgggctt ccagcgctgg cttcatcgcc acctccctgc 60
cttctccccc ttccgcatte ttagggtgac gatggtggca ccatcaagca aaagatcttc 120
accttcgacg ccatgtttct caccactac tcacacatgg agaactaccg caagcgagag 180
gacctggtgt accagtcac tgtgaggctg cccgagggtc ggtatcaga caatgggtccc 240
tatgagtgcc atgtgggcat ctacgaccgc gccaccaggg agaagggtgg cctggcatca 300
ggcaacatct tcctcaacgt catggctcct cccacctcca ttgaagtggg ggctgctgac 360
acaccagccc ccttcagccg ctaccaagcc cagaacttca cgctgggtctg catcgtgtct 420
ggaggaaaaa cagcaccat ggtttatttc aaacgagatg gggaaaccaat cgacgcagtg 480
cccctatcag agccaccagc tgcgagctcc ggccccctac aggacagcag gcccttccgc 540
agccttctgc accgtgacct ggatgacacc aagatgcaga agtcaactgt cctcctggac 600
gccgagaacc ggggtgggcg accctacacg gagcgcccct cccgtggcct gacccagat 660
cccaacatcc tcctccagcc aaccacagag aacataccag agacggctcg gagccgtgag 720
tttcccgcct ggggtccacg cgcgcagccc acctacttcc tgcgcccacg ccgcaccccc 780
agcagtgcag gcaactgtga agtacgtgcc ctgctcacct ggacctcaa cccacagatc 840
gacaacgagg cctcttccag ctgagagggt aagcaccag ctctgtcgat gcccatgcag 900
```

```

gcagaggtca cgctggttgc ccccaaagga cccaaaattg tgatgacgcc cagcagagcc 960
cgggtagggg acacagttag gattctgggc catgggtttc agaacgaagt cttcccggag 1020
cccatgttca cgtggacgcg ggttgggagc cgcctcctgg acggcagcgc tgagttcgac 1080
gggaaggagc tgggtgctgga gcgggttccc gccgagctca atggctccat gtatcgctgc 1140
accgcccaga acccactggg ctccaccgac acgcacaccc ggctcatcgt gtttgaaaac 1200
ccaaatatcc caagaggaac ggaggactct aatggttcca ttggcccccac tggtgcccgg 1260
ctcaccttgg tgctcgccct gacagtgatt ctggagctga cgtgaaggca cccgcccgg 1320
ccactccatc aggcactgac atctccacga ccggttttca tttcttttct aaactatttc 1380
cagtcttgtt cttagtctct ttccatctgt gtcttggctt cttcagtcgg ttttaattaa 1440
acaaacagaa caattttccc caaaaaaaaaa aaa 1473

```

<210> 57
 <211> 1591
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1656935CB1

```

<400> 57
attaaacgac tcaactatagg gaatttggcc ctcgaggcaa gaattcggca cgagggttcc 60
tataggagag ctttagaatg aagtcattta gaggagcagg cgaatcctaa ccgcatctct 120
ctcttttagt ggactgaacc caaacatgaa tgtcaacagc atggacatga ccggtggctt 180
gtcggtagaag gacccatccc agtcccagtc acgcctcccc cagtggacgc accccaactc 240
catggataaac ttgcccagtg ccgcttcccc cctggagcag aaccctagca agcatgggtg 300
tatccctgga ggtctaagca ttgggcctcc aggtaagtcc tccattgatg actcctatgg 360
ccggtacgat ttaatccaga acagtgagtc accagccagt cctcccgtag ctgttcccc 420
tagctgggtca cgtgccaaat ctgacagtga taaaatctca aatggctcta gcatcaactg 480
gccccagaa ttccatccgg gagttccatg gaaaggactg cagaatattg accctgagaa 540
tgaccctgac gtcactccct gcagtgtccc cactgggcct accatcaaca ccaccatcca 600
ggatgtcaac cgctacctcc tcaagagtgg agggctcctc ccgccatcat ctcaaatgc 660
cacgctgcct tcttcgagtg cctggccact cagtgcctcc ggctacagta gctctttcag 720
cagcattgca tccgcaccta gtgttgacag taaactgtca gacatcaaat cgacgtggtc 780
ctctggccct acctcccaca cgcaagcctc tctgtctcat gaactatgga aggtgcccag 840
aaacagtact gcacccacga ggccacctcc agggttaacc aatcccaagc cctcctccac 900
ctggggtgcc agccccctcg gctggaccag ctctactccc tcgggttctg cctggagcac 960
cgacacctca ggaagaacca gcagctggct cgttcttcga aacctcactc cccagattga 1020
tggtttctaa ctgcggacat tgtgtttgca acatgggcct cttatcacat tccacctgaa 1080
tctgactcaa ggcaatgctg tggtcgggta cagctccaag gaggagggct tgccaaaggc 1140
ccaggaagtc ctctgcacaa ttgtgcgtcc ttgggagaca cttagccatt ccttggggccc 1200
gagtttccgc ttggttggaa cgaaggaagt tggaattcgg gtttccttta agccgccaga 1260
gggcccaggc cgatttggcc agagcacaat ttccaaggc ctggcacaat tccacgacca 1320
gaggggctgt tccaagctca caggacgtgg cgggattcac aggccacgcg ggaggggcaa 1380
ggcttctcac caacttgcac atatgagaca ttgtgaatta acatttttaag agcgttgcca 1440
ctaattggga caatcatcgg acaaggaccc tgagttcaat tttgtaaagt gccctgttat 1500
aaatgtttgc cgtgtagaat aaaggtttct ttccactata tcccattgta tcgggatata 1560
tcacatcgta acagtgttat tttgggaaga a 1591

```

<210> 58
 <211> 1858
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1859305CB1

```

<400> 58
gcaggcgcgg cgaaacttcc ctctaccgac ccggcccgcg gcgcgcaccg ttggcgctgg 60
acgcttccctc cttggaagcg cctctccctc aggtctcaag aactgttttg agatgcagga 120
attcatctaa ttttactgac cgggcgaggt gtgagagccc tagcatctga aagtggctga 180
cttgcgagtt gttatggaga aaacttgtat agatgcactt cctcttacta tgaattcttc 240

```

```

agaaaagcaa gagactgtat gtatttttgg aactgggtgat tttggaagat cactggggatt 300
gaaaaatgctc cagtgtgggtt attctgttgtt ttttgggaagt cgaaaccccc agaagaccac 360
cctactgccc agtgggtgcag aagtcttgag ctattcagaa gcagccaaga agtctgacat 420
cataatcata gcaatccaca gagagcatta tgattttctc acagaattaa ctgaggttct 480
caatggaaaa atattggtag acatcagcaa caacctcaaa atcaatcaat atccagaatc 540
taatgcagag taccttgctc atttgggtgcc aggagccac gtggtaaaag catttaacac 600
catctcagcc tgggctctcc agtcaggagc actggatgca agtcggcagg tgtttgtgtg 660
tggaatgac agcaaagcca agcaaagagt gatggatatt gtctgtaatc ttggacttac 720
tccaatggat caaggatcac tcatggcagc caaagaaatt gaaaagtacc cctgcagct 780
atttccaatg tggaggttcc cttctattt gtctgctgtg ctgtgtgtct tcttgtttt 840
ctattgtgtt ataagagacg taatctaccc ttatgtttat gaaaagaaag ataatacatt 900
tcgtatggct atttccattc caaatcgtat ctttccaata acagcactta cactgcttgc 960
tttggtttac ctccctggtg ttattgtctg cattctacaa ctgtaccgag gcacaaaata 1020
ccgtcgattc ccagactggc ttgaccactg gatgctttgc cgaaagcagc ttggcttgggt 1080
agctctggga tttgccttcc ttcattgtct ctacacactt gtgattccta ttcgatatta 1140
tgtacgatgg agattgggaa acttaaccgt taccagggca atactcaaga aggagaatcc 1200
athtagcacc tctcagcct ggctcagtga ttcatatgtg gctttgggaa tacttgggtt 1260
ttttctgttt gtactcttgg gaatcacttc ttatgcatct gtttagcaatg cagtcaactg 1320
gagagagttc gattttgtcc agtccaaact gggttatttg accctgatct tgtgtacagc 1380
ccacaccctg gtgtacggtg ggaagagatt cctcagccct tcaaatctca gatggatatc 1440
tctgcagcc tacgtgttag ggcttatcat tccttgcaact gtgctgggtg tcaagtttgt 1500
cctaatactg ccattgttag acaacaccct tacaaggatc cgccagggtc gggaaaggaa 1560
ctcaaaacac tagaaaaagc attgaatgga aaatcaatat ttgaaacaaa gttcaattta 1620
gctggatttc tgaactatgg ttttgaatgt ttaaagaaga atgatgggta cagttaggaa 1680
agttttttcc tgaccctgt actgagggaa acattgcttg tctttgagaa attgactgac 1740
atactggaag agaacaccat tttatctcag gttagtgaag aatcagtgca ggtccctgac 1800
tcttattttc ccagaggcca tggagctgag attgagacta gccttgtggt tttcacac 1858

```

<210> 59
 <211> 1454
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1949083CB1

```

<400> 59
caccctgggtg tcttccctcg tggccctgca aggcataatca tgcttctctgt ctctaggacc 60
tgcttactag aaagcagcac ccgcctgaaa cctcacgaag ccagaaacta caggaagaag 120
gcattgtggg tgtcttggtt ctccatcatt gtcaccctgg ccctcgcggt ggctgccttt 180
actgtctccg ttatgaggta cagcgcctct gcttttgggt ttgcatttga tgccatcctg 240
gacgtctctg catcgcgcat tgtctctgtg cgttacagca acgcgccgcg tgtgcaactct 300
gcccataggg agtacatagc ctgtgtcatt ttgggggtga tattccttct gtcattccata 360
tgtatagtgg tcaaaagccat ccattgacctc tcaactaggc tgctccaga agtggacgat 420
ttcctgttca gtgtctccat tttaaagtgg attcttttga gcatcctggc cgtgttgaag 480
ttcatgctgg ggaaggttct gaccagtaga gcaactcataa cagatgggtt taactccctc 540
gtgggtggcg tgatgggctt ctccattctt ctgagcgcgg aagtgttcaa gcatgactcg 600
gcggtctggt acctggacgg cagcataggc gttctgatcg gctcaccat atttgcttat 660
ggggtcaaac tctcatcga catggtgccg aaggtgaggg agacacgtca ctacgagatg 720
tttgagtga gggggccagc atccgcatga gaccatcgag atgaggagtt cccacatagg 780
caaaggtgc caatatttaa ctgaacatct ggtttctttt tggaaagttt ctttcacatg 840
gtttgtcatt acaagacaag gtctgccag ccagggtggat ctaccttgcc cccatcacct 900
gccgccccca tcaaacatgt tgggacaatg cccataggaa tggacctcct tccccgtctc 960
cagctgggac tggcggtttt tagtctctgg agtatgatgg ttctcatggg taggatgaga 1020
tctttggcag aaaggtcttc ggtggtgctc tgagcctgag ctgcatagga ctgagcagac 1080
ccacctctc cagcttgggt ggccctgcca ctctgggtc caagtctctc ctttctctggc 1140
aggtcttaag ggaagattgt acccctcacc ctttacatac ccagaatcat cagtatgtca 1200
cttcttaatt tctatcagt tatctcatta tttcactact ttttactaat cctaagtcta 1260
aacagatttg ctcaaaagga gaccattcta ttttttaaag tacttagtga tacacgtata 1320
agctttgcat ggacgaatta aatagcaca ttgacctttt cttgtacatt cagaacctga 1380
acatccatgt gaaaactggg tccatttttg agagatgtga aactacagtt tatatgtaat 1440
aaataaatat aata

```

<210> 60
<211> 2310
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1996357CB1

<400> 60
ggcccgatg ttcgggtgcag ctgccagatc cgctgatcta gtgcttctcg aaaaaaacct 60
tcaggcggcc catggctgtc gatattcaac cagcatgcct tggactttat tgtgggaaga 120
ccctattatt taaaaatggc tcaactgaaa tatatggaga atgtggggta tgcccaagag 180
gacagagaac gaatgcacag aaatattgtc agccttgcac agaattcctt gaactttatg 240
attggctcta tcttggattt atggcaatgc ttcctctggt tttacattgg ttcttcattg 300
aatgggtactc ggggaaaaaag agttccagcg cacttttcca acacatcact gcattatttg 360
aatgcagcat ggcagctatt atcaccttac ttgtgagtga tccagttggt gttctttata 420
ttcgttcatg tgcgattatt atgctttctg actggtacac gatgctttac aaccaagtc 480
cagattacgt taccacagta cactgtactc atgaagccgt ctaccacta tataccattg 540
tatttatcta ttacgcattc tgcttggtat taatgatgct gctccgacct cttctggtga 600
agaagattgc atgtgggtta gggaaatctg atcgatttaa aagtatttat gctgcacttt 660
acttcttccc aattttaacc gtgcttcagg cagttggtgg aggccttcta tattacgctt 720
tcccatacat tatattagtg ttatctttgg ttactctggc tgtgtacatg tctgcttctg 780
aaatagagaa ctgctatgat cttctggtca gaaagaaaag acttattggt ctcttcagcc 840
actgggtact tcatgcctat ggaataatct ccatttccag agtggataaa cttgagcaag 900
atttgcctct tttggctttg gtacctacac cagccctttt ttactgttcc actgcaaaat 960
ttaccgaacc ttcaaggata ctctcagaag gagccaatgg acactgagtg tagacatgtg 1020
aaatgccaaa aacctgagaa gtgctcctaa taaaaaagta aatcaatctt aacagtgtat 1080
gagaactatt ctatcatata tgggaacaag attgtcagta tatcttaatg tttgggtttg 1140
tctttgtttt gtttatgggt agacttacag acttggaaaa tgcaaaactc tgtaatactc 1200
tggtacacag ggtaatatata tctgtacac tggaggcccg ctagggaagcc cttgcttctc 1260
tcaacagttc agctgttctt tagggcaaaa tcatgtttct gtgtacctag caatgtgttc 1320
ccattttatt aagaaaagct ttaacacgtg taatctgcag tccttaacag tggcgtaatt 1380
gtacgtacct gttgtgtttc agtttgtttt tcacctataa tgaattgtaa aaacaaacat 1440
acttgtgggg tctgtagtag aacatagaaa tgatgtatat tgttttttgt tatctattta 1500
ttttcatcaa tacagtattt tgatgtattg caaaaataga taataattta tataacaggt 1560
tttctgttta tagattgggt caagatttgt ttggattatt gttcctgtaa agaaaacaat 1620
aataaaaagc ttacctacat aaaatttcaa tgttttgaca cttaatgtgt gtttggcaca 1680
atagtatgga agtaattcaa actggttaaat agtttctctc catatctcgg gtatatatac 1740
ataccatatt ttattgatcc agagatactt atttcaactt gtgacatctc tgaattagga 1800
tgcattctac aactgatggc ttattaggtt taatgaaata cagaagatac acagtataaa 1860
aagggttttc ctgtgggttg tttgtggttt gtgatagggt ttctgtgatg tttatgcttt 1920
gaaggcctta agactcatgg ttgcaaccat ggaagcaaaa tgaaattttt agctcttaac 1980
ctaacaacct gaccatgttt atccattttt attgtttaga agtttattta ctgatacttg 2040
gtggagggtg tgtgaattag ttaaatttta aatgtttaag acttctatta acagctgcaa 2100
aatatgaaag taagtgcact cacttttccct gtatagtctt gtcttttgaa ttcacagcag 2160
ttgtatcctt gagttacttt gttaatgtat tttctcagc acatttaacc actgggaaat 2220
gaacccttgt acgaatgtgt ttcttcttct ctgtaggaat aaaaaataaa tataaaaaat 2280
ttatttgtat tgcacacaaa aaaaaaaaaa 2310

<210> 61
<211> 744
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2061330CB1

<400> 61
agtgggtgtg tacctgccga cagcataaag cgaggcaagg tccagccgtt ccgcccggca 60
tacaagctta tggagcagcc ccctttgaag atctccaggt ggacttcaca gagatgtcaa 120
agtgtagagg tgatcgagtg tggatcaaga actggaacgt agcctctttg tgtccactgt 180

```

ggaaaaggacc ccagactgtc gttctgagca ctcccaccgc tgtgaaggta gaaggaatcc 240
cagcctggat ccaccacagc catgtaaaac ctgcagcgcc tgaaacctgg gaggcaagac 300
caagcccaga caaccctgc agagtgaacc tgaagaagac gacaagccct gctccagtca 360
caccgcgaag ctgactgggc cacgcacggc cgaagcctga ggaagctcat catgagattc 420
atTTTTctta aattttggac ttatacagta agggcttcaa ctgatcttac tcaaactggg 480
gactgttccc agtgactca tcaggtcacc gaagtaggac agcaaattaa aacaatcttt 540
ctgtttcata gttattatga atgtatggaa acaataaaag aaacttgttt gtataatgcc 600
actcagtaca aggtatgtag cccgagaaat gaccgacctg atgtgtgtta taacccatct 660
gagccccctg caccaccgtt tttgaaataa gaataagaac tggccttttc ctagggtgata 720
caagtaaaat aataactaga acag                                     744

```

<210> 62
 <211> 1109
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2346947CB1

<220>
 <221> unsure
 <222> 30
 <223> a, t, c, g, or other

```

<400> 62
gaagcagtgc agagaggaga gcgagcgcn agtgccgctg agcaaaggcc ttcaccatgg 60
ccgagtcctcc cggtgctgc tccgtctggg cccgctgcct ccactgcctg tatagctgcc 120
actggaggaa atgccccaga gagaggatgc aaaccagcaa gtgcgactgt atctggtttg 180
gcctgctctt cctcaccttc ctcttttccc tgagctggct gtacatcggt ctgctccttc 240
tcaatgacct gcacaacttc aatgaattcc tcttcgccc ctggggacac tggatggact 300
ggtccttggc attcctgctg gtcatctctc tactgggtcac atatgcatcc ttgctattgg 360
tcctggccct gctcctcggt cttttagtag agccctgca tctgcacagc ctccacaagg 420
tgctgctgct cctcattatg ctgcttgtgg cggtggcct tgtgggactg gacatccaat 480
ggcagcagga gtggcatagc ttgctgtgtg cactgcaggc cacagcccca ttccttcata 540
ttggagcagc cgctggaatt gccctcctgg cctggcctgt ggctgatacc ttctaccgta 600
tccaccgaag aggtcccaag attctgctac tgctcctatt ttttggagtt gtcctggtca 660
tctacttggc cccctatgc atctcctcac cctgcatcat ggaaccaga gacttaccac 720
ccaagcctgg gctgggtggga caccgagggg cccccatgct ggctcccag aacaccctga 780
tgtccttgcg gaagacagct gaatgcggac tactgtgttt gagactgatg tgatggctag 840
ctccgatggg gtcccttccc tcacatgata tgagcacctc agcaggacca cgaatgtagc 900
ctctgtattc ccaaccgaa tcacagccca cagcagtgaac ttctcctgga ctgaactgaa 960
gagactcaat gctggatcct ggttcctaga gaggcgaccc ttctgggggg ccaaaccgct 1020
ggcaggccct gatcagaaag aggtgagag tcagtcggta ccagcattag aagagctatt 1080
ggaggaagct gcagccctca accctttca                                     1109

```

<210> 63
 <211> 2511
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2795577CB1

```

<400> 63
gcagccgtcg ccttcggagc gaaggggtacc gaccggcgag aagctcggag ctctcgggg 60
atcgaggagg caggcccgcg ggcgcacggg cgagcggggc gggagccgga gcggcgagg 120
agccggcagc agcggcgcgg cgggtccag gcgagggcgt cgacgctcct gaaaacttgc 180
gcgcgcgctc gcggcactgc gccggagcg atgaagatgg tcgcgccctg gacgcggttc 240
tactccaaca gctgctgctt gtgctgccat gtccgcaccc gcaccatcct gctcggcgtc 300
tggtatctga tcatcaatgc tgtggtactg ttgattttat tgagtgcctt ggctgatccg 360
gatcagtata acttttcaag ttctgaactg ggaggtgact ttgagttcat ggatgatgcc 420

```

```
aacatgtgca ttgccattgc gatttctctt ctcatgatcc tgatatgtgc tatggctact 480
tacggagcgt acaagcaacg cgcagcctgg atcatcccat tcttctgtta ccagatcttt 540
gactttgccc tgaacatggt gggtgcaatc actgtgctta tttatccaaa ctccattcag 600
gaatacatac ggcaactgcc tccataatct ccctacagag atgatgtcat gtcagtgaat 660
cctacctgtt tgggtccttat tattctctctg tttattagca ttatcttgac ttttaagggg 720
tacttgatta gctgtgtttg gaactgctac cgatacatca atggtaggaa ctctctctgat 780
gtcctggttt atgttaccag caatgacact acgggtgctgc taccctcgta tgatgatgcc 840
actgtgaatg gtgctgccaa ggagccaccg ccaccttacg tgtctgctta agccttcaag 900
tgggcggagc tgagggcagc agcttgactt tgcagacatc tgagcaatag ttctgttatt 960
tcacttttgc catgagcctc tctgagcttg tttgttgctg aaatgctact ttttaaaatt 1020
tagatgttag attgaaaact gtagttttca acatatgctt tgctggaaca ctgtgataga 1080
ttaactgtag aattcttctt gtacgattgg ggatataatg ggcttccacta accttcccta 1140
ggcattgaaa ctccccccaa atctgatgga cctagaagtc tgcttttcta cctgctgggc 1200
cccaaagtgg ggcatttttc tctctgttcc ctctcttttg aaaatgtaaa ataaaacca 1260
aaatagacaa ctttttcttc agccattcca gcatagagaa caaaacctta tggaaacagg 1320
aatgtcaatt gtgttaatcat tgttctaatt aggtaaatag aagtccttat gtatgtgtta 1380
caagaatttc ccccaacaac tcctttatga ctgaagtcca atgacagttt gtgtttgggtg 1440
gtaaaaggatt ttctccatgg cctgaattaa gaccattaga aagcaccagg ccgtgggagc 1500
agtgaccatc tgctgactgt tctgtgggat cttgtgtcca gggacatggg gtgacatgcc 1560
tcgtatgtgt tagaggggtg aatggatgtg tttggcgctg catgggatct ggtgcccctc 1620
ttctcctgga ttcacatccc caccaggggc ccgcttttac taagtgttct gccctagatt 1680
ggttcaagga ggtcatccaa ctgactttat caagtggaaat tgggatatat ttgatatact 1740
tctgcctaac aacatggaaa aggggttttct tttccctgca agctacatcc tactgctttg 1800
aacttccaag tatgtctagt caccttttaa aatgtaaaca ttttcagaaa aatgaggatt 1860
gccttctctg tatgcgcttt ttaccttgac tactgaatt gcaagggatt tttatatatt 1920
catatgttac aaagtcagca actctcctgt tgggtcatta ttgaatgtgc tgtaaattaa 1980
gtcgtttgca attaaaacaa ggtttgcccc caaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2040
aaaaaaaaaa aaaaaaaaaa caatacgaaa caacaagaac agcaagaatc aaaaagctta 2100
aaatcctcga gtgcacaatg aagaaaacag gggggcgccc ctctaggggt cccagtttctg 2160
gtcgggggtg ttgggggggac tgagcccttc ctatagggac cccctaattc tgatttcagg 2220
ggcggggcgg tttttaaaaa gcgcgggttaa cgggggaaac acccctgggg gttatcccc 2280
ctcttttaag cggcgtgttg gtgggaaaaga caccgcccct ttttgcgccc aagtgggggc 2340
atatactctc ccggtgaggg cgccctctac ccgatatctg gcctttcccc agaagttttg 2400
cccgcgccct ctactttggg gactttgggt gccgccacct ttattaaggg ggcgtttata 2460
ccagcggcgg gggtttgttg gcgtcctcgc gcgggaggcc cgctccaccc t 2511
```

<210> 64

<211> 788

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3255825CB1

<400> 64

```
cccacgcgtc cgcagctgcg tctctggggc cacgctccac ttgcccgtt ccaccggaa 60
agccccccag gctgagtgcg gcatgatctc catcacgaa ttggcagaaga ttggtgtggg 120
gatcacgggt ttcggcatct tcttcacctt ctttgaaca ctctgtact ttgattccgt 180
gtccttgccc tttggaaacc tgetgttctt gacgggcttg tccctcatca ttggcctgag 240
gaagaccttt tggttcttct tccaacggca caaactcaag ggaaccagct tctcctggg 300
gggtgtggtt atcgtgctcc tacgtggccc cctcctcggc atgttctctg aaacctacgg 360
attcttcagc ctctttaagg gctttttccc tgtcgcttcc ggttctctgg caatgtctgc 420
aacatccctt tcttgggtgc gctgttccgg agacttcaag gcactagctc gatggtctga 480
aaaacagaga tgagctcctt gaacttgatg cattgggtga gggggctaga gggagaatgg 540
gaaccacccc ctcagtcccc tgcactgact cactccccga catatccgga cctccccaa 600
tccagaagga aggaatggag ctgagcaact gacgtcaaat ccccaagtcg actcaagagg 660
ctgccaggaa gcagagatgc agacccaag gagactgggc tggggctggt atcacacct 720
cactctatat ttatgggagg aaaagtgaag attaaattcc caagttgtgc gtgtgtctaa 780
aaaaaaaa
```

<210> 65

<211> 1831

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3393430CB1

<400> 65

```
gccttataaaa gtagcctctg catctgcctg cctcgggcag aggaggggcta ccctgggggct 60
gagagttcac ctgtctcagg aaccacctga gccacacagat cctgtgggca ggggccaggg 120
cagccatggc ttgggcaagt aggctgggcc tgctgctggc actgctgctg ccggtggctg 180
gtgectccac gccaggcacc gtgggtccgac tcaacaaggc agcattgagc tacgtgtctg 240
aaattgggaa agcccccttc cagcggggccc tgcaggtcac tgtccctcat ttcttggaact 300
ggagtggaga ggcgcttcag cccaccagga tccggattct gaatgtccat gtgccccgcc 360
tccacctgaa attcattgct ggtttcggag tgcgcctgct ggcagcagct aattttactt 420
tcaaggtctt tcgcgcccc aagccctgg agctgacgct gcctgtggaa ctgctggctg 480
acaccgcgt gaccagagc tccatcagga cccctgtggt cagcatctct gcctgctctt 540
tattctcggg ccacgccaac gagtttgatg gcagtaacag cacctcccac gcgctgctgg 600
tcctggtgca gaagcacatt aaagctgtct tgagtaacaa gctgtgcctg agcatctcca 660
acctggtgca ggggtgtcaat gtccacctgg gcaccttaat tggcctcaac ccggtgggtc 720
ctgagtcacca gatccgctat tccatgggca gtgtgcccac tgtcaccagt gactacattt 780
ccctgggaagt caatgctgtt ctcttctctg tgggcaagcc catcatcctg cccacggatg 840
ccacccttt tgtgttgcca aggcattgtg gtaccgaggg ctccatggcc accgtggggc 900
tctcccagca gctgtttgac tctgcgctcc tgctgctgca gaaggccggt gccctcaacc 960
tggacatcac agggcagctg aggtcggatg acaacctgct gaacacctct gctctggggc 1020
ggctcatccc ggaggtggcc cgccagtttc ccgagcccat gcctgtggtg ctcaagggtg 1080
ggctgggtgc cacacctgtg gccatgctcc acacaaacaa cgccacctg cggtgcagc 1140
ccttcgtgga ggtcctggcc acagcctcca actcggcttt ccagtcctc ttctccctgg 1200
atgtggtagt gaacttgaga ctccagctct ctgtgtccaa ggtgaagctt caggggacca 1260
cgtctgtgct gggggatgtc cagctcacgg tggcctctc caacgtgggc ttcatgtata 1320
cagatcaggt gcgcacactg atgggcaccg tttttgagaa gccctgctg gaccatctca 1380
atgctctctt ggccatggga attgcctcc ctggtgtggt caacctccac tatgtcgccc 1440
ctgagatctt tgtctatgag ggctacgtgg tgatatccag tggactcttc taccagagct 1500
gaggcaagac cactgggagg cctgagagtg ggccagctcg ctgctcaggc gaatttctca 1560
tttcaagcca ctggggaaac tgaggcaaaa ccatacttag tcatcaccaa caagctggac 1620
tgcttagctg ggctgtttta tcttccctga gtgcctgggt ctccctccct cacttctgcc 1680
ctttcccttc ctctcctct tctcctccct ctccctccat ctccctccct ctctctctgc 1740
cccaccccag gggggagcag actgctcctc caggctgtat agacctgcc tcttgcatca 1800
aacaacttct cttgagctgc aaaaaaaaaa a 1831
```

<210> 66

<211> 1499

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3490990CB1

<400> 66

```
tgtttgagtt ggggtgaagaa ggcgggaagt taatcctctt ccaatcagct ccccatctcg 60
ccctctggcg gctccggacc acaggccgag ttactctgat taccgccatg ctccaagacc 120
cgcactgaca gtcccacact cccgttctct gtcacgggtt cctcfaatgg agtccacatg 180
tttgggcaga atctggaggt gcagctgagc tctgcgagga ccgagaacac gactgtggtg 240
tggaagagct tccatgacag catcacctc attgttctgt catctgaggt gggcatctct 300
gagctgaggc tggagagact actccaaatg gtgtttggag ccatggctct tcttgtggga 360
cttgaagaac tgaccaatat ccgcaacgtg gagagactga agaaggactt gagggccagt 420
tattgcctca tcgacagctt cctgggggac tcggagctca tcggggacct gaccagtggt 480
gtggactgcg tgattcctcc agagggtgct ctcttgagg aagccctctc cgggttcgct 540
gaggccgcgg gcacgacctt cgtcagctct gtggtgtccg gccgggtggt ggcagcaaca 600
gagggttggt ggcgggtggg gacgcccag gccgtgctgc tccctgggt ggtgggtgct 660
ctgccgcgcg agaccgctcg cgactaccog gtgtacctac cgacggggag cccacgggtc 720
ccacaccggt tcctgacctt gactctgctg ccgagcctgg agctgtgtct actctgcggg 780
```

```

ccgagccac ccctcagcca gttgtatcca cagcttctgg agcgtgtggtg gcagccactg 840
ctggaccogt tgcgggcctg tctgccgttg ggaccccggtg cgctgcccag tggcttcccc 900
cttcacacag acatcctcgg gctgctgctc ctccacctgg aactgaagcg ctgcctcttc 960
accgtggagc ccttggggga taaagagcct tcaccagaac agcgccggcg cctcctccga 1020
aactttctata ccctgggtcac ctccacgcac ttccaccag agccagggcc accagagaag 1080
acagaagatg aggtctacca ggcccagctg cccagagctt gctacctggt gttggggact 1140
gaggaaccag gcacaggagt gcgtctggtg gccttgacgc tggggcttcg gcggctgctg 1200
ctgctgctgt ctccccagag tccccccat gggctgcgaa gcctggccac ccacactctg 1260
catgccctca ccccaacttct ttgactacct agcagtgggt gatggacaca gacatggggc 1320
tgttagcgctc tctctgttta ttcgctcaca ataatacaca gcccctggat ggggaggggg 1380
taggaggggc tacaacaggg tgggggtggga ggggaggaga catccacttc cctggcccct 1440
ctccccctctg tgcttggggg ggaaagggag ggagggggac tccccctaac cccccagaa 1499

```

```

<210> 67
<211> 365
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 3635154CB1

```

```

<400> 67
gtttttttgc actaacttca ggagccagct cgtgatctca ggatgtatgg aaaaataatc 60
tttgtattac tattgtcaga aattgtgagc atatcagcat caagtaccac tgggtgtggca 120
atgcacactt caacctcttc ttcagtcaca aagagttaca tctcatcaca gacaaatgga 180
gaaacggggc aacttgtcca tcgtttcact gtaccagctc ctgtagtgat aatactcatt 240
attttgtgtg tgatggctgg tattattgga acgatcctct tattttctta cagttttcgc 300
cgactgataa agggctgagg gtgtagcctg catgtgccc atcttgctct gaaccggccg 360
cattg                                     365

```

```

<210> 68
<211> 1102
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 4374347CB1

```

```

<400> 68
gaggacagag caggcagcag agaccatggg gccccctca gcttgteccc acagagaatg 60
catcccctgg caggggctct tgetcacagc ctcaacttcta actttctgga acgcacccac 120
cactgcctgg ctctttattg catcagcgcc ctttgaagtt gctgaagggg agaattgtta 180
tctctctgtg gtttatctgc ccgagaatct ttacagctat ggctggtaca aagggaaaac 240
ggtggagccc aaccagctaa tcgcagcata tgtaatatag actcacgtta ggactccagg 300
gcctgcatac agcggctcgag agacaatatc acccagtggg gatctgcatt tccagaacgt 360
caccctagag gacacgggat actacaacct acaagtcaca tacagaaatt ctgagattga 420
acaggcatct caccatctcc gtgtatacga gtcagtggct cagccctcca tccaagccag 480
cagcaccaca gtcacagaga agggctccgt ggtcctgacc tgccacacaa ataacactgg 540
aacctctttc cagtggattt tcaacaacca gcgtctgcag gtcacgaaga ggatgaagct 600
gtcctggttt aacctgtgc tcaccataga ccccatcagg caggaggacg ctggggagta 660
tcagtgtgag gtctccaacc cagtcagctc caacaggagc gacccccctca agctgactgt 720
aaaatatgac aacactctag gcatactgat cggggctcctg gttgggagtc ttctgggtggc 780
tgcaattgtg tgtttcctgc tectccgaaa aactggcagg gccagcgatc agagtgactt 840
cagggagcag cagccccag cctccacccc cggccatgga cctctgaca gctctgacag 900
ctccatctcc taggaattgc tacactctga cacaaacatt tactgctgga tcgaccacaa 960
agcagatgtg gcttcttagg ttctctctgg agctgctcct gtgggttgat ggagcgtccc 1020
tgaagccccc agccctgggg atggggaagg acatggagcc tgagccagag aaccagctct 1080
gagtcctgag gagacacagg cc                                     1102

```

```

<210> 69
<211> 2546

```


<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4596747CB1

<400> 69
ccccggggcg ccccaccgcg ccgcatccat gttcgacacc acaccccaact ctggccggag 60
cacgccaagc agctccccc cgtcccgga acggctgcag ctccctgccc caagccggcc 120
cccacctgag ccagaaccag gcacatgggt ggagaaggga tcagatagct cctcagagaa 180
gggtgggggtg cctgggaccc ccagcaccca gagcctaggc agccggaact tcacccgcaa 240
cagaagaag atgcagagct ggtacagtat gctgagcccc acttataagc agcgtaatga 300
ggacttcgag aaactgttca gcaaactccc cgaagcagaa cgccctattg tggattactc 360
ctgcgccctg cagcgtgaga tcctgtcca tggggccctc tacctctctg agaactggat 420
ctgcttctac agcaacatct tccgctggga gaccacgac tccatccagc tgaagggaag 480
gacatgtctg aagaaggaaa agacggccaa gctgatcccc aacgccatcc agatctgcac 540
ggagagcgag aagcatttct tcaacttctt tggggccctg gaccgctgct tcctcctcat 600
cttccgcctc tggcagaatg cactgcttga aaagacgctg agtccccgag agctctggca 660
cctggtgcat cagtgtacg gctcagagct gggcctcacc agtgaggatg aggactatgt 720
ctcccccttg cagctgaacg gtctggggac ccccaaggaa gtgggagatg tgatcgccct 780
gagcgacatc acctcctcgg gggcagctga ccgagccag gagccaaagg cagtgggttc 840
gcgccgtggc catgtcacgc ccaaccttct ccgagccagc agcgacgcag accatggggc 900
agaggaggac aaggaggagc aggtagacag ccagccagac gcctcctcca gccagacagt 960
gaccccggtg gctgaacccc cgagcacaga gcccaccag cctgacgggc ccaccacct 1020
gggcccttg gatctgtgc ccagtggga gctattgaca gacacaagta actcctcttc 1080
atccactggg gaggaagcgg acttggctgc cctgcttccc gacctctcgg gccgcctct 1140
catcaactct gtcttccatg tgggcgctga cgggctccag cagatgctct tctcggactc 1200
gcccttctc cagggcttcc tacagcagtg caagttcaca gacgtgacgc tgagccctg 1260
gagtggggac agcaagtgcc accagcgccg ggtgctgacg tacaccatcc ccatcagcaa 1320
cccactgggc cccaagagcg cctccgtggg ggagacacag acgctgttcc ggcgcggccc 1380
ccaggccggc ggggtgtgtg tggactccga ggtgctgacg caggcatcc cctaccagga 1440
ctacttctac actgcccacc cctactgcat cctgggtctg gcccggaaca aggcgcggct 1500
ccgagtgtct tctgagatcc gctaccgaaa gcagccgtgg agcctggtga agtcgctcat 1560
tgagaagaac tcgtggagcg gcattgaaga ctatttccac catctggagc gagagctcg 1620
caaggctgag aagctgtctc tggaggaagg cgggaaggat gcccggggct tgctatccgg 1680
cctgcggcgg cggaagcgcc ccctgagctg gcgggctcac ggggacgggc cccagcacc 1740
agatcctgac ccctgtgccc gggccggcat tcacacctcg ggctccctca gctcccgctt 1800
ctccgaacca tctgtggacc agggcccgcc ggcaggcatc ccagtgccc tggttctcat 1860
cagcattgtg atctgtgtga gccttateat cctcatcgcc ctcaacgtcc tgctcttcta 1920
ccgctctg tccctgaaa ggacagccca cacttttgag tcctggcaca gcctggccct 1980
ggccaagggc aagttcccc agacggccac agagtgggag gagatcctgg cgctgcagaa 2040
gcaattccac agcgtggagg tgcacaagtg gaggcagatc ctgcccggct ccgtggagct 2100
cctggatgag atgaagttct cgctggagaa gctgcaccaa ggcatcacag tctcagaccc 2160
tccctttgac acccagcccc ggcccgatga cagcttttcc tgaggacccc ggccacgcag 2220
ctgttcccc acatggacag atggacacac agagcctcgg cggccactgc tggcacggtg 2280
tgagcgccag gcattcccc cccgccccct ccgacggccc aaccaggggc tgtgcagacg 2340
tggggaccac ggaaccgaga tgcactttag accagggagc tggcccggcc tctggcaggc 2400
ccccactaa cttattttgc ccggtgagg ttgtgggggg cgccctctgg ggtgcacgat 2460
tccctcagct ctgggtttaa tgtattatat ttatttgggg ccgacagtgc cccaataaag 2520
ggtcagaagt gaaaaataaa aaaaaa 2546

<210> 70
<211> 1845
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5052680CB1

<400> 70
ctctggcgga ctgcctggcg gaagcgggaa cgtcgcatcc tgaggtaaag gtgcacggca 60

tcctgggaca	tgtagtctgg	ccggggctcg	gacgccccct	cggatgaatg	ggaccgaagc	120
tgactgcgaa	ctacagcttc	ttggcagcgt	cgggtgtggc	cgcgggagaa	ggggagaccg	180
cggcgcccc	cagtgaagac	ggctttccag	gacggtgcga	tgtgctgcgc	agcgaagagg	240
caggaggccg	gcttcctggg	gtagcggtag	aggcggggcg	ttactctgtg	cgcttgcttc	300
cccaaccctg	caccggccat	gcgcccggcc	ttggcgggtg	gcctgggtgt	cgcaggctgc	360
tgcagtaacg	tgatcttcc	agagctcctg	gcccgggaagc	atccaggatg	tgggaacatt	420
gtgacatttg	cacaattttt	atttattgct	gtggaaggct	tcctctttga	agctgatttg	480
ggaaggaagc	caccagctat	cccaataagg	tactatgcc	taatggtag	catgttcttc	540
accgtgagcg	tggatgaaca	ctatgccctg	aatctcaaca	ttgccatgcc	cctgcatatg	600
atatttagat	cgggttctct	aattgccaac	atgattctag	gaattatcat	tttgaagaaa	660
agatacagta	tattcaata	tacctccatt	gccctgggtg	ctgtggggat	atttatttgc	720
acttttatgt	cagcaaaagc	ggtgacttcc	cagtcagct	tgaagtga	tgatggattc	780
caggcatttg	tgtgtgtgtt	actaggtatt	ggggcattga	cttttgcct	tctgatgtca	840
gcaaggatgg	ggatattcca	agagactctc	tacaaacgat	ttgggaaaca	ctccaaggag	900
gctttgtttt	ataatcacgc	ccttccactt	ccgggtttcg	tcttcttggc	ttctgatatt	960
tatgaccatg	cagttctatt	caataagtct	gagttatatg	aaattcccgt	catcggaagt	1020
accctgccca	tcattgtgtt	ctacctcctc	atgaacatca	tcactcagta	cgtgtgcatc	1080
cggggtgtgt	ttatcctcac	cacggaatgc	gcctccctca	ccgtcacgct	cgtcgtgacc	1140
ctacgcaaat	ttgtgagcct	catcttttcc	atcttgtagt	tccagaaccc	cttccaccctg	1200
tggcactggc	tgggcacctt	gtttgtcttc	attgggacct	taagtacac	agaggtgtgg	1260
aacaacctag	ggaccacaaa	aagtgaagct	cagaaggaca	gcaagaagaa	ctgaggcctg	1320
tctggagtac	gtagaccagt	gtcgtcgtga	gggtgggacc	ctgtgaaggt	ctgaccaccg	1380
tttcgctttt	gttaatgcgc	agctaccgcg	agtgcctgagc	cagccgtgca	aaaggaaatc	1440
ttcaggaggg	gacttctcac	gttgctcaga	ctgacacatg	tagactaaat	agaaaccctc	1500
cagccctaaa	atagaaaaaa	gaacaggtct	agactattga	acaggcgatt	catttatttc	1560
tgttttgttt	accgacatat	tcagtattat	tcttgttttc	tgaattctga	gtctcctgaa	1620
caaaaggctt	actatccatg	gtcttggaaa	agattgtccc	ttctcttgct	gttgtaatgc	1680
atctctgcta	cagcagtcac	tgtcatcctg	tgaccttacc	ccttttgtct	cccatcacct	1740
ttctcttctt	cagcagcacc	tcacagtcag	tcagacgctg	caggcaccac	tatcaccagg	1800
agtggccaag	taaaccgaga	caccagtaac	agcagggtgg	acttc		1845

<210> 71

<211> 1940

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5373575CB1

<400> 71

gtctcctgtc	aaaagccatg	ctcggcaggt	ctgggtaccg	ggcgtgccc	ctgggtgatt	60
ttgaccgctt	ccagcagtcg	agcttcggct	ttctgggctc	gcagaagggc	tgcttgctcc	120
cggagcgggg	cggcgtgggg	acaggggccc	atgtacccca	gagctggccc	tcctgcctct	180
gtcatggcct	catcagtttc	ctggggttct	tgtgctgtgt	ggtcaccttc	cccatttctg	240
gctggtttgc	cctgaagatt	gtgcccacct	acgagcggat	gattgtgttc	cgctggggcc	300
ggatccgcac	ccccaggga	ctggcatgg	ttctgctctt	gcccttcatt	gactccttcc	360
agagggtgga	tctgaggaca	cgagccttca	acgtccctcc	ctgcaagctg	gcctctaagg	420
acggggctgt	gctgtccgtg	ggagccgatg	tccagtttgc	catctgggac	ccgggtgctgt	480
cgggtgatgac	tgtgaaagac	ctgaacacag	ccacacgcat	gacagcccag	aacgccatga	540
ccaaggccct	gctcaagagg	ccgctgcggg	agatccagat	ggagaagctc	aagatcagcg	600
accagcttct	gctggagatc	aacgatgtga	ccagggcctg	ggggctggag	gtagaccgcg	660
tggagctggc	agtggaggcc	gtgctccagc	cgccccagga	cagcccagct	gggcccaccc	720
tggacagcac	ctgccagcag	ctggccctgc	acttctggg	aggaagcatg	aactcaatgg	780
caggaggtgc	cccgtccccg	gggcccagcag	acaccgtgga	gatggtgagt	gaagttgagc	840
cacctgcccc	tcaagttggt	gccaggtcca	gtccgaagca	gcctctggcg	gaggggctac	900
tgactgtctct	acagcccttc	ctgtctgagg	ccctggtcag	ccaagtcggg	gcctgctacc	960
agttcaatgt	cgctcctgcc	agcggcaccc	aaagcgccca	cttctctggac	ctcactacag	1020
gacgaggaag	agtgggacac	gggggtgcctg	atggcatccc	tgatgtgggtg	gtggagatgg	1080
ccgaggcaga	cctgcggggc	ctgctatgca	gagagctggg	gccccggggg	gcctacatga	1140
gtggacggct	gaaggtgaag	ggcgacctgg	ctatggccat	gaagctggag	gctgtcctca	1200
gggccttgaa	gtagcagcct	tggctgactt	tccagagccc	agtcccaagc	ctggcaccaa	1260
gccccagggg	cctcttgagg	gaggaggtgt	tcactgtcac	cacagagagt	tgaggcccta	1320

```

acaaatttca ggcccagcca agagcccatg aatggaggct gcaggaggct gagtccggct 1380
gccatgcacg tctcccttac agtggttctc tggacaaggc tttgtccatc ccggtcccca 1440
gctgagtgcc cagcgctgag ctgggtgcac ggtgtgattc caggaggaga gccaggcctg 1500
ccctgccctg ctggcttctc gactggagag acaggaccca cagaaacagc ctgacagcag 1560
ctggttttgt ccttgtgtga gggaccaagc atgtggccca ggctctaagc tctgcgggga 1620
ttggagaggg atggggaggg aagggaaggc agctccaaga agaggctcct gtggcgaagt 1680
tacctgggga tcctggctgg ccaccttcc tggctgcagt ccaggcccgt gctggcggga 1740
ttgggcattg gaaggagcag ggctgctgc ttcctggcg ctgctcccaa agatttctga 1800
ctcatctgcc agctctgtcc tgcattgctg gcgagctggg gccaggggca gcatgaagga 1860
gagccctgcg ttctgtgctt cttaccagag gtttgcaagc ctcagacaaa taaatgtggg 1920
gtttacaatg taaaaaaaaa 1940

```

<210> 72

<211> 880

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5524468CB1

<400> 72

```

gcgaagccga agggagcgcg gctaagagtg ccgcaccgcc tcacaacctg ggaaccggag 60
agttaggggc gtgggctggc aagaaccgc cgtgcctcct cggcaagggc catccgggtg 120
caccatgtc gactagagc agaagagggt gagtcctgga actgcaacct gcacagagct 180
gctctgtact gtccctgggt gtgcgcgcca tgacctggtt ggtgctgctg gggacactgc 240
tctgcatgct gcgcgttggg ttaggcaccc cggactccga gggtttcccg ccccgctgcg 300
tcacaactg cccctacaaa tgtatctgcg ctgcccacct gctaagctgc actggcctag 360
ggctgcagga cgtgccagcc gagttacctg ccgctactgc ggacctcgac ctgagccaca 420
acgcgctcca gcgcctgcgc ccgggctggt tggcgccctt ctccagctg cgcgccctgc 480
acctagacca caacgaacta gatgcgctgg gtgcggcgct ctctgtcaac gccagcgggc 540
tgaggctgct cgatctatca tctaaccgt tgcgggcgct tgcccgccac gacctcgacg 600
ggctgggggc gctgggagaag ctgcttctgt tcaataaccg cttggtgcac ttggacgagc 660
atgccttcca cggcctgcgc gcgctcagcc atctctacct gggctgcaac gaactcgct 720
cgttctcctt cgaccacctg caggtctga gcgccacca cctgcttact ctggacctct 780
cctccaaccg gctgggacac atctccgtac ctgagctggc cgcgctgccg gccttctcct 840
agaacggcct ctacctgcac gacaacacat aaaaaaaaaa 880

```

<210> 73

<211> 2403

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5944279CB1

<400> 73

```

agccacctgc tggccaggta cccctccctt acctggggca gtgtctgcct ggtggccact 60
agagacagcc cagcctgggc catggaagaa aaccgcacct tggaatcaga agcctggggc 120
tcctctaggg ggtggctggc ccccccggag gccagaggag gccatcgct gtcttctgtg 180
ctgaacgagc tgcccagtcg tgccaccctt cggtagcgag accctggggt gctgccttgg 240
ggggcgctgg aggaggagga ggaggatgga ggaaggagca gaaaggcctt cacagaagtc 300
acccagacag agctgcagga cctcaccct tcccgggaac tgccctggcc catgcaggcc 360
agacggggcac acaggcaaag aaatgccagc agggaccagg tgggtctatg ctctggaaact 420
aagacggacc gatgggcgcg gctacttcgg aggtccaagg agaaaacaaa ggaaggcttg 480
cgaagcctgc agccctgggc gtggacactg aagaggatcg ggggccagtt tggcgccggc 540
acggagtctt acttctcctt gctgcgcttc ctgtctcttc ttaacgtgct ggcctctgtg 600
ctcatggcct gcatgacgct gctgccacc tggttgggag gcgctcccc aggcctctcc 660
ggccccgaca tctctcgcc ctgcggctcc tataaccccc actcccagg cctggctacc 720
tttgccacc agctcttcaa cttgctctcg ggtgagggtt acctggaatg gtccctctc 780
ttctatggct tctaccgcc ccgcccacgc ctggcggtca cctacctgtg ctgggccttt 840
gccgttggcc tcattctgct cctgctcatc ctgcatcgct cgggtgtctg gctgaagcag 900

```

```

acactgctgg cggagtcgga ggctctgacc agctacagcc accgggtggt ctcggcctgg 960
gacttcggtc tctgcgggga cgtccacgtg cggctgcgcg agcgcatcat cttgtacgaa 1020
ttaaaggtgg agctggagga gacagtgggt cggcgccagg ctgcggtgcg gacgctgggc 1080
cagcaagcca gggtttggtt ggtgcgggtg ctgctcaacc tgctgggtgg cgcgctcctg 1140
ggggcagcct tctatggcgt ctactgggct acggggtgca ccgtggagct gcaggagatg 1200
ccccctgtcc aggagtggcc actgctgaag cttgggggtga attacctcc gtccatcttc 1260
atcgctgggg tcaattttgt gctgcccggc gtgttcaagc tcattgtctc actggagggc 1320
tacactcgga gtcgccagat cgtttttatc ctgctcagga ccgtgtttct tcgectcgcc 1380
tccctggtgg tccctgctct ctctctctgg aatcagatca cttgtggggg cgactccgag 1440
gctgaggact gcaaaaacct tggctacaat tacaacaac ttccgtgctg ggagactgtc 1500
ctgggccagg aaatgtacaa acttctgctc tttgatctgc tgactgtctt ggcagtcgag 1560
ctgctcatcc agtttcttag aaagctcttc tgtggcctct gtctggggg gctgggtcgt 1620
ctggcgggga cccaggagtt ccagggtccc gacgaggtgc tggggctcat ctacgcgcag 1680
acgggtggtc ggggtggggg ttttttctgc cctttactgc ccctgcttaa cacggtcaag 1740
tccctgctgc ttttctacct gaagaagctt accctcttct ccacctgtc cccggctgcc 1800
cgcaccttcc gggcctccgc ggcaaatctc ttttccctc tggctctct cctgggtctg 1860
gccatctcca gcgttccctt gctttacagc atcttctga tcccgcctc taagctttgt 1920
gggtccattcc gggggcagtc gtccatctgg gccacagatc ctgagtctat ttccagctc 1980
cctgagacca cccagaattt cctcttcttc ctggggaccc aggttttgc tgtgccccct 2040
ctgctgatct ccagcatcct gatggcgtag actgtggctc tggctaactc ctacggacgc 2100
ctcatctctg agctcaaacg tcagagacag acggagcgcg agaataaagt ctctctggca 2160
cggcgcgctg tggcgctgac ctccacaaaa cgggtcttct gacccccgca gcccacgtcc 2220
cgctttcaga ccccgagccc attgtaagcc taggtcaca catctgtaa ctaggagAAC 2280
tggagaagac tccacgcctt tccagctttg gtatctggag atttccaggg cccctcgccg 2340
ccacgtccct gactctcggt tgatcttctc tgtatcaata aatacagccg aggttgggtg 2400
gcc 2403

```

<210> 74
<211> 2850
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 6114480CB1

```

<400> 74
cggctcgagc ggctcgagtg aagagcctct ccacggctcc tgcgcctgag acagctggcc 60
tgacctccaa atcatccatc caccctgct gtcactctgt ttcatagtgt gagatcaacc 120
cacaggaata tccatggctt ttgtgctcat tttggttctc agtttctacg agctggtgtc 180
aggacagtgg caagtcaact gaccgggcaa gtttgtccag gccttgggtg gggaggacgc 240
cgtgttctcc tgctccctct ttccctgagac cagtgcagag gctatggaag tgcggttctt 300
caggaatcag ttccatgctg tggctccact ctacagagat ggggaagact gggaaatctaa 360
gcagatgcca cagtatcgag ggagaactga gtttgtgaag gactccattg cagggggggc 420
tgtctctcta aggtctaaaa acatcactcc ctgggacatc ggctgtatg ggtgctggtt 480
cagttccacg atttacgatg aggaggccac ctgggagctg cgggtggcag cactgggctc 540
acttcccttc atttccatcg tgggatatgt tgacggaggt atccagttac tctgcctgtc 600
ctcaggctgg ttccccagc ccacagccaa gtggaaaggt ccacaaggac aggtattgtc 660
ttcagactcc agagcaaatg cagatgggta cagcctgtat gatgtggaga tctccattat 720
agtccaggaa aatgctggga gcatattgtg ttccatccac cttgctgagc agagtcata 780
ggtggaatcc aaggtattga taggagagac gtttttccag ccctcacctt ggcgcctggc 840
ttctatttta ctggtgttac tctgtggtgc cctgtgtggt gttgtcatgg ggatgataat 900
tggtttcttc aaatccaaag ggaaaatcca ggcggaactg gactggagaa gaaagcacgg 960
acaggcagaa ttgagagacg cccggaaaca cgagtgagag gtgactctgg atccagagac 1020
ggctcaccgc aagctctgcg tttctgatct gaaaactgta acccatagaa aagctcccca 1080
ggaggtgcct cactctgaga agagatttac aaggaagagt gtggtggctt ctcagggttt 1140
ccaagcaggg agacattact gggagggtga cgtgggacaa aatgtagggt ggtatgtggg 1200
agtgtgtcgg gatgacgtag acagggggaa gaacaatgtg actttgtctc ccaacaatgg 1260
gtattgggtc ctcagactga caacagaaca tttgtatttc acattcaatc cccattttat 1320
cagctcccc cccagcacc ctcctacacg agtaggggtc ttctggact atgagggtgg 1380
gacctctcc ttcttcaata caaatgacca gtcccttatt tataccctgc tgacatgtca 1440
gtttgaaggc ttgttgagac cctatatcca gcatgcgatg tatgacgagg aaaaggggac 1500
tcccatattc atatgtccag tgtcctgggg atgagacaga gaagaccctg cttaaagggc 1560

```

cccacaccac	agaccagac	acagccaagg	gagagtgttc	ccgacaggtg	gccccagctt	1620
cctctccgga	gcctgcgcac	agagagtcac	gccccccact	ctcctttagg	gagctgaggt	1680
tcttctgccc	tgagccctgc	agcagcggca	gtcacagctt	ccagatgagg	ggggattggc	1740
ctgaccctgt	gggagtcaga	agccatggct	gccctgaagt	ggggacggaa	tagactcaca	1800
ttaggttttag	tttgtgaaaa	ctccatccag	ctaagcgatc	ttgaacaagt	cacaacctcc	1860
caggctcctc	atttgctagt	cacggacagt	gattcctgcc	tcacaggtga	agattaaaga	1920
gacaacgaat	gtgaatcatg	cttgacaggtt	tgagggcaca	gtgtttgcta	atgatgtgtt	1980
tttatattat	acattttccc	accataaact	ctgtttgctt	attccacatt	aattttactt	2040
tctctatacc	aaatcaccca	tggaaatagtt	attgaacacc	tgctttgtga	ggctcaaaga	2100
ataaagagga	ggtaggattt	ttcactgatt	ctataagccc	agcattacct	gatacaaaa	2160
ccaggcaaag	aaaacagaag	aagaggaagg	aaaactacag	gtccatatcc	ctcattaaca	2220
cagacacaaa	aattctaaat	aaaattttaa	caaattaaac	taaacaatat	atttaaagat	2280
gatataatac	tactcagtgt	ggttttgtccc	acaaatgcag	agttggttta	atatttaa	2340
atcaaccagt	gtaattcagc	acattaatag	agtaggaaa	aaaccatagg	aaggagagac	2400
gaaggaaa	gaaaaggggg	cgccccaact	agtgcgtgg	caccccgggg	attttcgggc	2460
goggactctc	aggggggtag	cgggtatcga	agcagctttg	aagacggaga	ctcggggggg	2520
gccgggacca	aattcccca	gaggggcggg	tatcggcgcg	cgctggcctc	ggtgttaa	2580
gccggagtgg	gaaccaggg	gtaccaagtt	agggcttgag	ccgaccgtat	gttgggcttc	2640
cagctagaga	ctgcaaggaa	agaagaagga	ggctagaggc	caggatagtg	gagagagaac	2700
agacgcgaga	acaaggggaa	atgaggacgc	gaggggagag	atgaggagag	atagagatgt	2760
gacgagcagc	gaggagagcc	gggaccggag	aggcaacgga	gaagtgacga	acgaagaa	2820
acaagggagc	gaaggaacgc	agaagtggaa				2850

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.